



The CCB Science 2 Service Distance Learning Program[®]

Please complete all required information and fax to 203.284.9500

For questions or assistance with the program, please email Jeff at JQuamme@ctcertboard.org

S2S Course 2042 TIP 49 Incorporating Alcohol Pharmacotherapies into Medical Practice Module 2 PRE TEST

1. Drinking alcohol enhances _____ opioid activity.
 - a) respiratory
 - b) endogenous
 - c) androgynous
 - d) heterogenous
2. Short term naltrexone treatment is linked to a:
 - a) lower percentage of drinking days
 - b) fewer drinks per drinking day
 - c) more days of abstinence
 - d) all of the above
3. Studies have shown that the use of naltrexone for adolescents with AUDs:
 - a) is safe and effective
 - b) is contraindicated
 - c) needs additional research
 - d) is widely recommended
4. Pharmacological interventions, such as naltrexone, in and of themselves, do not constitute comprehensive treatment.
 - a) True
 - b) False
5. Injectable naltrexone plasma levels tend to peak at two times, approximately 2 hours after injection and then _____ later.
 - a) 12-24 hours
 - b) 48-72 hours
 - c) 5 days
 - d) 7 days
6. According to TIP 49, _____, identifying and treating patients with AUDs has the potential to improve the healthcare of many primary care patients.
 - a) screening
 - b) encouraging
 - c) assessing
 - d) medicating
7. Biomarker/s which may reflect the presence of an AUD is/are :
 - a) CDT
 - b) AST
 - c) GGT
 - d) All of the above



The CCB Science 2 Service Distance Learning Program[®]

Please complete all required information and fax to 203.284.9500

For questions or assistance with the program, please email Jeff at JQuamme@ctcertboard.org

8. Any pharmacological treatment for alcohol dependence should be used _____ psychosocial treatment.

- a) as a replacement for
- b) as an adjunct to
- c) after completion of
- d) both a and c

9. Although some AA members may have negative attitudes toward medications, the organization itself supports appropriate medication use.

T F

10. Assessing a patient's response to medication includes the _____.

- a) efficacy of the medication
- b) cost of the medication
- c) side effects of the medication
- d) a and c only