

**RESTRICTED USE PESTICIDE
DUE TO HIGH ACUTE TOXICITY TO HUMANS**

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading or repair or cleaning of application equipment.

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

ACCEPTED

July 24, 2013



Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

DUPONT™ LANNATE® SP INSECTICIDE

EPA Reg. No. 352-342

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc id: 535933

**FOR CONTROL OF SPOTTED WING DROSOPHILA
ON BLUEBERRIES IN THE STATE OF NEW YORK**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DuPont™ LANNATE® SP insecticide is recommended for the control of spotted wing drosophila (SWD) in blueberries. Apply LANNATE® SP at 1/2 to 1 pound/acre when the first adult SWD are trapped locally or based on local university crop stage timing recommendations in order to protect ripening fruit.

APPLICATION

Apply by properly calibrated air or ground equipment using sufficient spray volume to obtain thorough coverage. For aerial application in dense growth blueberries, use a spray volume of 10 gallons per acre. Make repeat applications as needed based on scouting or on a 5 to 7 day schedule.

Note:

Do not apply during bloom.

Do not apply more than 4 pounds per acre per crop.

Do not make more than 4 applications per crop.

The field re-entry interval following application of LANNATE® SP is 48 hours.

The pre-harvest interval after the last application of LANNATE® SP is 3 days.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use. If the terms are not acceptable, return the product at once, unopened, for a refund of the purchase price. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2013 E.I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898