

ADM Animal Nutrition™
Gro Fast® 36 Calf Concentrate
RU 35245NKS

▶ **MEDICATED** ◀

For Growing Calves

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in calves (excluding veal calves.)

ACTIVE DRUG INGREDIENTS

Monensin..... (57.15 mg/lb)..... 114.3 gm/ton

GUARANTEED ANALYSIS

Crude Protein, Min. 36.0%
Crude Fat, Min. 2.00%
Crude Fiber, Max. 8.00%
Calcium (Ca), Min. 2.60% Max. 3.10%
Phosphorus (P), Min. 0.55%
Potassium (K), Min. 1.0%
Vitamin A, Min. 16,000 IU per pound
Vitamin D₃, Min. 3,800 IU per pound
Vitamin E, Min. 180 IU per pound

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Molasses Products, Animal Fat preserved with BHA, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Calcium Carbonate, Salt, Magnesium Oxide, Manganous Oxide, Manganese Sulfate, Magnesium Potassium Sulfate, Potassium Sulfate, Calcium Phosphate, Ferrous Sulfate, Cobalt Carbonate, Copper Sulfate, Calcium Iodate, Zinc Oxide, Zinc Sulfate, Sodium Propionate (Preservative) and Sodium Selenite.

FEEDING DIRECTIONS

Hand feed 2.0 lb of Gro Fast® 36 Calf Concentrate per head per day to calves weighing from 115 to 350 lb. This will provide from 0.99 to 0.33 milligrams monensin per pound of body weight per day. A maximum of 3.5 lb per head per day of this product may be fed (to provide up to 200 milligrams per head per day.) Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be a hazard. Allow free access to pasture or roughages. Provide clean, fresh water at all times.

CAUTION: Consumption of this product by sheep and goats may result in copper toxicity. Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding high concentrations of monensin could be fatal to cattle. Do not exceed the levels of monensin recommended in the feeding directions, as reduced daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. 11517

ADM Animal Nutrition, a division of

Archer Daniels Midland Company, Quincy, IL 62305-3115

Net Weight 50 lb (22.67 kg) Or As Invoiced