

COLON AND RECTAL SURGERY, LTD.

KENNETH J. BOYD, M.D. • SAMIR N. PARIKH, M.D. • SHIRLEY S. SHIH, M.D.

610-565-3435

Colonoscopy at _____ Date _____ @ _____ Arrival Time _____

MIRALAX AND DULCOLAX BOWEL PREP FOR COLONOSCOPY

A bowel prep is done to clear the bowel of all solid matter. Its purpose is to prepare the bowel for surgery or a procedure. Please follow these instructions. Begin the bowel prep one day before your scheduled procedure.

TO PREPARE:

- Tell your doctor if you have diabetes or take blood thinners. You may need to stop some of your medicine a few days before your test.
- Ask your doctor if you should take any of your medicines the morning of your test. If so, take with a sip of water only.
- One week before your test:
 - Do not take aspirin products or iron tablets.
 - Do not take fiber supplements like Metamucil, Citrucel or Fiberall.
 - Do not eat popcorn or any corn.
- *You will need these items from the pharmacy:*
 - **Miralax** in large bottle (8.3 ounces or 238 grams)
 - **4 Dulcolax** or bisacodyl tablets (5 milligram tablets)

There may be store brands of these products that cost less. Ask the pharmacist to help you find what you need.

- Buy a large 64-ounce bottle of sports drink such as Gatorade. **If you have diabetes**, buy a no or low calorie drink such as Crystal Light instead. This will be used to mix your Miralax the day before your test. Do not use carbonated beverages.
- Arrange to have an adult bring you to your appointment and take you home after your test.

ON THE DAY BEFORE YOUR TEST

1. Drink only clear liquids. **Avoid all red or purple colored liquids.** Do not eat any solid food or milk products until your test is done. Clear liquids include:
 - Water, Popsicles, Jell-O, Ice, Kool Aid, Gatorade, Coffee or tea (no milk or creamer),
 - Strained fruit juices (no pulp), Clear broth or bouillon
2. Follow the schedule below for your bowel prep. You may need to go to the bathroom right away. You will have many bowel movements through the day. They will become very watery. The bowels are clear and clean when there is only pale yellow fluid without flecks of stool.

1:00 PM	Take 4 Dulcolax tablets with a drink of clear liquids.
2:00 PM	Mix the Miralax in a 64-ounce bottle of Gatorade or other clear liquid of choice. Cap the bottle and shake to dissolve the powder. Most people prefer to drink the liquid chilled so you may want to place it in the refrigerator.
4:00 PM	Start to drink the Miralax. Drink it quickly rather than sipping small amounts because it does not taste that good. Finish drinking the liquid in 2 hours. Be sure to drink all of the liquid.
12:00 Midnight	Do not eat or drink anything after midnight. You may gargle but do not swallow any liquid. Do not smoke after 12 midnight.

THE MORNING OF THE COLONOSCOPY

- Do not eat or drink anything until after your test is done.
- If you are to take your morning medicines, take with a small sip of water only.
- Bring all medicines you usually take (in the original containers) to the hospital with you.
- You will need to have an adult with you to take you home after your test. You will not be able to leave by yourself and it will not be safe for you to drive because of the medicine you are given at the start of the test.
- You may be told to arrive 1 hour before your test is scheduled so that staff can get you ready. Expect to be here for 2 to 4 hours for your test and recovery time.

Talk to your doctor or nurse if you have any questions or concerns.