

# Fixed Point Theorem for Contraction Mapping Using Iteration Methods

Vandana Garg<sup>1</sup>, Chetan<sup>2</sup>

*Department of Mathematics, Dravidian University, Kuppam, (A. P)*

**Abstract:** Object in this paper to discuss about fixed point theory in metric spaces, also we established some fixed point theorems in complete metric spaces. Many problems in pure and applied mathematics have as their solutions the fixed point of some mapping. Therefore a number of procedures in numerical analysis and approximations theory amount to obtaining successive approximations to the fixed point of an approximate mapping

**Keywords:** Banach Space, Complete metric space.

## I. INTRODUCTION

Many problems in pure and applied mathematics have as their solutions the fixed point of some mapping. Therefore a number of procedures in numerical analysis and approximations theory amount to obtaining successive approximations to the fixed point of an approximate mapping. Our object in this paper to discuss about fixed point theory in metric spaces, also we established some fixed point theorems in complete metric spaces.

## II. BANACH FIXED POINT THEOREM

The Banach fixed point theorem (also known as the contraction mapping theorem or contraction mapping Principle.) is an important tool in the theory of metric spaces; it guarantees the existence and uniqueness of fixed points of certain self maps of Metric space, and provides a constructive method to find those fixed points.

### A. Contraction mapping

B. Let  $(X, d)$  be a Metric space. A mapping  $f : X \rightarrow X$  is called a contraction mapping if there is a real number  $k$ ,  $0 < k < 1$ , such that

Proof: Let  $x_0$  be an arbitrary point of  $X$  and let  $\{x_n\}$  where  $x_n = f^n(x_0)$  and  $n$  is a positive integer, be the sequence of iterates of  $f$  at  $x_0$ . If  $x_n = x_{n+1}$  for some  $n$  then the result is immediate. So let  $x_n \neq x_{n+1}$  for all  $n$ .

Now

$$d(x_{n+1}, x_n) = d(f(x_n), f(x_{n-1}))$$

$$\frac{d(x_{n+1}, x_n)}{d(x_n, x_{n-1})} = \frac{d(x_n, x_{n-1})}{1 - }$$

$$d(fx, fy) \leq kd(x, y) \text{ for all } x, y \in X$$

The well-known Banach [1] contraction principle states that a contraction mapping of a Complete Metric space  $X$  into itself has a unique fixed points. This theorem has been extensively used in proving existences and uniqueness of solutions to various functional equations, particularly differential and integral equations. Because of its wide spread applicability there has been a search for generalization of the Banach contraction Principle. This celebrated principle has been generalized by many author, viz. chu. and Diaz [2], Sehgal [3]. Holmes [4]. Reich [5],

Hardy and Rogers [6], Wong [7], Edelstein [11] and Kannan [10] others in various ways. In this paper we obtain yet an other generalization of this principle.

## III. MAIN RESULTS

Theorem: Let  $f$  be a continuous self-map defined on a complete metric space  $(X, d)$ . Further, let  $f$  satisfy the following condition.

$$\begin{aligned} & d(f(x), f(y)) && d(x, f(x)).d(y, f(y)) \\ & d(x, y) && \text{for all } x, y \in X, \text{ and for some } , \text{ unique fixed point in } X. \\ & [0,1] \text{ with a } + && \\ & < 1 \text{ and } && \text{is finite. Then } f \text{ has a} \end{aligned} \tag{A}$$

$$\frac{d(x_n, f(x)).d(x_{n+1}, f(x_{n+1}))}{d(x_n, x_{n+1})}$$

$$\frac{d(x_n, f(x_{n+1})).d(x_{n+1}, x_n)}{d(x_n, x_{n+1})}$$

$d(x_n, x_{n+1}) + d(x_n, x_{n-1})$  Which implies that

By the triangle inequality we have for

$m$   $n,$

$$\begin{aligned} & \frac{d(x_n, x_m)}{d(x_{n+1}, x_{n+2}) + \dots + d(x_{m-1}, x_m)} + \\ & \frac{d(x_1, x_0)}{d(x_n, x_{n+1}) + \dots + d(x_{m-1}, x_m)} \\ & < (k_{-k} + k_{-k} + \dots + k_{-k}) d(x_0, f(x_0)), \text{ where } k = \end{aligned}$$

$$\begin{cases} k_{-k} d(x_0, f(x_0))^n \\ 0 \text{ if } m, n \end{cases}$$

Since  $X$  is complete, therefore there exists a  $u \in X$  such that  $x_n \rightarrow u$ . Further, the continuity of  $f$  in  $X$  implies

$$\begin{aligned} f(u) &= f(\lim_{n \rightarrow \infty} x_n) \\ &= (\lim_{n \rightarrow \infty} f(x_n)) \\ &= u. \end{aligned}$$

Therefore  $u$  is a fixed point of  $f$  in  $X$ . Now, if there exists another point  $v \in X$  such that  $f(v) = v$ , then  $d(v, u) = d(f(v), u)$

$$\begin{aligned} & \frac{d(v, f(v)).d(u, f(u))}{d(v, u)} \\ &= \frac{d(v, u).d(u, u)}{d(v, u)} \\ &< d(v, u), \quad (< 1) \\ & \text{a contradiction. Hence } u \text{ is a unique fixed point of } f \text{ in } X. \end{aligned}$$

#### IV. REFERENCES

- [1]. Bonsall, F.F. (1962). Lectures on some fixed point theorems of functional analysis. Tata Institute of Fundamental Research, Bombay.
- [2]. Chu, S.C., and Diaz, J.B. (1965). Remark on a generalization of Banach's mappings.
- [3]. Sehgal, V.M., (1969). A fixed point theorem for mapping with a contractive iterate. Proc. Am. math. Soc., 23, 631-34.
- [4]. Holmes, R.D. (1969). On fixed and periodic points under sets of mappings. canad. math. Bull., 12, 183-22
- [5]. Reich, S (1971). Some remarks concerning contraction mapping. Canad. Math., Bull., 14, 121-24.
- [6]. Hardy, G.C. and Rogers, T. (1973). A generalization offixed point theorem of Reich. Canad. math. Bull., 16, 201-206
- [7]. Wong, C.S (1973). Common fixed points of two mapping. Pac. J. Math., 48, 299-312.
- [8]. Smart D.R. (1974). Fixed point Theorems. Cambridge University Press.
- [9]. Nadler (Jr.), S.B. (1968). Sequences of contractions and fixed points, Pac. J. Math., 27, 579-85
- [10]. Kannan, R. (1969). Some results on fixed Points- II Am. Math. Monthly, 76, 71-76.