



## ***The CCB Science 2 Service Distance Learning Program<sup>®</sup>***

*Please complete all required information and fax to 203.284.9500*

*For questions or assistance with the program, please email Jeff at JQuamme@ctcertboard.org*

### ***S2S 2055 DETOXIFICATION***

#### ***Module 2 Pre-Test***

1. The signs and symptoms of alcohol withdrawal generally start to take place\_\_\_\_\_.
  - a. 3-6 hours after the last drink
  - b. 6-24 hours after the last drink
  - c. 12-36 hours after the last drink
  - d. none of the above
  
2. The majority of alcohol seizures occur\_\_\_\_\_after the cessation or reduction of alcohol.
  - a. 12 hours
  - b. 24 hours
  - c. 36 hours
  - d. 48 hours
  
3. Signs of opioid intoxication include\_\_\_\_\_.
  - a. pinpoint pupils
  - b. fast pulse
  - c. high body temperature
  - d. sweating
  
4. The Office of National Drug Control Policy (ONDCP) reported that in 2002, fewer than 1 in \_\_\_\_\_persons with a substance use disorder were in treatment at any one time.
  - a. 5
  - b. 7
  - c. 10
  - d. 12
  
5. The highest dose of nicotine patch is\_\_\_\_\_mg.
  - a. 22
  - b. 33
  - c. 44
  - d. none of the above
  
6. Phencyclidine (PCP) remains in legitimate use for\_\_\_\_\_.
  - a. anesthesia for humans
  - b. veterinary anesthesia
  - c. psychiatric treatment
  - d. there is no legitimate use



## ***The CCB Science 2 Service Distance Learning Program<sup>®</sup>***

*Please complete all required information and fax to 203.284.9500*

*For questions or assistance with the program, please email Jeff at [JQuamme@ctcertboard.org](mailto:JQuamme@ctcertboard.org)*

7. Auricular (ear) acupuncture has been used as an adjunct treatment for \_\_\_\_\_ detoxification for about 30 years.
- opioid
  - alcohol
  - sedative hypnotic
  - all of the above
8. African Americans are at greater risk of the co occurrence of \_\_\_\_\_ which should be taken into account when placing and monitoring them on withdrawal medications.
- diabetes
  - hypertension
  - heart disease
  - both a and b
9. When engaging a/an \_\_\_\_\_ in the process of detoxification, moving through the process too quickly or abruptly can be perceived as showing a lack of caring and is considered contrary to trust building.
- Asian
  - American Indian
  - African American
  - Hispanic/Latino
10. The consensus panel recommends that if a person who is maintained on opioid agonist medications become detained or incarcerated for 30 days or less, then that individual should be \_\_\_\_\_.
- maintained on their usual dosage
  - increased in dose to prevent withdrawal
  - gradually tapered in dose
  - none of the above