

## RECOMMENDED APPLICATIONS

### ROW CROPS/PERMANENT CROPS

**Post Harvest/Pre-Plant:** 1 qt. - 2 qt. per acre.  
For fields with high residue apply up to 2 gallons per acre.

**Seed Treatment:** 8oz. - 16oz. per 100lbs of seed.

**Planting:** 1 qt. - 2qt. per acre.

**In Season:** For fields with high residue and/or problem areas - up to 2 qt. per acre.

### FOR BEST RESULTS IN AGRICULTURAL APPLICATIONS

- Ensure tank is free of residue, if using chlorinated water, let tank sit for 24 hrs (with lid removed) before mixing.
  - Do not apply when a crop is severely stressed.
  - Do not mix with fungicides or insecticides and do not apply within 2 weeks of applying fungicides.
  - May be mixed with fertilizer and/or herbicide.
- 
- Use only as directed on the label.
  - Shake well before using.
  - Keep out of the reach of children.
  - Eye Contact: Flush with water/call physician.
  - Skin Contact: Exposure may result in dry skin. Rinse/Apply lotion.
  - Ingestion: May cause nausea or vomiting. Seek immediate medical attention.
  - In case of spills disperse with water, treat like surfactant or soap spillage.
- Not classified according to US GHS in compliance with OSHA 1910.1200

# N<sup>®</sup>TEXX EDGE

N-TEXX<sup>®</sup> Edge is a symbiotic family of selective microbial species that function as a soil inoculant intended to introduce microbes in the soil and increase the active biomass. The product makes no nutrient claims, but it introduces microbes into the soil resulting in improved plant vigor and nutrient cycling.

### CONTAINS NONPLANT FOOD INGREDIENTS

#### Active Ingredients:

|                         |                       |        |
|-------------------------|-----------------------|--------|
| Pseudomonas putida      | 2.5 x 10 <sup>5</sup> | cfu/mL |
| Pseudomonas alcaligenes | 1 x 10 <sup>5</sup>   | cfu/mL |
| Pseudomonas fluorescens | 2.5 x 10 <sup>5</sup> | cfu/mL |
| Bacillus subtilis       | 1 x 10 <sup>4</sup>   | cfu/mL |
| Bacillus licheniformis  | 2.5 x 10 <sup>5</sup> | cfu/mL |
| Bacillus pumilus        | 2.5 x 10 <sup>5</sup> | cfu/mL |
| Acidovorax delafieldii  | 1 x 10 <sup>5</sup>   | cfu/mL |
| Hydrogenphaga - atypica | 1 x 10 <sup>5</sup>   | cfu/mL |

1.50 % Humic Acid (Derived from Leonardite)

#### Inert Ingredients:

97.10 % Water (Microbial Solution)

**Disclaimer or Conditions of Sale:** The information contained on this label is believed to be accurate and reliable at the time of shipping. Buyer and user acknowledge and assume all liability resulting from the use of the material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller. Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

 **MADE IN THE USA**  
www.cxinternational.com  
1100 E. Sandy Lake Rd.  
Coppell, TX 75019  
Ph: 972-471 7775

Store inside out of direct sun in moderate temperatures between 50° - 90° F. Expires 2 years after date on stamp.

**LOT NO.** \_\_\_\_\_ **DATE** \_\_\_\_\_

Net Contents: 275 Gallons | 1040 L      Weight: 2310 lbs | 1048 Kg