

**Solomon Press**  
**Core Mathematics C2**  
**Paper B**  
**(Mark Scheme)**

**All exam papers are issued free to students for education purpose only.  
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)  
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations  
Advanced Subsidiary

## Core Mathematics C2

Paper B

### MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



*Written by Shaun Armstrong*

© *Solomon Press*


*These sheets may be copied for use solely by the purchaser's institute.*

## C2 Paper B – Marking Guide

1.	$\log_5 \frac{4x+3}{x-1} = 2$ $\frac{4x+3}{x-1} = 5^2 = 25$ $4x + 3 = 25(x - 1)$ $21x = 28, \quad x = \frac{4}{3}$	M1 M1 M1 A1	(4)										
2.	$\int_1^3 (x^2 - 2x + k) \, dx = \left[ \frac{1}{3}x^3 - x^2 + kx \right]_1^3$ $= (9 - 9 + 3k) - \left( \frac{1}{3} - 1 + k \right) = 2k + \frac{2}{3}$ $\therefore 2k + \frac{2}{3} = 8\frac{2}{3}, \quad k = 4$	M1 A2 M1 M1 A1	(6)										
3.	<p>(a) <math>= 1 + n\left(\frac{1}{4}x\right) + \frac{n(n-1)}{2}\left(\frac{1}{4}x\right)^2 + \dots</math></p> <p><math>= 1 + \frac{1}{4}nx + \frac{1}{32}n(n-1)x^2 + \dots</math></p> <p>(b) <math>\frac{1}{4}n = \frac{1}{32}n(n-1)</math></p> <p><math>8n = n(n-1)</math></p> <p><math>n[8 - (n-1)] = 0</math></p> <p><math>n \neq 0 \therefore n = 9</math></p>	B1 M1 A1 M1 M1 A1	(6)										
4.	$3(1 - \sin^2 x) + \sin^2 x + 5 \sin x = 0$ $2 \sin^2 x - 5 \sin x - 3 = 0$ $(2 \sin x + 1)(\sin x - 3) = 0$ $\sin x = 3 \text{ (no solutions) or } -\frac{1}{2}$ $x = 180 + 30, 360 - 30$ $x = 210, 330$	M1 A1 M1 A1 B1 M1 A1	(7)										
5.	<p>(a) <math>(x+1)^2 + (y-6)^2 = (2\sqrt{5})^2</math></p> <p><math>(x+1)^2 + (y-6)^2 = 20</math></p> <p>(b) sub. <math>y = 3x - 1</math> into eqn of C:</p> <p><math>(x+1)^2 + [(3x-1)-6]^2 = 20</math></p> <p><math>(x+1)^2 + (3x-7)^2 = 20</math></p> <p><math>x^2 - 4x + 3 = 0</math></p> <p><math>(x-1)(x-3) = 0</math></p> <p><math>x = 1, 3</math></p> <p>(c) <math>x = 1 \Rightarrow y = 2 \therefore (1, 2), \quad x = 3 \Rightarrow y = 8 \therefore (3, 8)</math></p> <p><math>AB = \sqrt{(3-1)^2 + (8-2)^2} = \sqrt{4+36} = \sqrt{40} = \sqrt{4 \times 10} = 2\sqrt{10}</math></p>	M1 A1 M1 A1 M1 A1 B1 M1 A1	(9)										
6.	<p>(a) <math>\frac{dy}{dx} = 4 - x^{-2}</math></p> <p>for minimum, <math>4 - x^{-2} = 0</math></p> <p><math>x^2 = \frac{1}{4}</math></p> <p><math>x &gt; 0 \therefore x = \frac{1}{2} \therefore \left(\frac{1}{2}, 4\right)</math></p> <p>(b)</p> <table style="margin-left: 20px; border-collapse: collapse;"> <tr> <td style="padding-right: 10px;"><math>x</math></td> <td style="padding-right: 10px;">1</td> <td style="padding-right: 10px;">2</td> <td style="padding-right: 10px;">3</td> <td style="padding-right: 10px;">4</td> </tr> <tr> <td><math>4x + x^{-1}</math></td> <td>5</td> <td><math>8\frac{1}{2}</math></td> <td><math>12\frac{1}{3}</math></td> <td><math>16\frac{1}{4}</math></td> </tr> </table> <p>area <math>\approx \frac{1}{2} \times 1 \times [5 + 16\frac{1}{4} + 2(8\frac{1}{2} + 12\frac{1}{3})]</math></p> <p><math>= 31.5</math> (3sf)</p>	$x$	1	2	3	4	$4x + x^{-1}$	5	$8\frac{1}{2}$	$12\frac{1}{3}$	$16\frac{1}{4}$	M1 A1 M1 A2 B1 B1 M1 A1 A1	(10)
$x$	1	2	3	4									
$4x + x^{-1}$	5	$8\frac{1}{2}$	$12\frac{1}{3}$	$16\frac{1}{4}$									

7. (a)  $r = \frac{114}{120} = 0.95$  M1  
 $u_5 = 120 \times (0.95)^4 = 97.74$  M1  
 $\therefore$  1 hour 38 minutes A1
- (b)  $S_8 = \frac{120[1-(0.95)^8]}{1-0.95}$  M1 A1  
 $= 807.79\dots$  minutes  $\approx$  13 hours 28 minutes A1
- (c)  $120 \times (0.95)^{n-1} < 60$  M1  
 $(n-1) \lg 0.95 < \lg 0.5$  M1  
 $n > \frac{\lg 0.5}{\lg 0.95} + 1$  A1  
 $n > 14.51 \therefore$  15 papers A1 (10)

8. (a)  $BD^2 = 6^2 + 9^2 - (2 \times 6 \times 9 \times \cos 60)$  M1 A1  
 $BD^2 = 36 + 81 - 54 = 63$   
 $BD = \sqrt{63} = \sqrt{9 \times 7} = 3\sqrt{7}$  cm M1 A1
- (b)  $(3\sqrt{7})^2 = 3^2 + 8^2 - (2 \times 3 \times 8 \times \cos C)$  M1  
 $\cos C = \frac{9+64-63}{48} = \frac{5}{24}$   
 $\angle BCD = 78.0^\circ$  (1dp) M1 A1
- (c)  $= (\frac{1}{2} \times 6 \times 9 \times \sin 60) + (\frac{1}{2} \times 3 \times 8 \times \sin 77.975)$  M2  
 $= 35.1 \text{ cm}^2$  (3sf) A1 (10)

9. (a)  $f(1) = 1 - 9 + 24 - 16 = 0$  B1  
 $\therefore (x-1)$  is a factor of  $f(x)$  B1
- (b) 
$$\begin{array}{r} x^2 - 8x + 16 \\ x-1 \overline{) x^3 - 9x^2 + 24x - 16} \\ \underline{x^3 - x^2} \phantom{- 16} \\ -8x^2 + 24x \phantom{- 16} \\ \underline{-8x^2 + 8x} \phantom{- 16} \\ 16x - 16 \\ \underline{16x - 16} \\ 0 \end{array}$$
 M1 A1
- $f(x) = (x-1)(x^2 - 8x + 16)$   
 $f(x) = (x-1)(x-4)^2$  [ $p = -1, q = -4$ ] M1 A1
- (c)  B2
- (d)  $= \int_1^4 (x^3 - 9x^2 + 24x - 16) dx$   
 $= [\frac{1}{4}x^4 - 3x^3 + 12x^2 - 16x]_1^4$  M1 A2  
 $= [(64 - 192 + 192 - 64) - (\frac{1}{4} - 3 + 12 - 16)]$  M1  
 $= 6\frac{3}{4}$  A1 (13)

Total (75)