

## Math 2471 Calculus III – Sample Final Questions

1. Find the unit tangent and unit normal vector for  $\vec{r}(t) = \langle t, \frac{1}{2}t^2 \rangle$ .

2. (i) Prove the limits either exist or doesn't exist

$$(i) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 2y^2}{x^2 + y^2} \quad (ii) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y^4}{x^2 + y^2}$$

3. Find the equation of the tangent plane to the given surface at the specified point

$$x^2y + xz + yz^2 = 3, \quad P(1, 2, -1)$$

4. Find the directional derivative of  $z = x^2 + 3xy + y^2$  at  $(1, 1)$  in the direction of  $\langle -3, 4 \rangle$

5. Classify the critical points for  $z = x^2y - x^2 + y^2 - 18y$ .

6. Set up and evaluate an integral to calculate

(i) the volume bound by  $z = 0$ ,  $z = 1 - x^2 - y^2$

(ii) the volume inside both  $x^2 + y^2 + z^2 = 2$ ,  $x^2 + y^2 = 1$

(1)

7. Set of the triple integral  $\iiint f(x, y, z) dV$  in both cylindrical and spherical coordinates for the volume inside the cone  $z = \sqrt{x^2 + y^2}$  bound by  $z = 1$

8. Is the following vector field conservative?

$$\vec{F} = \langle y^2 + 3yz, 2xy + 3xz, 3xy \rangle.$$

If so, determine the potential function  $f$  such that  $\vec{F} = \vec{\nabla} f$ . Use this to evaluate

$$\int_c (y^2 + 3yz) dx + (2xy + 3xz) dy + 3xy dz.$$

where  $c$  is a curve that connects the points  $(0, 0, 0)$  and  $(1, 2, 3)$ .

9. Evaluate the following line integrals:

(i)  $\int_c 2xy dx + (x + 1) dy$  where  $c$  is the counterclockwise direction around the square with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(1, 1)$  and  $(0, 1)$

(ii)  $\int_c (x - y) dx + (x + y) dy$  where  $c$  is clockwise direction around a circle of radius 2.

10. Verify Green's Theorem

$$\oint_c Pdx + Qdy = \iint_R (Q_x - P_y) dA$$

where  $\vec{F} = \langle 3x^2y, x^3 + x \rangle$  and  $c$  the curve enclosing the area given by  $y = x$  and  $y = x^2$ .

11. Evaluate the following surface integrals

- (i)  $\iint_S xy dS$  where  $S$  is the portion of the plane  $2x + y + z = 6$  in the first octant.
- (ii)  $\iint_S (x + z) dS$  where  $S$  is the portion of  $y^2 + z^2 = 9$  in the first octant between  $x = 0$  and  $x = 4$ .

12. Verify the divergence theorem

$$\iint_S \vec{F} \cdot \vec{n} dS = \iiint_V \nabla \cdot \vec{F} dV$$

where  $\vec{F} = \langle x + yz, y + xz, z + xy \rangle$  and  $V$  is the volume of the tetrahedron bound by  $x + y + z = 1$  and the planes  $x = 0, y = 0$  and  $z = 0$ .