

## Sample Test 2 Solutions

1. Do the following converge (explain)?

$$(1.1) \sum_{n=1}^{\infty} \frac{\ln n}{n^4 + 1},$$

Since  $\ln n < n$  for  $n \geq 1$ , then  $\frac{\ln n}{n^4 + 1} < \frac{n}{n^4 + 1} < \frac{1}{n^3}$ . Since  $\sum_{n=1}^{\infty} \frac{1}{n^3}$  ( $p = 3$ ) converges, then by the DCT the original series converges.

$$(1.2) \sum_{n=1}^{\infty} \frac{1}{n^3 + 1},$$

Compare with  $\sum_{n=1}^{\infty} \frac{1}{n^3}$ . Since  $\lim_{n \rightarrow \infty} \frac{\frac{1}{n^3}}{\frac{1}{n^3 + 1}} = 1$  and  $\sum_{n=1}^{\infty} \frac{1}{n^3}$  converges ( $p$  series with  $p = 3$ ) then by the limit comparison test (LCT), the original series converges.

$$(1.3) \sum_{n=1}^{\infty} \left( \frac{1}{2} + \frac{1}{n} \right)^n,$$

Taking the limit  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \lim_{n \rightarrow \infty} \left( \frac{1}{2} + \frac{1}{n} \right) = \frac{1}{2} < 1$  then by the  $n^{\text{th}}$  root test, the original series converges.

$$(1.4) \sum_{n=1}^{\infty} \frac{e^n}{n!},$$

Consider  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{e^{n+1}}{(n+1)!} / \frac{e^n}{(n)!} = \lim_{n \rightarrow \infty} \frac{e^{n+1}}{(n+1)!} \cdot \frac{(n)!}{e^n}$   
 $= \lim_{n \rightarrow \infty} \frac{e}{n+1} = 0 < 1$  so by ratio test, the series converges

$$(1.5) \sum_{n=1}^{\infty} \frac{1}{\ln(n+1)},$$

Since  $\ln(n+1) < n+1$  for  $n \geq 1$  then  $\frac{1}{n+1} < \frac{1}{\ln(n+1)}$  for  $n \geq 1$  and since  $\sum_{n=1}^{\infty} \frac{1}{(n+1)}$  (harmonic) diverges, then by the DCT, original series does as well.

$$(1.6) \sum_{n=1}^{\infty} \frac{1}{n(n+1)},$$

Comparing with  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  then  $\lim_{n \rightarrow \infty} \frac{1}{n(n+1)} / \frac{1}{n^2} = \lim_{n \rightarrow \infty} \frac{n^2}{n(n+1)} = 1$ , and since  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges ( $p$ -series with  $p = 2$ ) then by the limit comparison test (LCT) the original series converges.

$$(1.7) \sum_{n=1}^{\infty} \frac{n-1}{n+1},$$

Since  $\lim_{n \rightarrow \infty} \frac{n-1}{n+1} = 1$ , then by the  $n^{\text{th}}$  term test for divergence, the series diverges.

$$(1.8) \sum_{n=1}^{\infty} \frac{(2n)!}{(n!)^2},$$

Consider  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{(2n+2)!}{(n+1)!^2} / \frac{(2n)!}{(n)!^2} = \lim_{n \rightarrow \infty} \frac{(2n+2)!}{(2n)!} \cdot \frac{(n+1)!^2}{n!^2}$   
 $= \lim_{n \rightarrow \infty} \frac{(2n+2)(2n+1)}{(n+1)(n+1)} = 4 > 1$  so by ratio test, the series diverges

$$(1.9) \sum_{n=2}^{\infty} \frac{1}{\ln^2(n)},$$

Since  $\ln n < n$  for  $n \geq 1$  then  $\ln^2 n < n \ln n$  for  $n \geq 1$  which gives  $\frac{1}{n \ln n} < \frac{1}{\ln^2 n}$  for  $n \geq 1$ . Since  $\sum_{n=1}^{\infty} \frac{1}{n \ln n}$  diverges, (see next question) then by the direct comparison test, original series does as well.

$$(1.10) \sum_{n=3}^{\infty} \frac{1}{n \ln n},$$

Let  $f(x) = \frac{1}{x \ln x}$ . Clearly  $f(x) > 0$  and  $f'(x) = -\frac{\ln x + 1}{(x \ln x)^2}$  for  $x \geq 3$  showing that  $f(x)$  is decreasing so that the integral test may be used. Consider

$$\int_3^{\infty} \frac{dx}{x \ln x} = \lim_{b \rightarrow \infty} \int_3^b \frac{dx}{x \ln x} = \lim_{b \rightarrow \infty} \ln \ln x \Big|_3^b = \infty.$$

Since the integral diverges, then by the integral test, the series does as well.

$$(1.11) \sum_{n=1}^{\infty} \frac{1}{2^n + 1}.$$

Compare with  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ . Then  $\lim_{n \rightarrow \infty} \frac{1}{2^n} / \frac{1}{2^n + 1} = \lim_{n \rightarrow \infty} \frac{2^n + 1}{2^n} = 1$  and since  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  converges (geometric series with  $r = 1/2$ ), the original series converges by the LCT.

2. Determine whether the following series converge absolutely, conditionally or diverge

$$(2.1) \sum_{n=1}^{\infty} \frac{(-1)^n(n-1)}{n+1},$$

Since  $\lim_{n \rightarrow \infty} \frac{(-1)^n(n-1)}{n+1} = (-1)^n \neq 0$  this series diverges.

$$(2.2) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n(n+1)'}}$$

We first consider  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n(n+1)'}}$  and by limit comparison with  $\sum_{n=1}^{\infty} \frac{1}{n}$  shows that we do not have absolute convergence. So we check the two conditions for conditional convergence. If we let  $a_n = \frac{1}{\sqrt{n(n+1)'}}$ , then clearly

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n(n+1)'}} = 0.$$

Next, we need to show  $a_{n+1} < a_n$ . We could show

$$\frac{1}{\sqrt{(n+1)(n+2)'}} \stackrel{?}{\leq} \frac{1}{\sqrt{n(n+1)'}}$$

but is easier to show that if

$$f(x) = \frac{1}{\sqrt{x(x+1)'}} \quad \text{then} \quad f'(x) = -\frac{2x+1}{2(x^2+x)^{3/2}} < 0 \quad \text{for } x \geq 1$$

so by the alternating series test (AST), the series converges conditionally.

$$(2.3) \quad \sum_{n=1}^{\infty} \frac{(-1)^n n^n}{n!}$$

We let  $a_n = \frac{n^n}{n!}$ . We stop the series from alternating and use the ration test.

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{(n+1)^{n+1}}{(n+1)!} / \frac{n^n}{n!} = \lim_{n \rightarrow \infty} \frac{(n+1)(n+1)^n}{(n+1)n!} \cdot \frac{n!}{n^n} = \\ \lim_{n \rightarrow \infty} \frac{(n+1)^n}{n^n} &= \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e > 1 \end{aligned}$$

so by ratio test, the series does not converge absolutely. Now we go the the AST. We will show that the terms are not decreasing. So we want to show that  $\frac{n^n}{n!} < \frac{(n+1)^{n+1}}{(n+1)!}$  so  $\frac{n^n}{n!} < \frac{(n+1)(n+1)^n}{(n+1)n!}$  and after canceling  $n^n < (n+1)^n$  or  $n < n+1$  which is true so the series diverges by the AST.

$$(2.4) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{2^n + 3^n}'$$

We first consider  $\sum_{n=1}^{\infty} \frac{1}{2^n + 3^n}$  and compare with  $\sum_{n=1}^{\infty} \frac{1}{3^n}$ . By the LCT

$$\lim_{n \rightarrow \infty} \frac{1}{3^n} / \frac{1}{2^n + 3^n} = \lim_{n \rightarrow \infty} \frac{2^n + 3^n}{3^n} = \lim_{n \rightarrow \infty} 1 + \left(\frac{2}{3}\right)^n = 1,$$

and since  $\sum_{n=1}^{\infty} \frac{1}{3^n}$  converges (geometric series  $r = 1/3$ ), the original series converges absolutely.

$$(2.5) \quad \sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2 + 1},$$

We first consider  $\sum_{n=1}^{\infty} \frac{n}{n^2 + 1}$  and compare with  $\sum_{n=1}^{\infty} \frac{1}{n}$ . By the LCT

$$\lim_{n \rightarrow \infty} \frac{1}{n} / \frac{n}{n^2 + 1} = \lim_{n \rightarrow \infty} \frac{n^2 + 1}{n^2} = 1,$$

which show that original series doesn't converge absolutely since we compared with the harmonic series that diverges. If we let  $a_n = \frac{n}{n^2 + 1}$ , then clearly

$$\lim_{n \rightarrow \infty} \frac{n}{n^2 + 1} = 0.$$

Next, we need to show  $a_{n+1} < a_n$ . If we let

$$f(x) = \frac{x}{x^2 + 1} \quad \text{then} \quad f'(x) = \frac{-x^2 + 1}{(x^2 + 1)^2} < 0 \quad \text{for} \quad x > 1$$

so by the alternating series test (AST), the series converges conditionally.

$$(2.6) \quad \sum_{n=1}^{\infty} \frac{(-1)^n n}{n + 1}.$$

Since  $\lim_{n \rightarrow \infty} \frac{(-1)^n n}{n + 1} = (-1)^n \neq 0$  this series diverges.

3. Determine the interval of convergence of the following.

$$\sum_{n=1}^{\infty} \frac{2^n x^n}{\sqrt{n+1}}, \quad \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2}, \quad \sum_{n=1}^{\infty} \frac{(x-1)^n}{n^2},$$

$$3(i) \quad \sum_{n=1}^{\infty} \frac{2^n x^n}{\sqrt{n+1}},$$

Choosing

$$a_n = \frac{2^n x^n}{\sqrt{n+1}}$$

then

$$\lim_{x \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{x \rightarrow \infty} \left| \frac{2^{n+1} x^{n+1}}{\sqrt{n+2}} / \frac{2^n x^n}{\sqrt{n+1}} \right| = \lim_{x \rightarrow \infty} 2 \frac{\sqrt{n+1}}{\sqrt{n+2}} |x| = 2|x| < 1$$

So  $|x| < \frac{1}{2}$  or  $-\frac{1}{2} < x < \frac{1}{2}$ . Checking the endpoints gives

$$x = -\frac{1}{2} \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n+1}}, \quad \text{which converges by AST}$$

$$x = \frac{1}{2} \quad \sum_{n=1}^{\infty} \frac{1}{\sqrt{n+1}}, \quad \text{which diverges by DCT with } p \text{ series } (p = 1/2)$$

Therefore the interval of convergence is  $-\frac{1}{2} \leq x < \frac{1}{2}$ .

$$3(ii) \quad \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2},$$

Choosing

$$a_n = \frac{(-1)^n x^{2n}}{2^{2n} n!^2}$$

then

$$\lim_{x \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{x \rightarrow \infty} \left| \frac{(-1)^{n+1} x^{2n+2}}{2^{2n+2} (n+1)!^2} / \frac{(-1)^n x^{2n}}{2^{2n} n!^2} \right| = \lim_{x \rightarrow \infty} \frac{1}{4(n+1)^2} |x^2| = 0 < 1$$

so series converges for all  $x$ .

$$3(iii) \quad \sum_{n=1}^{\infty} \frac{(2x-1)^n}{n^2},$$

Choosing

$$a_n = \frac{(2x-1)^n}{n^2}$$

then

$$\lim_{x \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{x \rightarrow \infty} \left| \frac{(2x-1)^{n+1}}{(n+1)^2} / \frac{(2x-1)^n}{n^2} \right| = \lim_{x \rightarrow \infty} \frac{n^2}{(n+1)^2} |2x-1| = |2x-1| < 1$$

So  $|2x-1| < 1$  or  $-1 < 2x-1 < 1$  or  $0 < 2x < 2$  or  $0 < x < 1$ . Checking the endpoints gives

$$x = 0 \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}, \quad \text{which converges absolutely, it's a } p \text{ series}$$

$$x = 1 \quad \sum_{n=1}^{\infty} \frac{1}{n^2}, \quad \text{which converges, it's a } p \text{ series}$$

Therefore the interval of convergence is  $0 \leq x \leq 1$ .

4. Calculate the  $n^{\text{th}}$  degree Taylor polynomial with remainder for the following. Expand about the point  $x = c$

(4.1)  $f(x) = e^x$ ,  $c = 0$ ,  $n = 2$

$$\begin{aligned} f(x) &= e^x & f(0) &= 1, \\ f'(x) &= e^x & f'(0) &= 1, \\ f''(x) &= e^x & f''(0) &= 1, \\ f'''(x) &= e^x & & \text{for the remainder,} \end{aligned}$$

The Taylor polynomial is

$$\begin{aligned} P_2(x) &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2, \\ &= 1 + x + \frac{x^2}{2!}, \end{aligned}$$

The remainder is given by

$$R_2(x) = \frac{e^z}{3!}x^3,$$

for  $0 < z < x$  or  $x < z < 0$

(4.2)  $f(x) = \sin x$ ,  $c = \frac{\pi}{2}$   $n = 4$ ,

In this example, we need only construct  $P_4$ .

$$\begin{aligned} f(x) &= \sin x & f\left(\frac{\pi}{2}\right) &= 1, \\ f'(x) &= \cos x & f'\left(\frac{\pi}{2}\right) &= 0, \\ f''(x) &= -\sin x & f''\left(\frac{\pi}{2}\right) &= -1, \\ f'''(x) &= -\cos x & f'''\left(\frac{\pi}{2}\right) &= 0, \\ f^{(4)}(x) &= \sin x & f^{(4)}\left(\frac{\pi}{2}\right) &= 1, \end{aligned}$$

The Taylor polynomial is

$$\begin{aligned} P_4(x) &= f\left(\frac{\pi}{2}\right) + \frac{f'\left(\frac{\pi}{2}\right)}{1!}\left(x - \frac{\pi}{2}\right) + \frac{f''\left(\frac{\pi}{2}\right)}{2!}\left(x - \frac{\pi}{2}\right)^2 + \cdots + \frac{f^{(4)}\left(\frac{\pi}{2}\right)}{4!}\left(x - \frac{\pi}{2}\right)^4, \\ &= 1 - \frac{1}{2!}\left(x - \frac{\pi}{2}\right)^2 + \frac{1}{4!}\left(x - \frac{\pi}{2}\right)^4, \end{aligned}$$

The remainder is given by

$$\begin{aligned} R_4(x) &= \frac{f^{(5)}(z)}{5!} \left(x - \frac{\pi}{2}\right)^5, \\ &= \frac{\cos z}{5!} \left(x - \frac{\pi}{2}\right)^5, \end{aligned}$$

for  $\frac{\pi}{2} < z < x$  or  $x < z < \frac{\pi}{2}$ .

(4.3)  $f(x) = \ln(x+1)$ ,  $c = 0$ ,  $n = 3$

$$\begin{aligned} f(x) &= \ln(x+1) & f(0) &= 0, \\ f'(x) &= \frac{1}{x+1} & f'(0) &= 1, \\ f''(x) &= \frac{-1}{(x+1)^2} & f''(0) &= -1, \\ f'''(x) &= \frac{2}{(x+1)^3} & f'''(0) &= 2, \\ f^{(4)}(x) &= \frac{-3!}{(x+1)^4} \text{ for the remainder.} \end{aligned}$$

The Taylor polynomial is

$$\begin{aligned} P_3(x) &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3, \\ &= 0 + x - \frac{1}{2!}x^2 + \frac{2!}{3!}x^3, \\ &= x - \frac{x^2}{2} + \frac{x^3}{3}, \end{aligned}$$

The remainder is given by

$$\begin{aligned} R_3(x) &= \frac{f^{(4)}(z)}{4!}x^4, \\ &= -\frac{3!}{(z+1)^4} \frac{x^4}{4!}, \end{aligned}$$

for  $0 < z < x$  or  $x < z < 0$ .

(4.4)  $f(x) = \frac{1}{2-x}$ ,  $c = 0$ ,  $n = 3$ .

In this case we only need  $P_3$ . The derivatives are:

$$\begin{aligned}f(x) &= \frac{1}{(2-x)}, & f(0) &= \frac{1}{2}, \\f'(x) &= \frac{1}{(2-x)^2}, & f'(0) &= \frac{1}{2^2}, \\f''(x) &= \frac{2}{(2-x)^3}, & f''(0) &= \frac{2}{2^3}, \\f'''(x) &= \frac{3!}{(2-x)^4}, & f'''(0) &= \frac{3!}{2^4}, \\f^{(4)}(x) &= \frac{4!}{(2-x)^4}, & \text{remainder,}\end{aligned}$$

The Taylor polynomial is

$$\begin{aligned}P_3(x) &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3, \\&= \frac{1}{2} + \frac{x}{2^2} + \frac{2!}{2^3} \frac{x^2}{2!} + \frac{3!}{2^4} \frac{x^3}{3!}, \\&= \frac{1}{2} + \frac{x}{2^2} + \frac{x^2}{2^3} + \frac{x^3}{2^4},\end{aligned}$$

The remainder is given by

$$\begin{aligned}R_3(x) &= \frac{f^{(4)}(z)}{4!}x^4, \\&= \frac{1}{(2-z)^5}x^4.\end{aligned}$$

for  $0 < z < x$  or  $x < z < 0$ .