

2017 Whatcom County BLS Protocol Test

Name: _____ Date: _____ Score: _____%

Circle or label the correct answers

1. **True or False**

BLS transport is appropriate for STABLE patients after EPI administration and where additional EPI is available to give if needed.

- A. True
- B. False

2. **True or False**

The maximum dose of Naloxone (Narcan) is 6ml or 3 complete doses.

- A. True
- B. False

3. If at any time you are not certain about a treatment or need guidance, you may contact_____. (**Select all that apply**)

- A. The Medical Control Physician, called the “STAR DOC” at 360-715-4149
- B. Anyone who has first aid/CPR training
- C. Retired Medics who happen to be on scene
- D. Doctors who happen to be on scene
- E. The incoming Medic Unit

4. **True or False**

In regards to head trauma, patients on blood thinners MUST be transported. The “STAR DOC” MUST be consulted if the patient refuses transport.

- A. True
- B. False

5. Number these steps (1-5) in the correct order for obtaining a DEX or blood glucose reading.

- _____ Use lancet
- _____ Wipe first blood drop with dry gauze
- _____ Clean area with alcohol wipe
- _____ Take sample
- _____ Allow to dry completely

6. **True or False**

CODE RED transports eliminate the need for the due regard of everyone's safety. Getting the patient to definitive care fast is our number one concern.

- A. True
- B. False

7. The EMT in charge of patient care is responsible for completing a CQI (Continuous Quality Improvement) online report for calls where which of the following were used, attempted (successful or unsuccessful), or administered? **(Select ALL that apply)**

- A. AED
- B. Res-Q-Pump and Pod
- C. Naloxone (Narcan)
- D. Oxygen
- E. 12 Lead ECG
- F. Epinephrine
- G. KED (Kendrick Extrication Device)
- H. Kingtube

8. **True or False**

Respirations greater than 30 breaths per minute and fractures at more than one location is an ALS Indicator?

- A. True
- B. False

9. What are the contraindications for Epinephrine during an allergic reaction?

- A. Difficulty Swallowing
- B. Past Stroke
- C. Blood Pressure greater than 100
- D. There are no contraindications when used in a life threatening emergency

10. **True or False**

After delivery of a baby, place two clamps on the cord two inches apart and six inches away from the baby. Cut the cord between the clamps.

- A. True
- B. False

11. Stroke signs less than _____hrs without ALS Indicators indicate a BLS Code Red Transport.

- A. 1.5hrs
- B. 2.5hrs
- C. 3.5hrs
- D. 4.5hrs

12. Please circle all the correct answers in regards to the Pelvic Sling.

- A. The Sling wraps the hips and buttocks, not the waist
- B. The Sling is a single-use, disposable item
- C. Once the Sling is in place don't remove it
- D. A large Sling fits a 32-50 inch circumference

13. Answer the following scenario and label Sick or Not Sick according to the Pediatric Assessment Triangle.

You and your partner are seeing a 4-year-old male with an obvious distal forearm fracture resulting from a fall from a swing onto a rubber mat. He is conscious and crying, without respiratory distress. His skin signs are pink, warm and dry. Chief complaint: arm pain, MOI, fall with obvious forearm fracture (IOS - low)

- *Appearance: conscious alert and crying*
- *Work of breathing: without complaint*
- *Circulation to skin: warm and dry, normal color*

- A. Sick
- B. Not Sick

14. When you complete a patient report, the report must be faxed in within:

- A. 2 Hours
- B. 4 Hours
- C. 6 Hours
- D. 12 Hours

15. According to the Whatcom County BLS Protocols, if a Diabetic Patient has a blood sugar less than _____ and unable to safely eat or drink, this would be an ALS Upgrade.

- A. 50
- B. 60
- C. 70
- D. 80

16. In the acronym AEIOU TIPS, the E and S Stands for:

- A. Evaluate & Survey
- B. Epilepsy & Stroke
- C. Euphoria & Sick
- D. EMS & Safety

17. **True or False**

Recent stroke symptoms with no airway compromise is considered a Code Red BLS Transport?

- A. True
- B. False

18. Circle all that apply to for ALS Indicators in regards to Amputation

- A. Poor Skin Signs (Pale, Sweaty)
- B. Sustained tachycardia
- C. Hypotension
- D. Evidence of ongoing bleeding
- E. Unstable vital signs

19. **True or False**

Before insertion of the King Tube, pre-oxygenate the patient with high flow nasal oxygen with NFNO at 15 L/min and maintain until airway is secured.

- A. True
- B. False

20. Patients between _____ months and _____ years with only 1 seizure of short duration suspected to be febrile may go POV.

- A. 3 and 6
- B. 6 and 3
- C. 12 and 5
- D. 5 and 12