

Sample Final - Solutions

1. Find the unit tangent and unit normal vector for the following vector functions

$$\vec{r}(t) = \left\langle t, \frac{1}{2}t^2 \right\rangle$$

Soln.

$$\begin{aligned}\vec{r} &= \left\langle t, \frac{1}{2}t^2 \right\rangle \\ \vec{r}' &= \langle 1, t \rangle \\ \|\vec{r}'\| &= \sqrt{t^2 + 1}.\end{aligned}$$

so

$$\vec{T} = \frac{\vec{r}'}{\|\vec{r}'\|} = \left\langle \frac{1}{\sqrt{t^2 + 1}}, \frac{t}{\sqrt{t^2 + 1}} \right\rangle$$

Further

$$\begin{aligned}\vec{T}' &= \left\langle \frac{-t}{(t^2 + 1)^{3/2}}, \frac{1}{(t^2 + 1)^{3/2}} \right\rangle \\ \|\vec{T}'\| &= \frac{1}{t^2 + 1}.\end{aligned}$$

so

$$\vec{N} = \frac{\vec{T}'}{\|\vec{T}'\|} = \left\langle \frac{-t}{\sqrt{t^2 + 1}}, \frac{1}{\sqrt{t^2 + 1}}, 0 \right\rangle$$

2. Prove the limits either exist or do not exist. In the former case use the squeeze theorem.

$$(i) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 2y^2}{x^2 + y^2} \quad (ii) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y^4}{x^2 + y^2}$$

Soln. 2 (i)

$$\text{Along } y = 0, \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 2y^2}{x^2 + y^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{x^2}{x^2} = 1$$

$$\text{Along } y = x, \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 2y^2}{x^2 + y^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{3x^2}{2x^2} = \frac{3}{2}.$$

Since following different paths lead to different limits, the limit DNE.

Soln. 2 (ii) From the inequalities

$$\begin{aligned}-\sqrt{x^2 + y^2} &\leq x \leq \sqrt{x^2 + y^2} \\ -\sqrt{x^2 + y^2} &\leq y \leq \sqrt{x^2 + y^2}\end{aligned}$$

we have

$$\begin{aligned}-(x^2 + y^2) &\leq x^2 \leq (x^2 + y^2) \\ -(x^2 + y^2)^2 &\leq y^4 \leq (x^2 + y^2)^2\end{aligned}$$

which gives

$$-(x^2 + y^2)^3 \leq x^2 y^4 \leq (x^2 + y^2)^3.$$

Thus,

$$-(x^2 + y^2)^2 \leq \frac{x^2 y^4}{x^2 + y^2} \leq (x^2 + y^2)^2$$

and

$$-\lim_{(x,y) \rightarrow (0,0)} (x^2 + y^2)^2 \leq \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y^4}{x^2 + y^2} \leq \lim_{(x,y) \rightarrow (0,0)} (x^2 + y^2)^2.$$

Since

$$\lim_{(x,y) \rightarrow (0,0)} (x^2 + y^2)^2 = 0$$

by the squeeze theorem

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y^4}{x^2 + y^2} = 0.$$

3. Find the equation of the tangent plane to the given surface at the specified point

$$x^2 y + xz + yz^2 = 3, \quad P(1, 2, -1)$$

Soln. If we define $F = x^2 y + xz + yz^2 - 3$ then $F_x = 2xy + z$, $F_y = x^2 + z^2$ and $F_z = x + 2yz$. Evaluating these at the point P gives $F_x = 3$, $F_y = 2$ and $F_z = -3$. The equation of the tangent plane is thus $3(x - 1) + 2(y - 2) - 3(z + 1) = 0$

4. Find the directional derivative of $z = x^2 + 3xy + y^2$ at $(1, 1)$ in the direction of $\langle -3, 4 \rangle$.

Soln. The gradient is given by $\nabla z = \langle 2x + 3y, 3x + 2y \rangle$ and at the point $(1, 1)$ it becomes $\nabla z = \langle 5, 5 \rangle$. The direction derivative is then given by

$$\nabla z \cdot \frac{\vec{u}}{\|\vec{u}\|} = \langle 5, 5 \rangle \cdot \frac{\langle -3, 4 \rangle}{5} = \frac{-15 + 20}{5} = 1$$

5. Classify the critical points for

$$z = x^2y - x^2 + y^2 - 18y$$

Soln. The derivatives are

$$z_x = 2xy - 2x = 2x(y - 1), \quad z_y = x^2 + 2y - 18.$$

Setting each of these to zero gives the following critical points: $(0, 9)$, $(-4, 1)$, and $(4, 1)$. The second derivatives are:

$$z_{xx} = 2(y - 1), \quad z_{xy} = 2x, \quad z_{yy} = 2$$

giving $\Delta = z_{xx}z_{yy} - z_{xy}^2 = 4(y - 1) - 4x^2$. We now test each critical point

$(0, 9)$	$\Delta = 32 > 0$	$z_{yy} > 0$	min
$(-4, 1)$	$\Delta = -64 < 0$		saddle
$(4, 1)$	$\Delta = -64 < 0$		saddle

6 (i). Find the volume bound by the paraboloid $z = 1 - x^2 - y^2$ and the plane $z = 0$

Soln. The two surfaces intersect when $z = 0$ so $x^2 + y^2 = 1$. The volume is then obtained from the integral

$$\iint_R (1 - x^2 - y^2) dA$$

As the region of integration is a circle of radius 1, we switch to polar coordinates giving

$$\int_0^{2\pi} \int_0^1 (1 - r^2) r dr d\theta = \frac{\pi}{2}$$

6 (ii). Find the volume inside the sphere $x^2 + y^2 + z^2 = 2$ and the cylinder $x^2 + y^2 = 1$

Soln. The surfaces intersect when $z^2 = 1$ or $z = \pm 1$. The volume is then obtained from the integral

$$\iint_R 2\sqrt{2 - x^2 - y^2} dA$$

As the region of integration is a circle of radius 1, we switch to polar coordinates giving

$$\int_0^{2\pi} \int_0^1 2\sqrt{2 - r^2} r dr d\theta = \frac{8\sqrt{2} - 4}{3} \pi$$

6 (iii). Find the surface area of the plane $x + 2y + 3z = 6$ for $x, y, z \geq 0$.

Soln. The general formula is

$$\iint_R \sqrt{1 + z_x^2 + z_y^2} dA$$

Since $z = 2 - \frac{1}{3}x - \frac{2}{3}y$, the $z_x = -1/3$ and $z_y = -2/3$ giving

$$\iint_R \sqrt{1 + \frac{1}{9} + \frac{4}{9}} dA = \frac{\sqrt{14}}{3} \iint_R dA$$

Thus

$$\frac{\sqrt{14}}{3} \int_0^3 \int_0^{6-2y} dx dy = 3\sqrt{14}.$$

7. Set of the triple integral $\iiint f(x, y, z) dV$ in both cylindrical and spherical coordinates for the volume inside the cone $z = \sqrt{x^2 + y^2}$ and below the plane $z = 1$.

Soln - Cylindrical Eliminating z between the equations gives $x^2 + y^2 = 1$. This is the region of integration

$$\int_0^{2\pi} \int_0^1 \int_r^1 f(r \cos \theta, r \sin \theta, z) r dz dr d\theta$$

Soln - Spherical From the picture we see that $\phi = 0 \rightarrow \pi/4$. Further, $\rho = 0 \rightarrow 1/\cos \phi$ and $\theta = 0 \rightarrow 2\pi$ so

$$\int_0^{2\pi} \int_0^{\pi/4} \int_0^{\sec \phi} f(\rho \cos \theta \sin \phi, \rho \sin \theta \sin \phi, \rho \cos \phi) \rho^2 \sin \phi d\rho d\phi d\theta$$

8. Is the following vector field conservative?

$$\vec{F} = \langle y^2 + 3yz, 2xy + 3xz, 3xy \rangle.$$

Soln. Since $\nabla \times \vec{F} = 0$ then yes, the vector field is conservative. Thus ϕ exists such that $\vec{F} = \vec{\nabla} \phi$ so

$$\phi_x = y^2 + 3yz \Rightarrow \phi = x y^2 + 3xyz + A(y, z)$$

$$\phi_y = 2xy + 3xz \Rightarrow \phi = x y^2 + 3xyz + B(x, z)$$

$$\phi_z = 3xy \Rightarrow \phi = 3xyz + C(x, y)$$

Therefore we see that

$$\phi = x y^2 + 3xyz + c$$

and

$$\int_C (y^2 + 3yz) dx + (2xy + 3xz) dy + 3xy dz = x y^2 + 3xyz \Big|_{(0,0,0)}^{(1,2,3)} = 22.$$

9 (i). Evaluate the following line integral $\int_C 2xy dx + (x+1) dy$ where C is the counterclockwise direction around the square with vertices $(0,0)$, $(1,0)$, $(1,1)$ and $(0,1)$.

Soln. Here we have 4 separate curves which we denote by C_1 , C_2 , C_3 and C_4 .

$$C_1 : \text{ Here } y = 0, dy = 0 \text{ so } \int_{C_1} 0 = 0$$

$$C_2 : \text{ Here } x = 1, dx = 0 \text{ so } \int_0^1 2 dy = 2$$

$$C_3 : \text{ Here } y = 1, dy = 0 \text{ so } \int_1^0 2x dx = -1$$

C_4 : Here $x = 0, dx = 0$ so $\int_1^0 dy = -1$

Thus $\int_C 2xy dx + (x + 1) dy = 0 + 2 - 1 - 1 = 0$.

9 (ii). Evaluate the following line integral $\int_C (x - y) dx + (x + y) dy$ where c is clockwise direction around the circle of radius 2.

Soln. Here we parameterize the curve by $x = 2 \cos t, y = -2 \sin t, 0 \leq t \leq 2\pi$. Note the -2 on the y term as we are going clockwise and not counterclockwise. So $dx = -2 \sin t dt$ and $dy = -2 \cos t dt$. Thus, the line integral becomes

$$\int_0^{2\pi} -(2 \cos t + 2 \sin t)2 \sin t dt - (2 \cos t - 2 \sin t)2 \cos t dt = -8\pi$$

10. Green's Theorem is

$$\int_C P dx + Q dy = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Verify Green's Theorem where $\vec{F} = \langle 3x^2y, x^3 + x \rangle$ where R is the region bound by the curves $y = x^2$ and $y = x$.

Soln. We have two separate curves which we denote by C_1 and C_2 .

C_1 : Here $y = x^2, dy = 2x dx$ so $\int_0^1 3x^4 dx + (x^3 + x)2x dx = 5/3$

C_2 : Here $y = x, dy = dx$ so $\int_1^0 3x^3 dx + (x^3 + x)dx = -3/2$

so

$$\int_C 3x^2y dx + (x^3 + x) dy = 5/3 - 3/2 = 1/6.$$

For the second part, since $P = 3x^2y$ and $Q = x^3 + x$ then $Q_x - P_y = 3x^2 + 1 - 3x^2 = 1$ so

$$\iint_R (Q_x - P_y) dA = \int_0^1 \int_{x^2}^x 1 dy dx = 1/6$$

11 (i). Evaluate $\iint_S x y dS$ where S is the surface of the plane $2x + y + z = 6$.

Soln. Since $z = 6 - 2x - y$ then $dS = \sqrt{1 + z_x^2 + z_y^2} dA = \sqrt{1 + 4 + 1} dA$ and thus

$$\int_0^3 \int_0^{6-2x} \sqrt{6} x y dy dx = 27\sqrt{6}/2$$

11 (ii). Evaluate $\iint_S (x + z) dS$ where S is the surface of the cylinder $y^2 + z^2 = 9$ bound between $x = 0$ and $x = 4$ in the first octant.

Soln. Here, we'll parameterize the surface by $x = u$, $y = 3 \cos v$ and $z = 3 \sin v$, $0 \leq u \leq 4$ and $0 \leq v \leq \pi/2$. If we let $\vec{r} = \langle u, 3 \cos v, 3 \sin v \rangle$ then $\|\vec{r}_u \times \vec{r}_v\| = 3$ and we have

$$\int_0^4 \int_0^{\pi/2} (u + 3 \sin v) 3 dv du = 3(4\pi + 12)$$

12. Verify the divergence theorem

$$\iiint_V \nabla \cdot \vec{F} dV = \iint_S \vec{F} \cdot \vec{n} dS$$

where $\vec{F} = \langle x + yz, y + xz, z + xy \rangle$ and V is the volume of the tetrahedron bound by $x + y + z = 1$ and the planes $x = 0$, $y = 0$ and $z = 0$.

Soln. We will first deal with the surface integrals. There are 4 of them.

S_1 : Bottom. Here $z = 0$ so $\vec{F} = \langle x, y, xy \rangle$ and $\vec{n} = \langle 0, 0, -1 \rangle$.

Thus $\vec{F} \cdot \vec{n} = -xy$ and $\int_0^1 \int_0^{1-x} -x y dy dx = -1/24$

S_2 : Left. Here $y = 0$ so $\vec{F} = \langle x, xz, z \rangle$ and $\vec{n} = \langle 0, -1, 0 \rangle$.

Thus $\vec{F} \cdot \vec{n} = -xz$ and $\int_0^1 \int_0^{1-x} -x z dz dx = -1/24$

S_3 : Back. Here $x = 0$ so $\vec{F} = \langle yz, y, z \rangle$ and $\vec{n} = \langle -1, 0, 0 \rangle$.

Thus $\vec{F} \cdot \vec{n} = -yz$ and $\int_0^1 \int_0^{1-y} -y z dz dy = -1/24$

S_4 : Plane. Here $x + y + z = 1$ and $\vec{n} = \langle 1, 1, 1 \rangle / \sqrt{3}$.

Thus $\vec{F} \cdot \vec{n} = (1 + x + y - x^2 - xy - y^2) / \sqrt{3}$ and

$$\int_0^1 \int_0^{1-x} (1 + x + y - x^2 - xy - y^2) dy dx = 5/8$$

Therefore $\iint_S \vec{F} \cdot \vec{n} dS = -1/24 - 1/24 - 1/24 + 5/8 = 1/2$.

Second part. $\nabla \cdot \vec{F} = 3$ so $\int_0^1 \int_0^{1-x} \int_0^{1-x-y} 3 dz dy dx = 1/2$. Verified!

13. Verify Stokes's theorem for

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_S \nabla \times \vec{F} \cdot \vec{n} dS$$

where $\vec{F} = \langle xz - y, yz, 1 \rangle$ and S is the surface of the paraboloid $z = 1 - x^2 - y^2$ for $z \geq 0$ and C is the curve where the paraboloid intersects the plane $z = 0$.

Soln. We will first do the line integral. Here

$$\oint_C (xz - y)dx + yz dy + dz$$

but since we are on the curve where the paraboloid intersects the plane, then $z = 0$ and so we have

$$\oint_C -y dx$$

If we let $x = \cos t$ and $y = \sin t$ this integral becomes

$$\int_0^{2\pi} \sin^2 t dt = \pi.$$

For the second part, $\nabla \times \vec{F} = \langle -y, x, 1 \rangle$. On the surface of the paraboloid

$$\vec{n} = \frac{\langle 2x, 2y, 1 \rangle}{\sqrt{4x^2 + 4y^2 + 1}}$$

and $dS = \sqrt{1 + 4x^2 + 4y^2} dA$ so

$$\begin{aligned} \iint_S \nabla \times \vec{F} \cdot \vec{n} dS &= \iint_S \langle -y, x, 1 \rangle \cdot \frac{\langle 2x, 2y, 1 \rangle}{\sqrt{4x^2 + 4y^2 + 1}} \sqrt{1 + 4x^2 + 4y^2} dA \\ &= \iint_S 1 dA = \pi. \end{aligned}$$

Verified!