


## Math 1497 - Sample Test 3 – Solutions

1. Sketch the following parametric curve and find the equation of the tangent at the point of self intersection

$$x = t^3 - 4t + 3, \quad y = t^2, \quad t = -3 \dots 3.$$

*Solution*

From the graph, it appears that they cross at the point (3, 4).



To determine the times where they cross we choose  $y$  (its easier) and set it to 4

$$y = 4 \Rightarrow t^2 = 4 \Rightarrow t = \pm 2.$$

Substituting both  $t = -2$  and  $t = 2$  into  $x$  shows both are 3 so yes, (3, 4) is the point the curve crosses itself. Next we find derivatives

$$\frac{dx}{dt} = 3t^2 - 4, \quad \frac{dy}{dt} = 2t,$$

and dividing gives

$$\frac{dy}{dx} = \frac{2t}{3t^2 - 4}.$$

At  $t = -2$ ,  $\frac{dy}{dx} = -\frac{1}{2}$  and at  $t = 2$ ,  $\frac{dy}{dx} = \frac{1}{2}$ , so the tangents are


$$y - 4 = -\frac{1}{2}(x - 3), \quad y - 4 = \frac{1}{2}(x - 3).$$

2. Graph the following polar equations


$$r = 2 + 2 \sin \theta, \quad r = 2 \sin 2\theta.$$

*Solutions*

$$r = 2 + 2 \sin \theta,$$



$$r = 2 \sin 2\theta,$$



3. Find the area inside one leaf of the rose described by

$$r = 2 \sin 3\theta.$$


*Solution*

Here we use

$$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$$

From the picture below, we find that we sweep out the area when  $\theta = 0 \rightarrow \frac{\pi}{3}$ , so these are the limits of integration. Thus,

$$A = \frac{1}{2} \int_0^{\frac{\pi}{3}} (2 \sin 3\theta)^2 d\theta = \frac{\pi}{3}$$



4. Find the area of the following:

- (i) inside  $r = 2 + 2 \sin \theta$ ,
- (ii) inside the outer loop and outside the inner loop of  $r = 1 - 2 \sin \theta$ ,
- (iii) outside  $r = \cos 2\theta$  and inside  $r = \sin 2\theta$  on  $[0, \frac{\pi}{2}]$ .

*Solutions*

(i)  $r = 2 + 2 \sin \theta$  The picture is above

$$A = \frac{1}{2} \int_0^{2\pi} (2 + 2 \sin \theta)^2 d\theta = 6\pi.$$


(ii) inside the outer loop and outside the inner loop of  $r = 1 - 2 \sin \theta$ ,

$$\begin{aligned} \text{InnerLoop} \quad & \frac{2}{2} \int_{\pi/6}^{\pi/2} (1 - 2 \sin \theta)^2 d\theta = \pi - \frac{3\sqrt{3}}{2} \\ \text{OuterLoop} \quad & \frac{2}{2} \int_{5\pi/6}^{3\pi/2} (1 - 2 \sin \theta)^2 d\theta = 2\pi + \frac{3\sqrt{3}}{2} \\ A = & 2\pi + \frac{3\sqrt{3}}{2} - \left( \pi - \frac{3\sqrt{3}}{2} \right) = \pi + 3\sqrt{3}. \end{aligned}$$

(iii) outside  $r = \cos 2\theta$  and inside  $r = \sin 2\theta$  on  $[0, \frac{\pi}{2}]$ .

In the first quadrant, the curves intersect at  $\theta = \pi/8$  and sweeps out half the area between  $\theta = \pi/8$  and  $\theta = \pi/4$ . The area is given by


$$A = \frac{2}{2} \int_{\pi/8}^{\pi/4} \sin^2 2\theta - \cos^2 2\theta d\theta = \frac{1}{4}.$$



Graphs for 4 (ii) and 4 (iii)

5. Find the projection of the vector  $\vec{u}$  onto  $\vec{v}$  where  $\vec{u} = \langle 2, 3 \rangle$ , and  $\vec{v} = \langle 4, 2 \rangle$ . Sketch both vectors, the projected vector and the orthogonal complement.

In the graph, the vectors  $\vec{u}$  and  $\vec{v}$  are shown



$$\vec{u} \cdot \vec{v} = 8 + 6 = 14, \quad \vec{v} \cdot \vec{v} = 16 + 4 = 20,$$

$$\text{proj}_{\vec{v}} \vec{u} = \left( \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \right) \vec{v} = \frac{7}{10} \langle 4, 2 \rangle$$

The orthogonal complement is given by

$$\vec{u} - \text{proj}_{\vec{v}} \vec{u} = \langle 2, 3 \rangle - \frac{7}{10} \langle 4, 2 \rangle = \left\langle -\frac{4}{5}, \frac{8}{5} \right\rangle.$$

6. (i) Find the equation of the plane that contains the vector  $\langle 1, 2, 4 \rangle$  and the points  $(1, 1, 1)$  and  $(-2, 3, 7)$ .

(ii) Find the equation of the plane that contains the points  $(1, 3, 5)$ ,  $(2, -1, 2)$  and  $(0, 4, 6)$ .

(i) We first construct a vector between the two points, this is  $\langle -3, 2, 6 \rangle$ . Next, cross the two vectors

$$\begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2 & 4 \\ -3 & 2 & 6 \end{vmatrix} = \langle 4, -18, 8 \rangle.$$

The equation of the plane is given by

$$4(x - 1) - 18(y - 1) + 8(z - 1) = 0.$$

or

$$2(x - 1) - 9(y - 1) + 4(z - 1) = 0.$$

(ii) Label the three points  $P(1,3,5)$ ,  $Q(2,-1,2)$  and  $R(0,4,6)$ . Next, we find two vectors that connects two pairs, i.e.  $\vec{PQ} = \langle 1, -4, -3 \rangle$  and  $\vec{PR} = \langle -1, 1, 1 \rangle$ . The cross product will give the normal

$$\vec{n} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -4 & -3 \\ -1 & 1 & 1 \end{vmatrix} = \langle -1, 2, -3 \rangle .$$

The equation of the plane is given by

$$(x - 1) - 2(y - 3) + 3(z - 5) = 0.$$

7. (i) Find the equation of the line that passes through the points  $(1, 2, 4)$  and  $(-2, 3, 7)$ .  
 (ii) Find the equation of the line perpendicular to the plane  $x + 2y - 3z = 6$  passing through the point  $(1, -1, 3)$ .

(i) The line will follow the vector  $\langle -3, 1, 3 \rangle$  so the equation of the line is

$$x = 1 - 3t, \quad y = 2 + t, \quad z = 4 + 3t.$$

and the symmetric form

$$\frac{x - 1}{-3} = \frac{y - 2}{1} = \frac{z - 4}{3}.$$

(ii) The line will follow the normal vector  $\langle 1, 2, -3 \rangle$  so the equation of the line is

$$x = 1 + t, \quad y = -1 + 2t, \quad z = 3 - 3t.$$

and the symmetric form

$$\frac{x - 1}{1} = \frac{y + 1}{2} = \frac{z - 3}{-3}.$$