

Journal of the Marine Biological Association of the United Kingdom

<http://journals.cambridge.org/MBI>


Additional services for *Journal of the Marine Biological Association of the United Kingdom*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Patterns of cleaning behaviour on coral reef fish by the anemoneshrimp *Ancylomenes pedersoni*

Lindsay K. Huebner and Nanette E. Chadwick

Journal of the Marine Biological Association of the United Kingdom / Volume 92 / Issue 07 / November 2012, pp 1557 - 1562
DOI: 10.1017/S0025315411001822, Published online: 06 December 2011

Link to this article: http://journals.cambridge.org/abstract_S0025315411001822

How to cite this article:

Lindsay K. Huebner and Nanette E. Chadwick (2012). Patterns of cleaning behaviour on coral reef fish by the anemoneshrimp *Ancylomenes pedersoni*. Journal of the Marine Biological Association of the United Kingdom, 92, pp 1557-1562 doi:10.1017/S0025315411001822

Request Permissions : [Click here](#)

Patterns of cleaning behaviour on coral reef fish by the anemoneshrimp *Ancylomenes pedersoni*

LINDSAY K. HUEBNER AND NANETTE E. CHADWICK

Department of Biological Sciences, 101 Rouse Life Sciences Building, Auburn University, Auburn, Alabama 36849

*Little is known about the cleaning behaviour of shrimps in comparison to that of cleaner fish, and only recently have cleaner shrimps been shown to remove parasites effectively from coral reef fish. Here we describe patterns of cleaning interactions between Pederson shrimp *Ancylomenes pedersoni* and fish clients in St Thomas, US Virgin Islands. Clients observed here were members of at least 16 fish families, including three previously unreported client families. Most cleans lasted <20 seconds; surgeonfish were cleaned most frequently, but lizardfish and groupers received the longest cleans. The shrimp formed social groups of varying sizes on individuals of the host sea anemone *Bartholomea annulata*, which served as the centres of their cleaning stations. The number and duration of cleans per station increased with the number of resident shrimp, however most anemones hosted small groups of fewer than four individuals, while larger groups of up to nine individuals were relatively rare. Some client fish chased away other fish and competitively excluded them from anemone stations. We conclude that these shrimp clean a wide diversity of clients, vary their clean duration with fish identity, and clean more when in large groups. In addition, clients in part control these patterns of interaction by interfering with access to these stations by other clients.*

Keywords: cleaning symbiosis, cleaner shrimp, *Ancylomenes pedersoni*, client fish, group size, sea anemone

Submitted 16 August 2011; accepted 29 September 2011; first published online 6 December 2011

INTRODUCTION

Cleaner organisms are ecologically important on coral reefs, in that they remove parasites and potentially enhance the healing of surface wounds on client fish (Foster, 1985; reviewed in Côté, 2000). Cleaner fish, primarily wrasses and gobies, have received the most attention from researchers, but a variety of shrimps also appear to function as fish cleaners (Becker & Grutter, 2004). Much of the information on cleaner shrimps is based on anecdotal observations (Limbaugh *et al.*, 1961; Mahnken, 1972), and whether they actually serve as effective cleaners of fish has been a topic of debate (Bunkley-Williams & Williams, 1998; Spotte, 1998; Côté, 2000).

Several recent studies demonstrate that cleaner shrimps can be effective in removing parasites from clients and also clean similar clients as do cleaner fish. Experiments performed under laboratory conditions have shown that two temperate species of *Palaemon* shrimps significantly reduce parasite loads on client plaice (Östlund-Nilsson *et al.*, 2005), *Ancylomenes holthuisi* (Bruce) reduces parasite loads on surgeonfish by 74.5% within two days (Becker & Grutter, 2004) and *A. pedersoni* (Chace) removes 100% of juvenile cymothoid isopods from fish clients (Bunkley-Williams & Williams, 1998). Experiments performed in macrocosms further revealed that *A. pedersoni* significantly reduce the

abundance and size of monogenean parasites on fish (McCammon *et al.*, 2010), and field observations in St Croix indicated that these shrimp also appear to clean a similar diversity of client fish in the Caribbean as do cleaning gobies *Elacatinus evelynae* (Böhlke & Robins) (Johnson & Ruben, 1988). On Red Sea coral reefs, niche overlap of client fish occurs between *A. longicarpus* (Bruce & Svoboda, 1983) and obligate cleaner wrasses *Labroides dimidiatus* (Valenciennes) (Chapuis & Bshary, 2009).

Conversely, some shrimps that initially were classified as cleaners recently have been shown to exert no significant impact on the abundance of certain fish parasites. Bunkley-Williams & Williams (1998) determined that the shrimps *Lysmata grabhami* (Gordon), *Stenopus scutellanus* Rankin and *S. hispidus* (Olivier) do not remove juvenile cymothoid isopods from clients; *S. hispidus* also has no significant effect on monogenean parasites (McCammon *et al.*, 2010). Additionally, *Periclimenes yucatanicus* (Ives) has been described previously as an obligate cleaner (Limbaugh *et al.*, 1961), but extensive field observations have failed to detect fish cleaning by members of this shrimp species (Gwaltney & Brooks, 1994; L.K. Huebner, personal observations), and they have no significant effect on monogenean parasites in a macrocosm setting (McCammon *et al.*, 2010). Further, Wicksten (2009) documented that at least one species of *Lysmata* shrimps, which clean larger fish, will kill and eat smaller fish. These studies demonstrate that quantitative field observations combined with experimental laboratory and macrocosm studies are needed to definitively classify shrimps as cleaners, especially as obligate cleaners.

Corresponding author:

L.K. Huebner

Email: lindsayhuebner@gmail.com

Proven cleaner shrimps on coral reefs vary widely among species in aspects of their biology (Table 1). In contrast to obligate cleaner fish, which perform ‘dances’ and have convergent blue and/or yellow striped body patterns (Potts, 1973; Cheney *et al.*, 2009), cleaner shrimps display neither convergent coloration nor stereotypical guild behaviours toward client fish (Becker *et al.*, 2005; Chapuis & Bshary, 2010). Cleaner shrimps also vary in their microhabitat preferences: some are crevice dwellers, while many are either facultative or obligate associates of sessile invertebrates, especially sea anemones and corals (Table 1). These cnidarian hosts serve as the centre of the shrimps’ cleaning station and can also be posing stimuli for clients (Wicksten, 1995a; Huebner & Chadwick, submitted). Interestingly, cnidarian-associated cleaner shrimp in the genus *Ancylomenes* (formerly within *Periclimenes*; Okuno & Bruce, 2010) all display some variation of transparent bodies with white and/or purple markings, which may serve as signals to clients of this congeneric group of cleaners. This pattern, as well as other similarities and differences among cleaner shrimps in terms of their microhabitat use, behavioural signals, and group cleaning behaviours (Table 1), highlight a need for further investigation into evolutionary and ecological patterns in this group.

Here we present field observations of interactions between client fish and Pederson cleaner shrimp *A. pedersoni* at cleaning stations on corkscrew sea anemones *Bartholomea annulata* (LeSueur) in St Thomas, US Virgin Islands. We identify client fish to family level, and quantify the rates and durations of cleans. We also report various behaviours observed among client fish, as well as include an examination of the effect of shrimp group size on cleaning rates and durations. Finally, we discuss the potential implications of our results for understanding the ecology and evolution of shrimp cleaning symbioses on coral reefs.

MATERIALS AND METHODS

We performed this study at two coral reef sites in St Thomas, US Virgin Islands: Brewers Bay (BB, ~6 m depth, 18°20’N 64°58’W) and Flat Cay (FC, ~7–9 m depth, 18°19’N 64°59’W; site details in Nelsen, 2008). Observations were conducted over about one week at FC in March 2009 only, and at BB in March, July and November 2009, and February 2010. We focused on BB after the initial sampling period, because these reefs are closer to shore, logistically easier to access for dawn observations, and also support a higher abundance of the anemone *Bartholomea annulata* and symbiotic cleaner shrimp *Ancylomenes pedersoni* than do reefs at FC (Nelsen, 2008).

Prior to cleaning observations during each sampling period, we conducted an initial census to identify and map all *B. annulata* in our study areas that each hosted at least one *A. pedersoni* (hereafter termed a cleaning station). We randomly selected a subset of these cleaning stations for observations; the sample size for each field period was determined by the duration of that period and the number of divers available to make observations (N = 15 stations at FC in March 2009, and 17, 16, 22, and 7

Table 1. Information on the six species of cleaner shrimps on coral reefs that have been examined quantitatively in terms of parasite removal ability and/or interactions with clients in the field. Note that *Ancylomenes anthophilus* may be a junior synonym of *A. pedersoni* (Wicksten, 1995b; Spotte, 1999).

| Shrimp | Region | Associate/habitat | Signal | Coloration | Group cleaner? | References |
|----------------------------------------------------|--------------|--------------------------------------------------------------------------------------------------|---------------------------------|-----------------------------------------------------------|--------------------------------------|----------------------------------------------------------------------------------------------------|
| <i>Brachycarpus biunguiculatus</i> (Lucas) | Pantropical | Stations on large sponges; shelter in crevices and/or with <i>Diadema</i> urchins | Not reported | Transparent to bright red-orange | Form pairs but do not clean together | Corredor, 1978; Humann & DeLoach, 2006 |
| <i>Urocaridella</i> sp. c | Indo-Pacific | No associate, crevice-dwelling | Rocking dance | Transparent, red and yellow spots, yellow chelae | Yes | Debelius, 1999; Becker <i>et al.</i> , 2005 |
| <i>Ancylomenes holthuisi</i> (Bruce) | Indo-Pacific | Obligate: various anemones, <i>Cassiopeia andromeda</i> jellyfish, fungiid corals | Not reported | Transparent, white and purple-red spots | Yes | Bruce & Svoboda, 1983 |
| <i>A. longicarpus</i> (Bruce & Svoboda) | Red Sea | Facultative diurnal: various anemones, <i>Xenia</i> soft and <i>Pterogyra</i> stony corals | Cheliped clapping, body swaying | Transparent, small white and violet-blue markings | Yes | Bruce & Svoboda, 1983; Lieske & Myers, 2004; Chadwick <i>et al.</i> , 2008; Chapuis & Bshary, 2010 |
| <i>A. pedersoni</i> (Chace) | Caribbean | Obligate: various anemones, primarily <i>Bartholomea annulata</i> , <i>Condyplactis gigantea</i> | Antennae lashing, body swaying | Transparent, purple lateral stripe, white antennae | Yes | Limbaugh <i>et al.</i> , 1961; Mahnken, 1972; Humann & DeLoach, 2006 |
| <i>A. anthophilus</i> (Holthuis & Eibl-Eibesfeldt) | Bermuda | Obligate: anemones <i>C. gigantea</i> , <i>Actinia bermudensis</i> | Antennae lashing, body swaying | Transparent, purple lateral stripe, white banded antennae | Yes | Sargent & Wagenbach, 1975; Spotte, 1999 |

stations at BB in March, July and November 2009, and February 2010, respectively; N = 77 stations total). Different stations were selected for observation during each of the four field periods.

We conducted cleaning observations in the morning, beginning just after sunrise at ~06:30 and concluding by 08:30, to coincide with the dawn peak in cleaning activity at these shallow reef sites (e.g. Sikkel *et al.*, 2006). Divers observed each cleaning station for 20 minutes from a distance of ~1.5 m or more (20 minutes × 77 stations = 25.67 hours of observation total on *A. pedersoni* stations). Observations began 2–3 minutes after the diver remained stationary on the sea floor, to allow fish to acclimate to diver presence and resume their normal activities. The following data were recorded for each station observed: number of cleaner shrimp at the station, number and identity (to at least family level) of fish clients cleaned, duration (in seconds) of each clean, and client behaviours and interactions. A clean was defined as *A. pedersoni* contact with a fish client body. Because neither the number of *A. pedersoni* per station nor the number or duration of cleans varied among the four sample periods (Kruskal–Wallis tests: $H_3 = 1.736, 6.689, \text{ and } 6.298, P = 0.629, 0.098, \text{ and } 0.083$, respectively), the data were pooled analyses. All data were analysed using SYSTAT 13.

RESULTS

We observed reef fish belonging to at least 14 families to participate in 123 cleans at 77 *Ancylomenes pedersoni* cleaning stations total during our four field periods (four clients were not identified to family level; Figure 1). Though not during our 20 minutes observation periods, we did observe members of Gerreidae (mojarra) and Holocentridae (squirrelfish) during incidental observations at *A. pedersoni* stations during fieldwork for this study; thus *A. pedersoni* clean members of at least 16 fish families at our sites in St Thomas. Acanthurids (surgeonfish) accounted for nearly 50% of fish cleaned; the second most common clients were mullids (goatfish) accounting for 13% of observed cleans. Members of all other families comprised less than 5% each of observed cleans. Despite being the most common clients, acanthurids did not receive the lengthiest cleans (Figure 1). Serranids (groupers) and synodontids (lizardfish) received the longest cleans on average (>120 seconds), while monacanthids (filefish) and mullids had moderately lengthy cleans, averaging 74 seconds. However, variability in the duration of cleans within each of these four families was high. Clean durations ranged from a few seconds up to 11 minutes, but most cleans lasted <20 seconds (Figure 2).

In some cases, after shrimps finished a clean and returned to their host anemones, we observed fish to remain posing at stations, possibly attempting to incite another cleaning bout. Similarly, some fish swam away from stations after the shrimp finished cleaning, but travelled only a few metres or less, and then returned to the same cleaning station. This occasionally resulted in another clean. Also, some fish lingered near stations (<1 m) for several minutes, up to the entire 20 minutes observation period, despite spending little of this time being cleaned.

We observed several types of interactions among fish that approached the cleaning stations. Some fish abandoned




Fig. 1. Variation among client fish families in the number of individuals cleaned and duration of cleans ($\bar{x} \pm SE$) at *Ancylomenes pedersoni* shrimp cleaning stations. Error bars represent variation among individual fish within each family; individuals were not identified to species. The number of individuals observed in each fish family is included in parentheses; a total of 123 cleans were observed. ‘Other’ includes fish that were not identified to family level, but were observed being cleaned; because they were not identified, we did not include durations for these cleans.

clean-seeking behaviour at a station that was already occupied, while others chased away clients that were in the process of receiving cleans. In addition, some clients interrupted their own clean to chase away incoming clients, then returned to the shrimp to continue the clean. These interactions were both inter- and intraspecific: surgeonfish *Acanthurus* spp. chased other *Acanthurus* spp., as well as goatfish and tobacco fish *Serranus tabacarius* (Cuvier), away from cleaning stations. In one such instance, an individual of *S. tabacarius* waited ~1 m from a cleaning station until the surgeonfish departed, then proceeded to pose at the station. One ocean surgeonfish *A. bahianus* Castelnau lingered <0.5 m from a station, and chased all visiting clients from the cleaning station for the duration of the 20 minutes observation, despite spending only 11 seconds being cleaned at the station. Damselfish whose territories overlapped with cleaning stations occasionally chased clients away from stations, consistent with observations of damselfish near cleaning goby stations (Arnal & Côté, 1998).

The number (simple linear regression (SLR): $F_{1,75} = 7.094, P = 0.009, R^2 = 0.086$) and total duration (SLR: $F_{1,75} = 7.775, P = 0.007, R^2 = 0.082$) of cleans per station significantly increased with the number of *A. pedersoni* at that station (Figure 3). However, the duration of individual cleans did not vary significantly with the number of shrimp cleaning at that station ($F_{1,121} = 0.273, P = 0.602, R^2 = 0.002$); thus more shrimp engaged in a clean does not necessarily result




Fig. 2. Variation in clean durations on client fish by *Ancylomenes pedersoni* shrimp.




Fig. 3. Number and duration of all cleans ($\bar{x} \pm \text{SE}$) during 20 minutes observation periods at *Bartholomea annulata* anemone cleaning stations with the number of *Ancylomenes pedersoni* cleaner shrimp per station. Error bars represent variation among stations (i.e. shrimp social groups). Sample sizes of stations observed with each number of shrimp are shown in parentheses. These relationships are significant ($P < 0.01$) in linear regression analyses.

in a shorter clean. One station contained only three shrimp, but 15 cleans occurred there: a large school of surgeonfish visited this station and remained together around the station as members of the school rotated to receive cleans. This station was excluded from these analyses as an outlier. In almost all cleans (97.6%, $N = 123$ cleans), all of the *A. pedersoni* present engaged in the clean, while in only three cleans a subset of the shrimp at the station participated.

DISCUSSION

Individuals of *A. pedersoni* cleaned a wide diversity of fish in St Thomas, US Virgin Islands, including fish in families heretofore unreported to be clients of *A. pedersoni*: Gerreidae, Gobiidae (gobies) and Ostraciidae (boxfish). If we combine our observations with those of other studies (Limbaugh *et al.*, 1961; Mahnken, 1972; Criales, 1979; Johnson & Ruben, 1988; Wicksten, 1995a, 1998), *A. pedersoni* is known to clean fish in 22 families: the 16 reported here (Figure 1) in addition to Aulostomidae (trumpetfish), Balistidae (triggerfish), Bothidae (left-eye flounders), Muraenidae (moray eels), Pomacanthidae (angelfish) and Tetraodontidae (puffer fish). This large diversity of clients is likely because *A. pedersoni* are effective removers of fish parasites (Bunkley-Williams & Williams, 1998; McCammon *et al.*, 2010).

The relative frequencies of client families at our stations differed somewhat from those observed at other localities in the Caribbean. In Bonaire, serranids (~40%) and scarids (parrotfish, ~20%) comprised the majority of clients in 13 families (Wicksten, 1995a), while our most common clients, acanthurids and mullids, represented the third and fourth most common client groups there. In St Croix, fish in only five families were observed at *A. pedersoni* stations (listed in descending frequency): Pomacentridae (damselfish), Serranidae, Scaridae, Holocentridae and Labridae (wrasses; Johnson & Ruben, 1988). These differences may reflect variation in the relative abundances of client fish among reef sites in the Caribbean, or regional variation in cleaner preferences by clients. Due to time constraints, we were not able to quantify relative fish abundances at our field sites, or to estimate the diversity of fish that visited goby stations. However, data from other studies indicate that *A. pedersoni* clean an assemblage of fish clients that overlaps considerably, but not completely, with that of cleaner gobies: fish belonging to 16 of

the 22 client families reported here (excepting Balistidae, Bothidae, Gerreidae, Gobiidae, Monacanthidae and Muraenidae) have been observed visiting cleaner gobies (Johnson & Ruben, 1988; Arnal & Côté, 1998; Côté *et al.*, 1998). However, members of Carangidae (jacks) and Sphyraenidae (barracudas) have been observed at goby stations and not at *A. pedersoni* stations (Johnson & Ruben, 1988).

The large observed variation in the duration of individual cleans, particularly among fish in families with the longest clean durations overall, likely relates in part to client body size. We did not estimate the lengths of fish clients in our study, and clean durations did not correlate significantly with the maximum known sizes of client fish that we observed in each family. However, members of the four fish families with the lengthiest cleans—Monacanthidae, Mullidae, Serranidae and Synodontidae—achieve some of the largest body sizes of the fish observed at these cleaning stations (Humann & DeLoach, 2006). Large fish generally have more parasites (Grutter, 1995), and higher parasite load has been shown to affect cleaning behaviour in some cleaner shrimps (Becker & Grutter, 2005). Thus, these shrimps likely spend more time cleaning large clients because of increased parasite foraging success, as do cleaner wrasse *Labroides dimidiatus* (Grutter, 1995; Bansemer *et al.*, 2002; Bshary & Grutter, 2002). As we only observed one fish to wait for a cleaning station, the presence of waiting fish did not appear to impact cleaning duration in our study. Finally, individual clean durations did not decrease with an increasing number of clients at a station; thus, clean durations, at least in the time span of our observations, appeared to be independent of the number of cleans that a group of shrimp performed. Overall, the durations of cleans observed in this study are similar to those observed by Johnson & Ruben (1988) for several Caribbean cleaners (both fish and shrimp) in nearby St Croix, US Virgin Islands.

The variation in cleans per station (and durations thereof) that we observed when the shrimp occurred in large groups may be due to the small numbers of anemones that hosted >4 *A. pedersoni* (Figure 3). It is not clear why so few anemones contained large groups of shrimp: if the shrimp engage in more cleans when they are part of a large group, why do most of them occur in small groups? It may be that social interactions among the shrimp (Mahnken, 1972), or other dynamics of sea anemone host choice, have a stronger influence on shrimp group size than optimizing their potential number of cleans. Also, these shrimp may not be limited by clients at our study sites and need not optimize cleans by forming large groups. More research is needed on the factors that influence cleaner shrimp social group size on sea anemones.

Client fish may visit stations that host more shrimp because there are more cleaners to inspect their body surface area. It is also possible that shrimp in groups regulate each other's clean quality by minimizing cheating in cooperative cleans, as do cleaner wrasse *L. dimidiatus* that clean in pairs (Bshary *et al.*, 2008). We infrequently observed jolts by client fish, but did not observe client fish to leave or 'punish' *A. pedersoni* after a jolt, suggesting either little to no cheating by these cleaner shrimp and a lack of partner control by the client fish in this system. Though some cleaner shrimp may have a cheating-control system with their clients (Chapuis & Bshary, 2009), *A. pedersoni* may enjoy a system without

punishment, suggesting more similarity to the symbiosis of Caribbean cleaner gobies with clients than to that of the Indo-Pacific cleaner wrasse *L. dimidiatus* (Soares *et al.*, 2008).

This study adds to the information available on cleaner shrimp interactions with client fish in a natural field setting. *Ancylomenes pedersoni* participate in multi-level symbiotic systems, both with client fish and their parasites, and with sea anemones and their other associated symbionts (endosymbiotic microalgae and other crustaceans; Huebner & Chadwick, submitted). The dynamics of how these organisms interact with one another, as well as possible intraspecific interactions among individuals of *A. pedersoni*, suggest a level of complexity in this cleaning system that does not occur with other cleaners that do not form large group sizes on symbiotic hosts. The patterns described here thus provide a foundation for future inquiry into the mechanisms that drive these several types of cleaner shrimp associations: among shrimp that share anemones, between shrimp and client fish, and among the clients themselves. Because cleaner shrimp symbioses are less well understood than those of cleaner fish, these aspects of cleaner shrimp biology likely will be fruitful areas for future research.

ACKNOWLEDGEMENTS

We thank the University of the Virgin Islands (UVI) for the use of their facilities. S. Ratchford at UVI provided insight and field support, and A. Isbell, K. Morrow, and B. and J. Titus contributed valuable field assistance. We thank P. Sikkell, C. Johnston, S. Woodard and three anonymous referees for helpful advice on the manuscript. Funding was provided by a NOAA Puerto Rico SeaGrant (N.E.C., grant number R-101-1-08). This is contribution No. 85 of the Auburn University Marine Biology Program.

REFERENCES

- Arnal C. and Côté I.M. (1998) Interactions between cleaning gobies and territorial damselfish on coral reefs. *Animal Behaviour* 55, 1429–1442.
- Bansemer C., Grutter A.S. and Poulin R. (2002) Geographic variation in the behaviour of the cleaner fish *Labroides dimidiatus* (Labridae). *Ethology* 108, 353–366.
- Becker J.H. and Grutter A.S. (2004) Cleaner shrimp do clean. *Coral Reefs* 23, 515–520.
- Becker J.H.A. and Grutter A.S. (2005) Client fish ectoparasite loads and cleaner shrimp *Urocaridella* sp. c hunger levels affect cleaning behaviour. *Animal Behaviour* 70, 991–996.
- Becker J.H.A., Curtis L.M. and Grutter A.S. (2005) Cleaner shrimp use a rocking dance to advertise cleaning service to clients. *Current Biology* 15, 760–764.
- Bruce A.J. and Svoboda A. (1983) Observations upon some pontonine shrimps from Aqaba, Jordan. *Zoologische Verhandlungen* 205, 3–44.
- Bshary R. and Grutter A.S. (2002) Parasite distribution on client reef fish determines cleaner fish foraging patterns. *Marine Ecology Progress Series* 235, 217–222.
- Bshary R., Grutter A.S., Willener A.S.T. and Leimar O. (2008) Pairs of cooperating cleaner fish provide better service quality than singletons. *Nature* 455, 964–967.
- Bunkley-Williams L. and Williams E.H. Jr (1998) Ability of Pederson cleaner shrimp to remove juveniles of the parasitic cymothoid isopod, *Anilocra haemuli*, from the host. *Crustaceana* 71, 862–869.
- Chadwick N.E., Đuriš Z. and Horká I. (2008) Biodiversity and behavior of shrimps and fishes symbiotic with sea anemones in the Gulf of Aqaba, northern Red Sea. In Por F.D. (ed.) *The Improbable Gulf: history, biodiversity, and protection of the Gulf of Aqaba (Eilat)*. Jerusalem: Magnes Press, Hebrew University, pp. 209–223.
- Chapuis L. and Bshary R. (2009) Strategic adjustment of service quality to client identity in the cleaner shrimp, *Periclimenes longicarpus*. *Animal Behaviour* 78, 455–459.
- Chapuis L. and Bshary R. (2010) Signalling by the cleaner shrimp *Periclimenes longicarpus*. *Animal Behaviour* 79, 645–647.
- Cheney K.L., Grutter A.S., Blomberg S.P. and Marshall N.J. (2009) Blue and yellow signal cleaning behavior in coral reef fishes. *Current Biology* 19, 1283–1287.
- Corredor L. (1978) Notes on the behavior and ecology of the new fish cleaner shrimp *Brachycarpus biunguiculatus* (Lucas) (Decapoda Natantia, Palaemonidae). *Crustaceana* 35, 35–40.
- Côté I.M., Arnal C. and Reynolds J.D. (1998) Variation in posing behaviour among fish species visiting cleaning stations. *Journal of Fish Biology* 53, 256–266.
- Côté I.M. (2000) Evolution and ecology of cleaning symbioses in the sea. *Oceanography and Marine Biology: an Annual Review* 38, 311–355.
- Criales M.M. (1979) Ecology and ethology of the fish cleaner shrimps *Periclimenes pedersoni* Chace and *Lysmata grabhami* (Gordon) in Santa Marta Bay (Colombia). *Acta Científica Venezolana* 30, 570–576.
- Debelius H. (1999) *Crustacea guide of the world*. Frankfurt: Ikan.
- Foster S.A. (1985) Wound healing: a possible role of cleaning stations. *Copeia* 4, 875–880.
- Grutter A.S. (1995) Relationship between cleaning rates and ectoparasite loads in coral reef fishes. *Marine Ecology Progress Series* 118, 51–58.
- Gwaltney C.L. and Brooks W.R. (1994) Host specificity of the anemoneshrimp *Periclimenes pedersoni* and *P. yucatanicus* in the Florida Keys. *Symbiosis* 16, 83–93.
- Humann P. and DeLoach N. (2006) *Reef creature identification: Florida, Caribbean, Bahamas*. 2nd edition. Jacksonville, FL: New World Publications, Inc.
- Johnson W.S. and Ruben P. (1988) Cleaning behavior of *Bodianus rufus*, *Thalassoma bifasciatum*, *Gobiosoma evelynae*, and *Periclimenes pedersoni* along a depth gradient at Salt River Submarine Canyon, St Croix. *Environmental Biology of Fishes* 23, 225–232.
- Lieske E. and Myers R.F. (2004) *Coral reef guide: Red Sea*. London: HarperCollins.
- Limbaugh C., Pederson H. and Chace F.A. Jr (1961) Shrimps that clean fishes. *Bulletin of Marine Science of the Gulf and Caribbean* 11, 237–257.
- Mahnken C. (1972) Observations on cleaner shrimps in the genus *Periclimenes*. *Bulletin of the Natural History Museum of Los Angeles County* 14, 71–83.
- McCammon A., Sikkell P.C. and Nemeth D. (2010) Effects of three Caribbean cleaner shrimps on ectoparasitic monogeneans in a semi-natural environment. *Coral Reefs* 29, 419–426.
- Nelsen M. (2008) *Modeling of population dynamics in the corkscrew anemone Bartholomea annulata on Caribbean coral reefs*. Master's thesis. Auburn University, Auburn, AL, USA.
- Okuno J. and Bruce A.J. (2010) Designation of *Ancylomenes* gen. nov., for the '*Periclimenes aesopius* species group' (Crustacea: Decapoda:

- Palaemonidae), with the description of a new species and a checklist of congeneric species. *Zootaxa* 2372, 85–105.
- Östlund-Nilsson S., Becker J.H.A. and Nilsson G.E. (2005) Shrimps remove ectoparasites from fishes in temperate waters. *Biology Letters* 1, 454–456.
- Potts G.W. (1973) The ethology of *Labroides dimidiatus* (Cuv. & Val.) (Labridae, Pisces) on Aldabra. *Animal Behaviour* 21, 250–291.
- Sargent R.C. and Wagenbach G.E. (1975) Cleaning behavior of the shrimp, *Periclimenes anthophilus* Holthuis and Eibl-Eibesfeldt (Crustacea: Decapoda: Natantia). *Bulletin of Marine Science* 25, 466–472.
- Sikkel P.C., Schaumburg C.S. and Mathenia J.K. (2006) Diel infestation dynamics of gnathiid isopod larvae parasitic on Caribbean reef fish. *Coral Reefs* 25, 683–689.
- Soares M.C., Côté I.M., Cardoso S.C. and Bshary R. (2008) The cleaning goby mutualism: a system without punishment, partner switching or tactile stimulation. *Journal of Zoology* 276, 306–312.
- Spotte S. (1998) “Cleaner” shrimps? *Helgoländer Meeresuntersuchungen* 52, 59–64.
- Spotte S. (1999) Possible synonymy of the Western Atlantic anemone shrimps *Periclimenes pedersoni* and *P. anthophilus* based on morphology. *Bulletin of Marine Science* 65, 407–417.
- Wicksten M.K. (1995a) Associations of fishes and their cleaners on coral reefs of Bonaire, Netherlands Antilles. *Copeia* 2, 477–481.
- Wicksten M.K. (1995b) Within-species variation in *Periclimenes yucatanicus* (Ives), with taxonomic remarks on *P. pedersoni* Chace (Crustacea: Decapoda: Caridea: Palaemonidae). *Proceedings of the Biological Society of Washington* 108, 458–464.
- Wicksten M.K. (1998) Behaviour of cleaners and their client fishes at Bonaire, Netherlands Antilles. *Journal of Natural History* 32, 13–30.
- and
- Wicksten M.K. (2009) Interactions with fishes of five species of *Lysmata* (Decapoda, Caridea, Lysmatidae). *Crustaceana* 82, 1213–1223.

Correspondence should be addressed to:

L.K. Huebner
Department of Biological Sciences
101 Rouse Life Sciences Building
Auburn University, Auburn, Alabama 36849
email: lindsayhuebner@gmail.com