## **Constipation**

- **Diagnosis:** Constipation, although very common in our country, is <u>not</u> normal or healthy. Stooling every day or several times a day <u>doesn't</u> mean you are not constipated. A normal stooling pattern is typically 1 to 3 soft, nonbulky, nonpainful stools per day without apparent blood.
  - : Signs of constipation include
    - abdominal pain
    - pellet-like stools
    - very large/bulky stools
    - clogging the toilet
    - straining while stooling
    - bleeding (including blood noted on the toilet paper, on the stool, or in the toilet water)
    - stool steaking in the diaper or on the underwear ("tire tracks")

**Treatment:** Effective treatment is not complicated, but is long and tedious.

: A proper diet is important, but rarely curative by itself.

- Provide adequate fiber such as **whole** grain breads and cereals, fresh fruits (especially prunes, pears, and apples), and vegetables.
- Avoid <u>excess</u> milk product intake. Limit milk and milk products to 16 ounces per day (24 ounces if 4 to 8 years old, 36 ounces if 9 to 18 years old).
  - \*1 slice of American cheese equals 4 oz. of milk
  - \*1 piece of string cheese equals 6 oz. of milk
  - \*1 oz. of yogurt equals 1 oz. of milk
- Minimize juice, pop, and Gatorade since these are empty calories and it is easy to fill-up on these at the expense of good food.
- : Medications, such as those listed below, are frequently necessary for effective treatment, but

we recommend that that be used only under the direction of a Physician:

- mineral oil senna Metamucil (polyethylene glycol) MiraLax Colace (docusate sodium) Benefiber
- : Sitting on the toilet 4 times per day (breakfast, lunch, dinner, bedtime) and trying to pass a a stool for about 2 minutes is an important part of the retraining process.
- : We will titrate (adjust) the therapy so as to obtain a normal stooling pattern. The medications will usually be continued for at least 6 months prior to <u>weaning</u>. This is necessary so the bowels have enough time to be retrained. **Do not stop the medication abruptly or without talking with us.**
- : Please call us with a progress report in \_\_\_\_\_\_. Also call whenever a problem or question arises. Good communication and consistent treatment is absolutely necessary for a successful outcome.