

What to expect after treatment

Now that you've had your BOTOX® Cosmetic (onabotulinumtoxinA) treatment, here are some reminders:

- Since there's **minimal downtime or recovery**, you should be able to resume your normal activities
- You can expect to see **real, noticeable results**, but you'll still look like yourself
- BOTOX® Cosmetic **starts to work within 24 to 48 hours**¹ – in clinical trials, a majority of people saw significant improvement at 7 days, continuing for up to a month¹
- **Long-lasting:** Proven to last up to 4 months,¹ with a physician-reported response of 25% (102/403) vs 2% (2/128) for placebo¹

Other important information you should know

- Side effects associated with the injection include localized pain, infection, inflammation, tenderness, swelling, redness, and/or bleeding/bruising¹
- **What to avoid:** BOTOX® Cosmetic may cause loss of strength or general muscle weakness, or vision problems within hours to weeks of taking BOTOX® Cosmetic. If any of this happens, do not drive a car, operate machinery, or do other dangerous activities¹

See additional Important Safety Information below.

Be sure to contact your aesthetic provider with any questions or concerns

BOTOX® Cosmetic (onabotulinumtoxinA) Important Information

Approved Use

BOTOX® Cosmetic is a prescription medicine that is injected into muscles and used to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in people 18 to 65 years of age for a short period of time (temporary).

IMPORTANT SAFETY INFORMATION

BOTOX® Cosmetic may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX® Cosmetic:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice (dysphonia), trouble saying words clearly (dysarthria), loss of bladder control, trouble breathing, trouble swallowing. **If this happens, do not drive a car, operate machinery, or do other dangerous activities**

The dose of BOTOX® Cosmetic is not the same as, or comparable to, another botulinum toxin product.

There has not been a confirmed serious case of spread of toxin effect when BOTOX® Cosmetic has been used at the recommended dose to treat frown lines.

Serious and/or immediate allergic reactions have been reported.

They include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Tell your doctor or get medical help right away if you are wheezing or have asthma symptoms, or if you become dizzy or faint.

Please see additional Important Safety Information on reverse side.

BOTOX®
—Cosmetic
onabotulinumtoxinA injection

There's only one BOTOX® Cosmetic

Appointment Reminder

Office name

Office phone number

Next appointment date


onabotulinumtoxinA injection

There's only one BOTOX® Cosmetic

IMPORTANT SAFETY INFORMATION (continued)

Do not take BOTOX® Cosmetic if you: are allergic to any of the ingredients in BOTOX® Cosmetic (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as *Myobloc*® (rimabotulinumtoxinB), *Dysport*® (abobotulinumtoxinA), or *Xeomin*® (incobotulinumtoxinA); have a skin infection at the planned injection site.

Tell your doctor about all your muscle or nerve conditions, such as amyotrophic lateral sclerosis (ALS or Lou Gehrig's disease), myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects including severe dysphagia (difficulty swallowing) and respiratory compromise (difficulty breathing) from typical doses of BOTOX® Cosmetic.

Tell your doctor about all your medical conditions, including: plans to have surgery; had surgery on your face; weakness of forehead muscles, such as trouble raising your eyebrows; drooping eyelids; any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX® Cosmetic can harm your unborn baby); are breast-feeding or plan to breast-feed (it is not known if BOTOX® Cosmetic passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal products. Using BOTOX® Cosmetic with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX® Cosmetic in the past.**

Especially tell your doctor if you: have received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin, such as *Myobloc*®, *Dysport*®, or *Xeomin*® in the past (be sure your doctor knows exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; or take a sleep medicine.

Other side effects of BOTOX® Cosmetic include: dry mouth, discomfort or pain at the injection site, tiredness, headache, neck pain, and eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information, including Boxed Warning and Medication Guide.

1. BOTOX® Cosmetic Prescribing Information, November 2012.