TYLENOL® Professional Dosing Information for Infants & Children

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving TYLENOL®. A healthcare professional should be consulted for dosing in children under the age of two years.



			Your child's weight and age								
Always ask your healthcare professional which product is right for your child.		Age	0-3 mos	4-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs	12 yrs
		Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	95 lbs & ove
Infants Dop Augusta	ape & Cherry dye-free flavo gredient: Acetaminophen 8		0.4mL	0.8mL	1.2mL (0.8mL + 0.4mL)	1.6mL (0.8mL + 0.8mL)	_	-	_	-	-
Cherry Bla Bubblegu dye-free f	n's TYLENOL® Susp ast, Very Berry Strawberry, m Yum, Grape Splash & Che lavors gredient: Acetaminophen 16 mL or 1 tsp)	erry	-	_	_	1 tsp or 5mL	1 ¹ /2 tsp or 7.5mL	2 tsp or 10mL	2 ¹ /2 tsp or 12.5mL	3 tsp or 15mL	-
Childrenk TYLENOL Bubblegui Grape Pur	n's TYLENOL® ay Tablets m Burst, Wacky Watermelo och flavors gredient: Acetaminophen 8 ablet)		-	-	-	2 tablets	3 tablets	4 tablets	5 tablets	6 tablets	-
The Action of th	ENOL [®] Meltaway Tal m Burst & Grape Punch flav gredient: Acetaminophen 16 ablet)	rors	-	_	-	-	- sp = teaspoon*	2 tablets	2 ¹ /2 tablets	3 tablets	4 tablets

Important Instructions for Proper Use

- Read and follow the label instructions on all TYLENOL* products
- Take every 4 hours as needed
- Do not exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Keep all medicines out of the reach of children
- Do NOT administer adult medicines to children

- *Use only the dosing device that comes with a specific product
- Concentrated TYLENOL® Infants' Drops are more concentrated than Children's TYLENOL® Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dropper.
- Children's TYLENOL® Liquids are less concentrated than Concentrated TYLENOL® Infants' Drops. The Children's TYLENOL® Liquids have been specifically designed for use with the enclosed measuring cup.
- Children's TYLENOL® Meltaway Tablets are not the same concentration as Jr. Strength TYLENOL® Meltaway Tablets. Jr. TYLENOL® Meltaway Tablets contain 160mg of acetaminophen while Children's TYLENOL® Meltaway Tablets contain 80mg of acetaminophen.
- If you have any questions, contact your healthcare professional or call 1-877-895-3665

MOTRIN[®] Professional Dosing Information for Infants & Children

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving MOTRIN[®]. A healthcare professional should be consulted for dosing in children under the age of six months.



		Your child's weight and age							
Always ask your healthcare professional which product is right for your child.	Age Weight	0-5 mos 6-11 lbs	6-11 mos 12-17 lbs	12-23 mos 18-23 lbs	2-3 yrs 24-35 lbs	4-5 yrs 36-47 lbs	6-8 yrs 48-59 lbs	9-10 yrs 60-71 lbs	11 yrs 72-95 lbs
Concentrated MOTRIN® Infants' DropsBerry & Berry dye-free flavorsActive Ingredient: (in each 1.25mL)	ıg	-	1.25mL	1.875mL	_	-	-	_	-
Children's MOTRIN® SuspImage: Supple stateImage: Supple state <td>ors</td> <td>-</td> <td>-</td> <td>-</td> <td>1 tsp or 5mL</td> <td>1 ¹/2 tsp or 7.5mL</td> <td>2 tsp or 10mL</td> <td>2 ¹/2 tsp or 12.5mL</td> <td>3 tsp or 15mL</td>	ors	-	-	-	1 tsp or 5mL	1 ¹ /2 tsp or 7.5mL	2 tsp or 10mL	2 ¹ /2 tsp or 12.5mL	3 tsp or 15mL
MOTRIN® Junior Strengt Easy-to-Swallow Caplet Active Ingredient: Ibuprofen 100 (in each caplet) Use only as directed.	5	-	_	-	_	-	2 caplets	2 ¹ /2 caplets	3 caplets
WOTRIN® Junior Strengt Chewable Caplets Orange & Grape flavors Active Ingredient: Ibuprofen 100 (in each tablet)		-	_	_	1 tablet	1 1/2 tablets tsp = teaspoon*	2 tablets	2 ½ tablets	3 tablets
	Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours.								

Important Instructions for Proper Use

- Read and follow the label instructions on all MOTRIN[®] products
- Take every 6-8 hours as needed
- Do not exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or nonprescription)
- Keep all medicines out of the reach of children
- *Use only the dosing device that comes with a specific product
 - Household items such as measuring spoons are less accurate
- Do NOT administer adult medicines to children
- Concentrated MOTRIN[®] Infants' Drops are more concentrated than Children's MOTRIN[®] Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dosing device.
- Children's MOTRIN[®] Liquids are less concentrated than Concentrated MOTRIN[®] Infants' Drops. The Children's MOTRIN[®] Liquids have been specifically designed for use with the enclosed measuring cup.
- The OTC dosing: the recommended dose is 7.5mg/kg every 6-8 hours. The recommended daily dose is 30mg/kg.
- If you have any questions, contact your healthcare professional or call 1-877-895-3665