

RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL TOXICITY and TO GROUND WATER CONTAMINATION

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's Certification.

AgLogicTM 15G Aldicarb Pesticide

For Control of Certain Insects, Mites, and Nematodes

ACTIVE INGREDIENT:

Aldicarb [2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl) oxime]..... 15.0%

OTHER INGREDIENTS: 85.0%

TOTAL: 100.0%

EPA Reg. No. 87895-1

EPA Est. No. 87895-GA-001

KEEP OUT OF REACH OF CHILDREN

DANGER  **POISON**

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with warm water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	

Aldicarb is an N-methyl carbamate.

See inside label booklet for Precautionary Statements, Directions for Use and Storage and Disposal.

MEDICAL INFORMATION

Contact a physician immediately in all cases of suspected poisoning. Illness may be produced rapidly following overexposure to AgLogic 15G. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN: AgLogic 15G is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. AgLogic 15G poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of the administration of parenteral atropine sulfate. Caution should be exercised to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until the patient is fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs such as (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples must be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by AgLogic Chemical LLC.

Consultation on therapy can be obtained at all hours by calling CHEMTREC: 1-800-424-9300.

Manufactured By:

AgLogic Chemical LLC
121 S. Estes Drive, Suite 101
Chapel Hill, NC 27514

NET WEIGHT: _____ POUNDS
LOT NO.: _____

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

DANGER

FATAL IF SWALLOWED. Causes cholinesterase inhibition. May be fatal or harmful by skin or eye contact or by breathing dust. Rapidly absorbed through skin or eyes. Do not get in eyes, on skin or on clothing. Do not breathe dust. Keep away from domestic animals.

Always stand up-wind from hopper when loading.

SIGNS AND SYMPTOMS OF OVEREXPOSURE:

Salivation, Muscle tremor, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache.

In severe cases, convulsions, unconsciousness, and respiratory failure may occur.

PERSONAL PROTECTIVE EQUIPMENT (PPE) FOR HANDLERS NOT USING CLOSED LOADING SYSTEMS

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Personal Protective Equipment (PPE):

All handlers (including loaders and applicators) must wear a minimum of coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear plus socks. In addition, during loading, equipment cleaning or repair, or spill clean up, handlers must wear protective eyewear (goggles or face shield), a chemical-resistant apron, and a NIOSH-approved respirator with a dust/mist filter with the MSHA/NIOSH approval number prefix TC-21C or any N, R, P or HE filter.

Engineering Controls for Enclosed Cab Vehicles:

Applicators using an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] may wear reduced personal protective equipment provided they wear a long-sleeved shirt, long pants, and shoes plus socks and, are provided, have immediately available, and use in an emergency, such as a broken package, spill, or equipment breakdown: chemical-resistant gloves made of any waterproof material, a chemical-resistant apron, chemical-resistant footwear, protective eyewear (goggles or face shield), and a NIOSH-approved respirator with dust/mist filter with the MSHA/NIOSH approval number prefix TC-21C or any N, R, P, or HE filter. Applicators must take off any PPE that was worn in the treated area before reentering the cab, and store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE: This pesticide is extremely toxic to birds and other wildlife. Birds feeding on exposed granules may be killed. Cover or immediately soil incorporate granules spilled during loading, at row ends, or elsewhere to ensure the granules are completely covered with soil.

This pesticide is toxic to fish and aquatic invertebrates. Run-off from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is potentially toxic to honey bees through translocated residues in pollen and nectar if application is made during bloom.

Aldicarb is known to leach through soil into ground water under certain conditions. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Apply this product only as specified on this label. Read the use directions and the appended Environmental Precautions and Soil Type Restrictions Tables prior to making applications. If there are any questions, contact AgLogic Chemical LLC at 1-919-932-5800.

DECOMPOSITION AND MOVEMENT IN SOIL: This product is readily decomposed into harmless residues under most use conditions. However, a combination of permeable and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperatures below 50°F (10°C) at application time tend to reduce degradation and promote movement of residues to ground water. If the above describes your local use conditions and ground water in your area used for drinking, do not use this product without first contacting AgLogic Chemical LLC at 1-919-932-5800.

NOTICE: Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

This Act protects Attwater's Greater Prairie Chicken in the Texas counties of Aransas, Austin, Brazoria, Colorado, Galveston, Goliad, Harris, Refugio, and Victoria.

Prior to making applications in these counties, the user must determine that this species is not located in or immediately adjacent to the area to be treated. If the user is in doubt whether or not the above named endangered species may be affected, he should contact either the regional U.S. Fish & Wildlife Service office (Endangered Species Specialist) or personnel of the State Fish and Game office.

DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Exception: If the product is soil-incorporated or soil-injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Once the REI has expired, do not allow persons to come into direct contact with treated wet soil as the result of the initial irrigation or rainfall after treatment unless they are wearing the PPE specified below for early entry. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls worn over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks, and protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store unused AgLogic 15G in original container only, in secure, well ventilated clean dry area out of reach of children and animals. Do not store in areas where temperature averages 115°F (46°C) or greater. Do not store in or around the home or home garden.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If container is burned, stay away from and do not breathe or contact smoke.

IN CASE OF TRANSPORTATION OR WAREHOUSE EMERGENCY INVOLVING A SPILL, FIRE, OR EXPOSURE, CALL CHEMTREC 1-800-424-9300 TWENTY-FOUR HOURS A DAY IN THE USA.

INFORMATION

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

AgLogic 15G controls listed insects, mites, and nematodes. When applied into moist soil at planting and/or postemergence the active ingredient aldicarb is rapidly absorbed by roots and translocated to all parts of the plant. Rainfall or irrigation soon after application will ensure prompt uptake of aldicarb; however, if irrigation is necessary, care should be taken not to over irrigate to reduce the potential for residues leaching to ground water. Control often lasts more than six weeks varying with growing conditions, rate of use, and pests. Crop yields are usually increased with treatments of AgLogic 15G.

RESTRICTIONS

Use AgLogic 15G only in accordance with label directions, and restrictions. **DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING.** Use higher rates on heavy organic or clay soils. Do not exceed the maximum label rate.

Application may only be made using motorized ground application equipment. Application using aircraft, backpack spreaders, or push-type spreaders is prohibited. Additional restrictions on application equipment pertain to use on sweet potatoes.

Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, AgLogic 15G and water must be on the same side of the plant row.

Calibrate and adjust ground application equipment to ensure proper rate and accurate placement. Clean application equipment thoroughly after use. For any leftover material, see instructions for STORAGE AND DISPOSAL in this booklet.

Deep disk any spills at row ends immediately to prevent birds from feeding on exposed granules.

GROUND WATER RESTRICTIONS

- Observe Environmental Hazard statements regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since over-irrigation promotes the leaching of chemicals.
- **Do not apply within 50 feet of any drinking water well. More restrictive setbacks may apply. See the *Environmental Precautions and Soil Type Restriction Tables*.**
- Do not wash, load or empty application equipment near any well, as this practice is a potential source of ground water contamination.
- Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.
- For State Specific Ground water Restrictions, see the *Environmental Precautions and Soil Type Restriction Tables*.
- Apply only between March 1 and September 1 when used in California.

ROTATIONAL CROPS RESTRICTIONS

Do not plant any crop not listed on this label in soil treated with AgLogic 15G within 10 months after the last application with the following exceptions:

Six-month plantback restriction

- Do not plant wheat or barley within 6 months after last application.
- Do not plant bulb crops (such as onions or garlic) within 6 months after the last application.
- Do not plant brassica crops (such as broccoli or cabbage) within 6 months after the last application.

Eight-month plantback restriction

- Do not plant corn within 8 months of last application.
- Do not plant melons within 8 months after last application.
- Do not plant other cucurbits (such as cucumbers and squash) within 8 months after last application.
- Do not plant fruiting vegetables (such as tomatoes or eggplant) within 8 months after the last application.

MEAT AND MILK

- Do not allow livestock to graze in treated areas before harvest.

COMPATIBILITY

Pesticidal activity of AgLogic 15G is not affected by normal applications of fertilizers or other pesticides. Its effectiveness may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix AgLogic 15G with other materials before application.

DIRECTIONS FOR USE

To provide maximum performance and to minimize hazard to birds, granules must be placed into bottom of furrow and immediately covered with soil by mechanical means. For all applications, cover or immediately soil incorporate granules spilled during loading, at row ends, or elsewhere to ensure the granules are completely covered with soil. When a range of rates are specified, use the higher rate if pest infestations are expected to be severe.

COTTON

Crop & Time Of Application	Pests Controlled	Pounds /Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions	
<p>COTTON At planting</p> <p>In the States of AL, FL, GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	Aphids, Thrips	3.5 to 5 (Except Texas, Oklahoma and New Mexico)	40" row spacing 4.5 to 6	<p>Apply granules in the seed furrow and immediately cover with soil by mechanical means.</p> <p>If seeds and AgLogic 15G are hill-dropped, AgLogic 15G rates may be reduced by one-half.</p>	
	Fleahoppers, Leafminers, Mites, Overwintering boll weevil (adults feeding on foliage), Plant bugs, including Lygus	2 to 5 (Texas, Oklahoma, and New Mexico Only)	2.5 to 6		
		4 to 7	5 to 8.5		
	Nematodes	3.5 to 7 (Except California)	4.5 to 8.5	<p>Apply granules in seed furrow and immediately cover with soil by mechanical means.</p> <p>OR</p> <p>Apply granules in a 4 to 6-inch band (T-Band) over open seed furrow and immediately cover with soil by mechanical means.</p>	
	Aphids, Thrips, Mites, Nematode Suppression (California Only)	5 to 7 (California Only)	6 to 8.5	Apply in seed furrow and cover with soil.	
<p>Side Dress Application Only or Split Application</p> <p>From 3 weeks after planting through first squaring</p> <p>Side Dress Application only</p>	Suppression of Leafhoppers, Fleahoppers, Mites, Lygus and Nematodes	5 (except California)	6	Side dress granules in a furrow that is 6 to 10 inches to one or both sides of plant row to a depth of 2 to 3 inches. Adjust applications to minimize root pruning.	
<p>In the States of AL, FL, GA and SC, if a Side Dress application only is planned and if a vulnerable soil is</p>	Suppression of Leafhoppers, Fleahoppers, Mites, Boll weevil, Plant	14 (California only)	17 (California only)	Apply before populations reach an economic threshold. Apply for control of moderate populations. Only suppression may be expected for heavy populations of sweet potato	

Crop & Time Of Application	Pests Controlled	Pounds /Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
<p>present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>Split Application</p> <p>In the States of AL, FL, GA and SC, if an At Planting and Side Dress application is planned and if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	<p>bugs including Lygus, Cotton leaf perforator, Whiteflies and Nematodes (suppression) . California only.</p>			<p>whitefly, especially the poinsettia (B) strain (silver leaf).</p>

RESTRICTIONS FOR COTTON

- The maximum single at-plant application rate is 7 pounds per acre. See application restrictions on vulnerable soils.
- The maximum single side-dress application rate is 14 pounds per acre in California and 5 pounds per acre in other states. See application restrictions on vulnerable soils.
- Do not exceed a total of 21 pounds per acre per year in California or 12 pounds per acre per year in other states for all applications to cotton. See application restrictions on vulnerable soils where a combination of an At-Planting and Side Dress application is planned.
- Apply only between March 1 and September 1 when used in California.
- Do not make more than one at-planting application and one postemergence application per crop per year.
- Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning.
- Do not apply within 90 days of harvest.
- Do not feed cotton forage to livestock or allow livestock to graze in treated area.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

DRY BEANS

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
DRY BEANS At planting FOR USE ONLY IN: Colorado, Oregon, Washington, Idaho, and Michigan	Seedcorn maggot	3.5 to 5	36 inch <u>row spacing</u> 4.0 to 5.5	Apply granules in seed furrow and immediately cover with soil by mechanical means.
	(Michigan Only)			OR
	Aphids	5 to 7	5.5 to 7.5	Apply in a furrow that is 2 to 3 inches to side of seed row and 2 to 3 inches deep and immediately cover with soil by mechanical means.
	Leafhoppers Mexican bean beetle Mites	7 to 14	7.5 to 15	OR Granules can be placed in seed furrow if the rate does not exceed 5.5 ounces per 1000 feet of row (5 pounds per acre on 36 inch rows).
	Nematodes	7 to 14	7.5 to 15	Apply granules in a 4 to 6 inch band and immediately cover with soil by mechanical means. Plant into treated zone. OR Where furrow irrigation is used, apply granules 3 to 4 inches deep and 3 inches from seed row on the water furrow side.

RESTRICTIONS FOR DRY BEANS

- For use only in Colorado, Oregon, Washington, Idaho, and Michigan.
- Do not exceed a total of 14 pounds per acre per season.
- Treatments in excess of 5.5 ounces per 1000 feet of row (5 pounds per acre on 36 inch rows) made directly to the seed furrow may delay plant emergence and reduce plant stand.

- Do not make more than one application per crop per year.
- Do not harvest within 90 days of treatment.
- Do not feed green forage, hay or straw to livestock.
- Do not allow livestock to graze in treated areas before harvest.
- Do not use green pods as food for humans.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

PEANUTS

Crop & Time Of Application	Pests Controlled	Pounds /Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
<p>PEANUTS At planting</p> <p>In the States of AL, FL, GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	Thrips	7	36" <u>row spacing</u> 7.5	Apply granules in seed furrow and immediately cover with soil by mechanical means. In Southwest United States use high rate only.
<p>Split Application (Alabama, Florida, Georgia, North Carolina, Oklahoma, Texas and Virginia Only)</p> <p>At time of planting</p> <p>and/or</p> <p>Post-emergence (pegging application) at or just prior to peg initiation but no later than 40 days after emergence and prior to last cultivation.</p> <p>In the States of AL, FL, GA and SC, if an At Planting and/or a Pegging application is planned</p>	Nematodes	7 At-planting	7.5 At-planting	<p>At-Planting: Apply granules in a 4 to 6 inch band over open seed furrow (T-band) and immediately cover with soil by mechanical means.</p> <p>OR</p> <p>Apply granules in a 6 to 12 inch band and immediately cover with soil by mechanical means to a depth of 2 to 4 inches. Plant seed into treated zone.</p> <p>Post-Emergence:</p> <p>Apply granules in a band 12 to 18 inches wide on the row and into the plant canopy. Ensure that plant foliage is dry prior to application. Dislodge granules from foliage by suitable means that will not damage the plant.</p>
		10 Post-emergence	11 Post-emergence	

Crop & Time Of Application	Pests Controlled	Pounds /Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
and if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.				Irrigate immediately after application and complete within 24 hours unless rainfall is received.

RESTRICTIONS FOR PEANUTS

- Do not make more than one application per crop per year in states other than Alabama, Florida, Georgia, North Carolina, Oklahoma, Texas and Virginia.
- The maximum single At-Plant application rate is 7 pounds per acre. See application restrictions on vulnerable soils.
- The maximum single Pegging application rate is 10 pounds per acre. See application restrictions on vulnerable soils.
- Do not exceed a total of 17 pounds per acre per year for all applications to peanuts. See application restrictions to vulnerable soils where a combination of an At-Planting and Pegging application is planned.
- Post-emergence applications are permitted only in fields where overhead irrigation is available.
- Do not make the split application to Spanish peanuts or other short season varieties (a minimum of 90 days is required between pegging applications and harvest).
- Do not harvest within 90 days of application.
- Do not hog-off treated fields.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed hay or vines to livestock.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

SOYBEANS

Crop & Time Of Application	Pests Controlled	Pounds /Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
<p>SOYBEANS At planting</p> <p>FOR USE ONLY IN: Georgia, North Carolina, South Carolina, and Virginia</p> <p>In the States of GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>[NB: No split or pegging language]</p>	<p>Mexican bean beetle Thrips Suppression of three cornered alfalfa hopper</p>	<p>5 to 7</p>	<p>30" <u>row spacing</u> 4.6 to 6.4</p>	<p>Apply granules in seed furrow and immediately cover with soil by mechanical means.</p> <p>Granules may be applied in seed furrow if rate does not exceed 5.5 ounces per 1000 feet of row OR If rate exceeds 5.5 ounces per 1000 feet of row (6 pounds per acre on 30 inch rows), apply a 4 to 6 inch band over open seed furrow and immediately cover with soil by mechanical means.</p>
	<p>Nematodes</p>	<p>7</p>	<p>6.4</p>	<p>Apply a 4 to 6 inch band (T-Band) over open seed furrow and immediately cover with soil by mechanical means.</p>

RESTRICTIONS FOR SOYBEANS

- For use only in Georgia, North Carolina, South Carolina, and Virginia.
- Do not make more than one application per crop per year.
- Do not harvest within 90 days of treatment.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed green forage, hay, or straw to livestock.
- Treatments in excess of 5.5 oz per 1000 feet of row (6 pounds per acre on 30 inch rows) made directly to the seed furrow may delay plant emergence and reduce plant stand.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

SUGAR BEETS

Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
SUGAR BEETS At planting or within one week before planting. FOR USE ONLY IN: California, Colorado, Idaho, Montana, Nebraska, Oregon, Washington and Wyoming.	Nematodes	27 to 33 (Except California)	22" <u>row spacing</u> 18-22	Apply granules in a 4 to 6 inch band and immediately cover with soil by mechanical means. Plant seed into or above treated zone; OR Where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
		14 (California Only)	9.5	
At planting	Aphids	7 to 14	4.5 to 9.5	Drill granules 1 to 3 inches below seedline. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre.
	Leafminers Leafhoppers	14 to 20 (Except California)	9.5 to 13.5	
		14 (California Only)	9.5	
Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules in a 2 to 3 inch band over seed row and immediately cover with soil by mechanical means. OR Where furrow irrigation is employed for seed germination, drill granules 2 inches deep and 2 inches from seed row on water furrow side. For the 7 lb. rate, apply granules in a 1 to 2-inch band in front of the press wheel as the furrow is closing.	
At planting plus postemergence (split applications)	Nematodes	14 to 20 at planting and 14 to 20 at postemergence (Except California)	9.5 to 13.5 at planting and 9.5 to 13.5 at postemergence	AT PLANTING: Apply granules in a 4 to 6-inch band and immediately cover with soil by mechanical means. Plant seed into or above treated zone; OR Where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches
		14 at planting and 14 at postemergence (California Only)	9.5 at planting and 9.5 at postemergence	

Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
				from seed row on water furrow side. POSTEMERGENCE: Apply granules to both sides of plant row and immediately cover with soil by mechanical means OR Where furrow irrigation is employed side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.
Postemergence Do not make any postemergence application if 27 to 33 lbs./acre were applied at planting or one week before planting. Do not make more than one at planting application and two postemergence applications per crop.	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil by mechanical means OR Where furrow irrigation is employed side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.
	Aphids	7 to 14	4.5 to 9.5	Apply as above. A repeat application may be required for continued protection against virus vectors (aphids, leafhoppers). Apply within 60 days of planting.
	Leafminers Leafhoppers	14 to 20 (Except California)	9.5 to 13.5	
		14 (California Only)	9.5	
	Nematodes	27 (Except California)	18	Apply as above. Apply within 60 days after planting. In California, apply within 30 days after planting.
14 (California Only)		9.5		

RESTRICTIONS FOR SUGAR BEETS

- For use only in California, Colorado, Idaho, Montana, Nebraska, Oregon, Washington and Wyoming.
- Apply only between March 1 and September 1 when used in California.
- Do not exceed a total of 28 pounds per acre per year in California and 33 pounds per acre per year in other states for all applications to sugar beets.
- Do not make more than one at planting application and two postemergence applications per crop per year.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.
- Treatments in excess of 4.5 ounces per 1000 feet of row (7 pounds per acre on 22 inch rows) made directly in the seed furrow may delay plant emergence and reduce plant stand.

- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

SWEET POTATOES

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
SWEET POTATOES Pre-plant or at planting FOR USE ONLY IN: Louisiana and Mississippi	Nematodes	10 to 20	<u>48"</u> <u>row spacing</u> 15 to 30	Apply granules in a 12-inch band over open furrow or soil surface and cover immediately during bed forming by mechanically hilling up 8 to 10 inches. Place transplant in center of treated zone.

REQUIRED APPLICATION EQUIPMENT FOR SWEET POTATOES

All applications must be made with Positive Displacement Applicator such as Horstine Microband® or Gandy PDM® or with other Positive Displacement Applicators.

RESTRICTIONS FOR SWEET POTATOES

- For use only in Louisiana and Mississippi.
- Maximum single application rate is 20 pounds per acre per year.
- Do not make more than one application per crop per year.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

PLANT GROWTH STIMULATION: In the absence of recognizable target pests and under certain growing conditions, stimulation of plant growth by AgLogic 15G on certain crops has been demonstrated under laboratory, greenhouse, and field conditions. Plants may be visibly taller, greener, and denser. Faster grow-off and an increase in fruiting rate and size often are results of the growth enhancement by AgLogic 15G. However, such effects do not occur with all crops or under all conditions. Therefore, this product should not be used solely as a plant growth regulator, and its effect as one should be considered a side benefit which may or may not occur as a consequence of its use.

LIMIT OF WARRANTY AND LIABILITY

AgLogic Chemical LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

To the extent consistent with applicable law, buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise. To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL TOXICITY and TO GROUND WATER CONTAMINATION

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's Certification.

AgLogic 15G Aldicarb Pesticide

EPA Reg. No. 87895-

Environmental Precautions and Soil Type Restriction Tables

Refer to the container label for additional use precautions and directions.

Ag Logic LLC
121 S. Estes Drive, Suite 101
Chapel Hill, NC 27514

ENVIRONMENTAL PRECAUTIONS

GENERAL USE RESTRICTIONS

AGRICULTURAL CHEMICALS HAVE THE POTENTIAL TO MOVE INTO SHALLOW GROUND WATER.

THE FOLLOWING RESTRICTIONS HAVE BEEN DEVELOPED TO PROTECT DRINKING WATER SUPPLIES.

DO NOT APPLY WITHIN 50 FEET OF ANY DRINKING WATER WELL TO MINIMIZE CONTAMINATION BY SURFACE RUNOFF.

MORE STRINGENT RESTRICTIONS MAY BE REQUIRED, AS DISCUSSED BELOW.

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination. In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.

Contact your state pesticide regulatory authority for further information on state requirements for the use of this product.

Some states have or may be developing more restrictive regulations regarding the use and application of AgLogic 15G. Follow all state regulations restricting the use and application of this product, including limitations on applications near drinking water sources. In all cases, the more restrictive requirements must be followed. It is the responsibility of the applicator to document the construction of wells claimed not to be shallow.

STATE SPECIFIC GROUND WATER LIMITATIONS

Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.

FOLLOW THE LISTED "ADDITIONAL RESTRICTIONS" if the following conditions are present:

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS
CO, DE, KS, KY, LA, MD, MO, MS, NC, TN, VA, WV	Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight. See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.
IA, IL, IN, MI, MN, MT, ND, NE, OH, SD, WY	Sandy loam, loamy sand, or sand surface soils, and loamy sand or sand subsoils, with an average organic matter in the upper 12 inches of less than 2% by weight. See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.
AL, GA, SC	Vulnerable Cotton and Peanut Soils in AL, GA and SC and Vulnerable Soybean Soils in GA and SC. Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	<p>At planting application only to cotton, peanuts or soybeans</p> <p>If AgLogic 15G is applied to cotton, peanuts or soybeans as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>Side Dress Application Only or Split Application to cotton</p> <p>If AgLogic 15G is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>If AgLogic 15G is applied to cotton as an At-Plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>At-Planting and/or Post emergence application to peanuts</p> <p>If AgLogic 15G is applied to peanut as an At-Plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Do not make a post emergence application to peanuts in South Carolina.</p>
FL	Vulnerable Peanut and Cotton Soils Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight. Any soil classified with a soil hydrologic group of C or D is not considered a restricted soil regardless of other properties.	<p>At planting application only.</p> <p>If AgLogic 15G is applied At-planting to peanuts or cotton and a vulnerable soil is present (see soils listed below) and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.</p> <p>Side Dress Application Only or Split Application to cotton</p> <p>If AgLogic 15G is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet</p>

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS							
		<p>below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>If AgLogic 15G is applied to cotton as an At-Plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>At-Planting and/or Post emergence application to peanuts</p> <p>If AgLogic 15G is applied to peanut as an At-Plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Do not make a post emergence application to peanuts in South Carolina.</p> <p>Additional Information</p> <p>In Florida, it is the responsibility of the applicator to document the construction of wells claimed not to be shallow. This must consist of: (a) a copy of the well completion report issued by the appropriate water management district; or (b) a statement certified as to accuracy by a licensed well contractor. The U.S.D.A. Natural Resources Conservation Service which serves your county can tell you if the soils on your farm fall within the following types of vulnerable soils. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. See Section 5E-2, 028, Florida Administrative Code (F.A.C.), for additional restrictions.</p> <table border="1" data-bbox="753 995 1528 1163"> <tr> <td data-bbox="753 995 899 1163">Adamsville Alaga Alpin Archbold Astatula Bigbee Candler</td> <td data-bbox="899 995 1062 1163">Eglin Florahome Fort Meade Foxworth Gainesville Kershaw Klej</td> <td data-bbox="1062 995 1224 1163">Lake Lakeland Neihurst Orlando Orsino Ortega Paola</td> <td data-bbox="1224 995 1528 1163">Penney Quartzipsamments Satellite Tavares Valdosta</td> </tr> </table>				Adamsville Alaga Alpin Archbold Astatula Bigbee Candler	Eglin Florahome Fort Meade Foxworth Gainesville Kershaw Klej	Lake Lakeland Neihurst Orlando Orsino Ortega Paola	Penney Quartzipsamments Satellite Tavares Valdosta
Adamsville Alaga Alpin Archbold Astatula Bigbee Candler	Eglin Florahome Fort Meade Foxworth Gainesville Kershaw Klej	Lake Lakeland Neihurst Orlando Orsino Ortega Paola	Penney Quartzipsamments Satellite Tavares Valdosta						
FL	Other Soils, All Crops	<p>State regulations require that AgLogic 15G not be applied within 300 feet of any drinking water well. See Section 5E-2, 028, F.A.C., and additional restrictions for peanut and cotton above.</p>							

SOIL TYPE RESTRICTION TABLE

FOLLOW THE LISTED ADDITIONAL RESTRICTIONS IF THE SOIL TYPES ARE PRESENT IN YOUR STATE:

ALABAMA					
<p>If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. At planting application only: if AgLogic 15G is applied to cotton or peanuts as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Side Dress Application Only: If AgLogic 15G is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Split Application: If AgLogic 15G is applied to cotton as an At-Plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. At-Planting and/or Post emergence application to peanuts: If AgLogic 15G is applied to peanut as an At-Plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
alaga alpin americus bassfield bigbee	blanton bruno buncombe chiplely chipola	corolla crevasse duckston eustis flomaton	foxworth fripp garcon gorgas jones	lakeland latonia leon nugent osier	pactolus plummer saffell scranton tarboro
COLORADO					
<p>If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
bangston bankard barcus baroid batterson bijou blakeland boel brad breece bresser canlodore cascajo chappell chaseville clark fork columbine comad corlena corlett costilla	cotopaxi coyet crestman crustown dailey dix dunday dunul dwyer eachuston ecklund eghalm ellicott els elsmere falcon farb farisita gilcrest gracot gretdivid	grieves grimm grimstone gunbarrel hiwan inavale ipage ironsprings juget julesburg kandaly kassler kerber kettle kippen laird laney las animas layoint legault lincoln	littlebear lonetree luning mathis maybell medano mespun mido mirror lake moenkopie moosed mosca munjor mysten nakai nesda newcomb osgood ouray pando patna	peetz pescar pineguest platte redcreek resort roswell ruedloff ruhe ryark san isabel sawcreek schamber schooner sheppard siebert southace space city sphinx spool stapleton	stecum tassel thoroughfare tipper tipperary tivoli tomah tomichi trail tricera tullock valent valentine wigton willwood yetull zeona
DELAWARE					
<p>If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
downer evesboro fort mott	galestown hammonton hooksan	hurlock kenansville klej	osier plummer pocomoke	rumford	

SOIL TYPE RESTRICTION TABLE (continued)

GEORGIA					
<p>If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. At planting application only: if AgLogic 15G is applied to cotton, peanuts or soybeans as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Side Dress Application Only: If AgLogic 15G is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Split Application: If AgLogic 15G is applied to cotton as an At-Plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. At-Planting and/or Post emergence application to peanuts: If AgLogic 15G is applied to peanut as an At-Plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
americus bigbee blanton boulogne buncombe cainhoy centenary	chiple chipola duckston echaw eustis flomaton foxworth	fripp gorgas hurricane johns kershaw kingsferry kureb	lakeland leon lynn haven mandarin meadowbrook molena osier	ousley penney plummer pottsburg ridgeland ridgewood sapelo	scranton valdosta wateree
ILLINOIS					
<p>If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
algansee alvin beavercreek billet bloomfield boone boyer brems burkhardt	carmi carr casco chelsea chute coloma dickinson disco el dara	eleva flagler grable granby hodge homer hononegah hoopeton lamont	landes lanier lorenzo matherton minneiska morocco oakville oshtemo plainbo	plainfield princeton roby rodman saffell sarpy seafield sparta spinks	stonelick thetford watseka zumbro
INDIANA					
<p>If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
abscola algansee alvin billet bloomfield bobtown boyer brady branch brems bronson	bruno carmi casco chatterton chelsea coloma conotton conrad desker dickinson elston	granby gravelton hanna homer hononegah hoopeton junius kosciusko landes leoni maumee	morocco moundhaven nesius newton oakville ormas oshtemo ouiatenon plainfield princeton prochaska	riverdale roby rodman saugatuck seafield shipshe sisson sparta spinks stockland stonelick	tedrow thetford tyner wasepi watseka zaborosky zadog

SOIL TYPE RESTRICTION TABLE (continued)

IOWA					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
backbone billet boone brady buckney burkhardt	carr chelsea chute dickinson dickman elrick	estherville finchford flagler fruitfield granby hawick	hoopston klum lamont lands lilah montieth	oesterle perks salida sarpy sparta toolesboro	watseka zenor
KANSAS					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
aline bankard dillwyn dix	dwyer els elsmere goltry	goodnight gracemore haxtun inavale	kanza las animas likes lincoln	pratt sarpy schamber simeon	thurman tivoli valent valentine
KENTUCKY					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
bruno crevasse	flomaton iuka	lakin molena	potomac saffell	smithdale	
LOUISIANA					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alaga bassfield betis bienville	bigbee bruno cheniere crevasse	darden eustis felicity flo	hackberry haggerty hainesville kenney	lakeland latonia lotus nugent	osier palm beach peveto
MARYLAND					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
aura carteret colts neck downer dragston	duckston evesboro fisherman fort mott fripp	galestown hammorton hooksan hurlock ingleside	klej lakeland leetonia leon molena	newhan osier plummer pocomoke remlik	runclint

SOIL TYPE RESTRICTION TABLE (continued)

MICHIGAN					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
abbaye	chesaning	granby	mahtomedi	padus	spinks
abscota	coloma	grattan	mancelona	pelkie	springlake
adams	covert	graycalm	manitowish	pemene	st. ignace
alcona	crosswell	grayling	matherton	pence	steuben
algansee	cunard	guardlake	maumee	pequaming	sturgeon
allouez	deer park	halfaday	mecosta	perrin	summerville
alpena	deerton	hodenpyl	menahga	peshekee	sundog
alvin	desker	ingalls	millersburg	pipestone	tedrow
arcadian	dickinson	ionia	monico	plainfield	tekenink
arkport	dixboro	ishpeming	montcalm	rapson	thetford
au gres	duel	kalkaska	morocco	richter	trimountain
banat	dunbridge	karlin	nadeau	ridgeville	tyre
battlefield	east lake	kawbawgam	newton	riverdale	velvet
bixler	eastport	keweenaw	nottawa	rodman	vilas
blue lake	eleva	kingsville	oakville	rousseau	wainola
bohemian	elston	kiva	oconto	rubicon	waiska
boyer	ensign	klacking	ocqueoc	sarona	wallace
brady	esau	koontz	omega	saugatuck	wasepi
branch	evart	kosciusko	onota	sayner	watseka
brems	fabius	landes	ormas	seafield	winterfield
bronson	finch	leelanau	oshtemo	shawano	zeba
channing	freesoil	leoni	otisco	shelldrake	zimmerman
chatham	gilchrist	liming	ottokee	sisson	
chelsea	gladwin	lode	paavola	sparta	
MINNESOTA					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
abscota	chelsea	finchford	langola	omega	soderville
alban	chetek	flagler	lasa	osakis	sparta
algansee	claire	fossum	leola	pierz	spinks
alvin	cloquet	friendship	lilah	plainbo	sugarbush
andrusia	conic	gotham	lino	plainfield	torning
anoka	copaston	granby	lohnes	poppleton	ulen
arvilla	cormant	graycalm	maddock	quetico	venlo
beavercreek	cowhorn	hamar	mahtomedi	redby	verndale
becker	cromwell	hawick	marquette	renshaw	wamduska
bellchester	deer park	hecia	meehan	rosholt	watseka
bergekeller	dickinson	hiwood	menahga	rosy	waupaca
billet	dickman	hoopeston	mesaba	salida	wawina
boone	egeland	insula	minneiska	sanburn	winterfield
burkhardt	eleva	kanaranzi	nemadji	sartell	zenor
burnsville	emmert	karlstad	northfield	serden	zimmerman
carmi	evart	kost	nymore	shawano	zippel
caryville	faunce	lamont	oesterle	sioux	zumbro
MISSISSIPPI					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alaga	bigbee	eustis	latonia	pactolus	st. lucie
bassfield	bruno	flomaton	nugent	plummer	
beulah	crevasse	lakeland	osier	saffell	
MISSOURI					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alvin	carr	eustis	maldden	scotco	
beulah	clana	finchford	plainfield	shelldrake	
bruno	crevasse	hodge	sandbur	sparta	
canalou	diehlstadt	landes	sarpy	wideman	

SOIL TYPE RESTRICTION TABLE (continued)

MONTANA					
If a vulnerable soil is present and the water table is less than 25, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alberton	calicott	glendive	mirror	sachet	trey
ambrant	castner	hanly	moiese	sawcreek	tulloch
apeldorn	cheadle	haskill	nelson	scravo	turnercrest
ashbon	cheyenne	haverdad	nemote	selle	tusier
ashuelot	chinook	haverly	nesda	selway	twilight
assinniboine	clark fork	havrelon	nippt	seroco	upsata
bangston	cohagen	hiwan	oceanet	sheege	usine
bankard	colburn	jugson	oraid	shingle	utica
banks	comad	kalsted	ovando	shook	valentine
baxendale	como	kirby	pend orielle	shurley	vebar
bearmouth	cowood	krause	poin	splitro	victor
beaverton	cozberg	laporte	redcreek	stecum	virgelle
beenom	dast	larim	rencot	tally	wabek
beisigl	dayschool	lihen	rentsac	tamely	wilsonville
belain	dominic	lisk	rhame	tassel	woodgulch
biglake	dwyer	lolopeak	ridge	terry	woodside
blackhall	elkner	lone rock	riedel	tinsley	worland
blanchard	elmira	lowercreek	riverside	tipperary	yellowbay
bonner	ervide	macfarlane	rivra	tomichi	yetull
brandenburg	flasher	manning	rochester	travessilla	zeona
branham	fleak	mccaffery	ryell	travson	
busby	glaciercreek	mcilwaine	ryorp	treble	
cabba	glenberg	mires	sacheen	trembles	
NEBRASKA					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alda	canyon	duda	hoffland	ord	talmo
almeria	carr	dunday	holt	orphan	tassel
anselmo	cass	dwyer	inavale	ortello	thurman
bankard	chappell	eckley	ipage	orwet	trelona
barney	cheyenne	els	laird	phiferson	treon
bigwinder	craft	elsmere	las animas	pivot	tryon
blendon	dailey	fonner	lute	platte	tuthill
boel	darr	gannett	manter	ronson	valent
bolent	dickinson	glenberg	marlake	sarpy	valentine
boone	dix	gothenburg	mckelvie	schamber	whitelake
brunswick	doger	haverson	meadin	selia	wildhorse
busher	draknab	hedville	munjor	shingle	
calamus		hersh	o'neill	simeon	
NORTH CAROLINA					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alaga	centenary	galestown	mandarin	pocomoke	tomahawk
alpin	charleston	immokalee	molena	potomac	valhalla
baymeade	chipley	johns	nansemond	pottsburg	wake
biltmore	conetoe	kenansville	newhan	remlik	wakulla
blanton	corolla	klej	norfolk	rimini	wando
bojac	dragston	kureb	ona	rumford	wateree
buncombe	duckston	lakeland	osier	seabrook	
butters	echaw	leon	ostin	st. lucie	
cainhoy	eustis	lillington	ousley	suncook	
candor	foreston	louisburg	pactolus	tarboro	
carteret	fripp	lynn haven	plummer	toisnot	

SOIL TYPE RESTRICTION TABLE (continued)

NORTH DAKOTA					
<p>If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
appam	chinook	fossum	lohnes	seroco	verendrye
arvilla	claire	glendive	maddock	sioux	virgelle
aylmer	coe	hamar	manning	stirum	wabek
banks	cohagen	hanly	metigoshe	telfer	wamduska
bantry	cormant	haskill	minnewaukan	tinsley	yecross
beisigl	dast	havrelon	osakis	trembles	yetull
binford	desart	hecla	poppleton	trey	zeona
blackhall	egeland	inkster	renshaw	tusler	
blanchard	ekalaka	karlsruhe	rhame	twilight	
brandenburg	falsen	ladner	ruso	ulen	
breien	flasher	lemert	schaller	vebar	
cabba	fleak	lihen	serden	venlo	
OHIO					
<p>If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
abskota	bronson	elnora	lakin	perrin	spinks
algansee	casco	galen	landes	pinegrove	stafford
alvin	coloma	granby	lanier	plainfield	steinsburg
arkport	colonie	hazleton	maumee	princeton	stonelick
barkcamp	conotton	homer	oakville	rodman	tedrow
bixler	dekalb	hoopeston	oshtemo	schaffenaker	tyner
boyer	dixboro	ionia	otisville	sisson	watertown
brady	dunbridge	kingsville	ottokee	sparta	
SOUTH CAROLINA					
<p>If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. At planting application only: if AgLogic 15G is applied to cotton, peanuts or soybeans as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Side Dress Application Only: If AgLogic 15G is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Split Application: If AgLogic 15G is applied to cotton as an At-Plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>					
alaga	carteret	eustis	louisburg	pactolus	tomahawk
alpin	centenary	foreston	lynn haven	plummer	wakulla
baratari	charleston	foxworth	molena	pocomoke	wando
baymeade	chiple	fripp	mouzon	ridgeland	wateree
blanton	corolla	johns	murad	rimini	witherbee
brogdon	crevasse	kenansville	nansemond	rosedhu	
buncombe	dragston	kershaw	newhan	scranton	
butters	duckston	kureb	olanta	seabrook	
cainhoy	echaw	lakeland	osier	seewee	
candor	elloree	leon	ousley	tarboro	

SOIL TYPE RESTRICTION TABLE (continued)

SOUTH DAKOTA					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
almeria	chinook	flasher	lute	rockoa	tryon
alwilda	cohagen	fleak	maddock	ronson	tusler
anselmo	claire	fossum	manning	sarpy	tuthill
arvilla	craft	gannett	manter	schamber	twilight
assinniboine	dailey	glenberg	marlake	serden	ulen
aylmer	dix	glendive	mawer	seroco	valent
bankard	doger	hamar	mccaffery	shingle	valentine
banks	dogiecreek	hanly	meadin	simeon	vebar
bantry	duda	hecla	minnewaukan	sioux	venlo
beisigl	dunday	henkin	munjor	stirum	wabek
blackhall	dwyer	holt	murdo	storia	whitelake
blendon	eckley	hopdraw	o'neill	talmo	yecross
boel	egeland	inavale	orton	tassel	zeona
butche	ekalaka	ipage	orwet	tefer	
cabba	els	kirby	platte	thurman	
canyon	elsmere	ladner	renshaw	travessilla	
cass	evridge	las animas	reva	trembles	
chappell	fedora	lihen	rhame	trey	
TENNESSEE					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
barkcamp	bruno	crevasse	nugent		
bigbee	buncombe	eustis	potomac		
VIRGINIA					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
alaga	catpoint	fripp	lakin	osier	seabrook
assateague	chipley	galestown	leetonia	ousley	spessard
aura	conetoe	johns	leon	pactolus	tarboro
biltmore	corolla	kenansville	magotha	plummer	wakulla
bojac	dragston	klej	millrock	pocomoke	wando
buncombe	drall	kureb	molena	remlik	wateree
camocca	duckston	lakehurst	nansmond	rumford	
carteret	fisherman	lakeland	newhan	schaffenaker	
WEST VIRGINIA					
If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
drall	lakin	landes	leetonia	potomac	schaffenaker

SOIL TYPE RESTRICTION TABLE (continued)

WYOMING					
If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.					
ashbon	chinook	glenton	meadowlake	rentsac	tipper
adilis	clarkelen	grenoble	means	rivra	tipperary
alcova	comad	grieves	millburne	rogert	tolman
alflack	conpeak	grimm	milok	rohonda	travessilla
anasazi	cordes	grimstone	mishak	roxal	travson
anselmo	corlett	gunbarrel	mosroc	ryark	trelona
apeldorn	cotha	handran	motoqua	ryorp	trembles
assinniboine	cotopaxi	hanks	mudray	sachet	treoff
bankard	cowestglen	hanly	nathale	sawcreek	treon
banks	cowood	hapjack	nelson	schamber	trook
barcus	coyet	haterton	nesda	schooner	troutville
barnum	cragosen	haverdad	newfork	scravo	tulloch
baroid	crestman	haverly	nizina	seedskadee	turnback
barrett	crownest	havermom	norrison	sharland	turnercrest
bearmouth	cryluha	hazton	norte	sheege	twilight
beaverton	curabith	hechtman	oceanet	shingle	valent
bigwin	dailey	henrieville	onason	shoshone	valentine
bigwinder	dast	herbman	orphan	shurley	vible
billycreek	derrick	hiland	otterson	sobson	vonason
blackhall	devore	hiwan	ouray	southace	walcott
bobtail	dix	hoot	pahlow	southfork	walden
bodorumpe	draknab	huguston	peetz	space city	wall
bosler	dunday	irigul	pendergrass	splitro	wardboro
bottle	dunul	julesburg	pepton	spool	wendover
boyle	dwyer	kandaly	pescar	stecum	wetmore
branham	eckley	koonich	phiferson	sudworth	whaley
breece	edlin	labou	pilotpeak	sunup	wiggleton
brownsto	elk mountain	lamarsh	pineguest	taluce	willwood
bruja	ellicott	lambman	poin	tassel	wilsonville
busher	emblem	laney	preatorson	tasselman	wint
butche	enos	laporte	pugsley	teagulf	worf
byrnie	ethelman	lariat	quealman	teewinot	worfka
calicott	eyre	larim	quealy	tellman	worfman
cambarge	farisita	las animas	randman	tenorio	worfstone
canwall	farson	lazear	ratake	terada	worland
carbol	feltner	littlebear	redbank	terro	yetull
cass	fleak	littsan	redcreek	terry	youjay
castner	folavar	lone rock	redfeather	tetonville	zeomont
cathedral	frontier	macfarlane	rekop	theedle	zeona
cestnik	gas creek	manter	relsob	thermopolis	
chappell	glenberg	mathers	rencot	tieside	
cheadle	glendive	mccaffery	reno	tine	