

The Effect of Self-Care Practices on Pregnant Cardiac Women Outcome and Nursing Implications

Amany Hamed Gad Mohamed¹ and Nadia Ahmed Eladham²

¹Obstetrics and Gynecological Nursing Department, Faculty of Nursing, Zagazig University, Zagazig, Egypt

²Community Health Nursing Department, Faculty Of Nursing, Fayoum University, Fayoum, Egypt

Email: amanygad92@yahoo.com and [ndia.adham5@gmail.com](mailto:nadia.adham5@gmail.com)

Abstract: Cardiac diseases are a major cause of morbidity and mortality among pregnant women in the developing world. Most women with cardiac disease can go through pregnancy and delivery safely, so long as they are adequately evaluated, counseled and received high quality care. This study aimed to identify the self-care practices performed by pregnant cardiac women in Zagazig university hospital, to find out the birth outcome behind the self-care practices performed by pregnant cardiac women, and to conduct a mother classes, and provide a handout booklet titled standard self-care practices for educating cardiac pregnant women. **Research design:** An exploratory descriptive design was utilized in this study. **Setting:** The study was conducted in antenatal high risk ward & outpatient clinic in obstetric department, at Zagazig university hospital. **Subjects:** A total of 180 pregnant cardiac women was collected over one year period, starting from the 1st of January 2017 to 31 December 2017. **Tools of data collection:** A structured interview questionnaire form, it entailed three parts: Part I: This includes: Socio-demographic characteristics, present pregnancy data, current pregnancy data regarding practices, current treatment regimen, current pregnancy investigations. Part II: Self-Care Assessment (SCAS). Part III: Assessing newborn and labor outcome. The study **Results:** indicates that (100.0%, 100.0%, 96.7%, 91.7%, 81.7%, 80.0%) respectively of the studied cardiac pregnant women had unsatisfactory score levels of reported practices related to practicing exercise, foot care, nutrition, personal hygiene, teeth care, as well as treatment and follow up of pregnancy. Also, (56.7% & 63.3%) of the studied women had high level of psychological and social stressors and only 3.3% of the women had high total coping level. As regards outcome, 86.7% of the studied cardiac pregnant women had full term pregnancy. Also, the same percentage 86.7% of the women had conducted normal vaginal delivery & (18.3%) of the studied women had delivered babies with unhealthy Apgar score less than 7. **Conclusion:** The study showed that the studied cardiac pregnant women had unsatisfactory score levels of reported daily practices related to practicing exercise, foot care, nutrition, personal hygiene, teeth care, as well as treatment and follow up of pregnancy, and only few percentages of the women had high total coping level toward psychological and social stressors. **The study recommended** that, carrying out educational program about cardiac disease during pregnancy in MCH centers and outpatient clinics to provide the women with necessary knowledge and daily practices. Regular check and follow up visits in the proper settings. The educational booklet should be distributed and implemented in the antenatal clinic, in the hospital and family health centers. Also further researches is needed to investigate the long-term effect of such educational intervention on improvement of the mothers and her babies.

[Amany Hamed Gad Mohamed and Nadia Ahmed Eladham. **The Effect of Self-Care Practices on Pregnant Cardiac Women Outcome and Nursing Implications.** *Biomedicine and Nursing* 2018;4(4): 57-68]. ISSN 2379-8211 (print); ISSN 2379-8203 (online). <http://www.nbmedicine.org>. 9. doi: [10.7537/marsbnj040418.09](https://doi.org/10.7537/marsbnj040418.09).

Key word: Pregnant woman – Cardiac disease – Outcome – Daily practices.

1. Introduction:

The types of the cardiac diseases during pregnancy includes; congenital, acquired, ischemic cardiac disease, cardiomyopathy, maternal placental syndromes, and hypertension in pregnancy (Harris, 2017). Pregnant women who have heart disease have a 100-fold higher mortality than the normal pregnant population, according to new data from the first – ever formal registry on pregnancy and heart disease. Acquired heart diseases are most common in developing countries, these include: rheumatic heart diseases (RHD), cardiomyopathies and ischemic heart disease (Titia et al., 2013).

The prominence of cardiac disease in maternal mortality and morbidity is attributed to several risk

factors; as age, lifestyle trends as work, way of follow up, diagnosis and treatment options, obesity and tobacco use, physical and psychological stressors. Also, the presence of chronic medical condition as diabetes (Bowater and Thorne, 2016).

The cardiac disease may affect pregnancy by polyhydramnios, preterm labor, postpartum hemorrhage, abortion, fetal hypoxia and death, small for gestational age, and low Apgar. While, the women with mild and low risk heart lesion can do well during pregnancy, and are adequately evaluated, and receives high quality care, but some women with high-risk cardiac lesion will not tolerate the hemodynamic changes of pregnancy and become worse (Johns, 2013).

Pregnancy makes a significant demand on the cardiovascular system. Therefore, it follows that woman with cardiovascular compromise due to cardiac disease needs specialist input and careful management, pre-, peri-, and post-partum. Cardiac disease is the most common cause of indirect maternal deaths and the most common cause of death overall (Lewis, 2017).

According to Mekhemar (2015) once pregnant women becomes a patient with cardiac disease, she should be seen early in the first trimester by obstetricians to document gestational age. This plays an important role in decisions regarding timing of delivery. Assessment by a cardiologist early in pregnancy also is indicated. Thereafter, patients are seen for prenatal visits every two weeks more frequently if necessary.

The midwifery and the community health nurses can act together and play a central role in working with the patients to promote the best outcome, they act as counselors, assessors, educators, health care providers, advocates, and researchers. A woman in previous studies identifies numerous inhibitors and facilitators in making changes. Nurses may utilize this provided information to assist in helping women to identify obstacles and inhibitors and to develop strategies to initiate health behaviors changes.

Significance of the study:

The incidence of heart disease with pregnancy worldwide is between 0.2 and 3.7%. Rheumatic heart diseases account for about half of the cases, while congenital heart defects are responsible for most of the second half. Maternal mortality in cardiac pregnant patients is 10% worldwide and 8% in Egypt (Gelson & Johnson, 2016). On the other hand, most of the studies about the cardiac disease during pregnancy consider with the medical aspect as diagnosis, treatment, management and safe delivery, not considered with daily practices and its effect on outcome.

Aim of the study:

- To identify the self-care practices performed by pregnant cardiac women in Zagazig university hospital.
- To find out the birth outcome behind the self-care practices performed by pregnant cardiac women.
- To conduct a mother classes, and provide a handout booklet titled standard self-care practices for educating cardiac pregnant women.

Research questions:

- What are the self-care practices that will be performed by pregnant cardiac women during pregnancy?
- What are the birth outcomes that will be behind the self-care practices performed by pregnant cardiac women?

2. Subjects and methods:

Research design:

An exploratory descriptive design was utilized in this study.

Study setting:

The study was conducted in antenatal high risk ward & outpatient clinic in obstetric department, at Zagazig university hospital. Whereas these settings include the high risk pregnant cardiac women in almost Sharkia governorate, the service provided in this ward includes physical examination, noninvasive test (echo & ultrasound), laboratory investigation, ECG, and follow up.

Study subjects:

A total of 180 cardiac pregnant women were collected over one year period, starting from the 1st of January 2017 to 31 December 2017. Women were eligible for recruitment in the study sample if they met the following inclusion criteria:

- cardiac pregnant woman at second and third trimesters of pregnancy
- free from any other chronic diseases
- diagnosed and registered as cardiac pregnant patient in the pre-mentioned settings

Tool of data collection:

A structured interview questionnaire form was designed for pregnant cardiac women by the researcher based on reviewing related literature and expert opinions, tested for validity and reliability, written in the simple Arabic language, and utilized to collect the necessary data. It entailed three parts:

Part 1: This includes:

- Socio-demographic characteristics such as; age, education, occupation, residence, monthly income.
- Present pregnancy data as; last menstrual period (LMP), expected date of delivery (EDD), weight of pregnant women, type of expected delivery....ect.
- Current pregnancy data regarding practices for receiving services and follow up, and person providing the health education.
- Current treatment regimen as; anti-rheumatic drugs, iron supplements & multivitamins, anti-arrhythmic drugs, anti-hypertensive drugs, anti-coagulant drugs.
- Current pregnancy investigations as; complete blood count (CBC), cardiac enzymes, blood group & RH factor, urine analysis for sugar and albumin, blood sugar, noninvasive test (ECO & ECG), sonar, fetal heart rate (FHR).

Part 11: Self-Care Assessment (SCAS) including:

- Self-care daily living practices performed by pregnant cardiac women. It comprised questions related to women's self-care practices in relation to

personal hygiene, teeth care, foot care, nutrition, rest & sleep, exercises, treatment and follow up. Presented in two answers (yes and no).

- Self-care practices related to social and psychological stressors and their coping patterns.

Part 111:

Assessing newborn and labor (outcome): It was completed immediately after birth and follows the ticket for one week after birth. It included an assessment of gestation weeks at delivery, mode of delivery, newborn condition, newborn problems, weight, and length. It is composed of 10 close ended questions in the form "Yes or No" answers. Answers were coded "1 for Yes and 0 for No". APGAR score also used for assessing the new born condition, it was done at 1st and 5th minutes after delivery, and they were considered healthy if the score is equal or more than "7" and unhealthy if it less than "7" (**Finster & Wood, 2015**).

A scoring system for pregnant women self-reported practices was done as follows, each answer scored as one mark for (Yes) and zero for (No). Total score of self-reported practices had been 40 marks, which represent 100%. Final self-reported practices will be considered satisfactory level if the percent score will be equal to or more than 24 points (60% or more) and unsatisfactory level if less than 24 points (60% or less).

Assessment of social and psychological stressors of cardiac pregnant women this is another part of self-reported practices; the original sheet adopted from (**Fraster, 2003**) and modified by the researcher. It is composed of 15 close-ended questions, and 15 coping strategies itemsto overcome these stressors. Presented in three answers (always, sometimes, and never) and range from "zero to two" as follows; 0 for never, 1 for sometimes, and 2 for always.

The total scoring of coping patterns equal "30" was calculated and interpreted as: A) - women who had a total score from "0-10" were classified as having a low coping level. B)-women who had a total score from "11-20" were classified as having a moderate coping level. And C) -women who had a total score from "21-30" were classified as having a high coping level.

In mother classes; pregnant cardiac women were open, speak comfortably and acceptable and expressed a sincere appreciation for education and handouts that were provided. In addition, the researcher used the computer for show media and printouts.

The educational booklet:

General objective:

At the end of reading this booklet, each cardiac pregnant woman will be able to: know the basic

information concerning cardiac disease during pregnancy.

Specific objectives:

At the end of reading this booklet, each cardiac pregnant woman will be able to:

- Acquire knowledge concerning cardiac disease to improve health and outcome.
- Acquire healthy practices to improve health and outcome.

Content validity and reliability:

The questionnaire was reviewed by a panel of five experts in the field of obstetrics and gynecological nursing, community health nursing, and cardiologist from faculty of medicine to test its content validity. Modifications were done accordingly based on their judgment. Cronbach's alpha coefficient was calculated to assess the reliability of the developed tool through their internal consistency.

Field study:

Once permission was granted to proceed with the study, the researcher contacted each woman individually in previous mentioned setting and explained to her the purpose and nature of the study.

Upon obtaining her written consent, the researcher started the interview. The interview time ranged from 20 to 25 minutes in the first meeting to fill part 1 & part 2 in the tool, and the another meeting to collect data about the outcome immediately after birth and follow the ticket for one week after birth. Collection of data covered a period of 12 consecutive month from the 1st of January to end of December 2017, three days per week.

Pilot study:

A pilot study was conducted on a sample of 10% of cases who were not included in the total sample size. It was done to test the study tool in terms of clarity and feasibility, and the time required to be applied. Following the pilot study the questionnaire was reconstructed and necessary modifications were done to reach the final form.

Administrative & ethical consideration:

All ethical issues were taken into consideration during all phases of the study; the researcher maintained an anonymity and confidentiality of the subjects. She introduced herself to the woman and briefly explained the nature and aim of the study to every woman before participation, and women were enrolled voluntarily after written consent. Women were also assured that all information obtained during the study was confidential and used for the research purpose only and they have the chance to withdraw from the study when they want.

Statistical analysis:

Data entry and statistical analysis were done using SPSS statistical software package. Data were presented using descriptive statistics in the form of

frequencies and percentages for qualitative variables, and means and standard deviations for quantitative variables.

As for the limitation of the study, the sample size was too small together with the inability to perform statistical equation to select the study sample, because the low rate of women attendance according to the previously mentioned inclusion criteria. This affected the generalization of the study results;

however, it gave an overview about the effect of self-care practices on pregnant cardiac women outcome. Provided that, this number of 180 pregnant cardiac woman in this study, is know consider larger than the previous time, because the opening of the new ward of cardiac disease in Zagazig University Hospital.

3. Results:

Table (1): Distribution of The Studied Cardiac Pregnant Women According to Their Socio-Demographic Characteristics (n=180)

General characteristics	No=180	%
Age (years):		
18 < 25	33	18.3
25 -31	93	51.7
≥ 32	54	30.0
Educational level:		
illiterate	6	3.3
primary	30	16.7
secondary	105	58.3
university	39	21.7
Occupation:		
housewife	99	55.0
employee	81	45.0
Residence:		
rural	78	43.3
urban	102	56.7
Monthly income:		
sufficient & save	18	10.0
just enough	39	21.7
insufficient	123	68.3

Table (2): Distribution of the Studied Cardiac Pregnant Women According to Their Practices Regarding Laboratory Investigations (n=180)

Lab. investigation	No (180)	%
Complete blood count (CBC)		
- Done	180	100.0
Cardiac enzyme		
- Done	150	83.3
- Not done	30	16.7
Blood group & Rh Factor		
- Done	180	100.0
Urine analysis (sugar & albumin)		
- Done		70.0
- Not done	12654	30.0
Blood sugar		
- Done	69	38.3
- Not done	111	61.7
Non invasive test (Echo, ECG)		
- Done	180	100.0
Ultrasonography & FHR		
- Done	180	100.0

Table (1): Shows the socio-demographic characteristics of the studied cardiac pregnant women. It shows that 51.7% of the studied women were within the age group ranged from 25-31 year. Also 58.3% of the studied women had secondary education and 55.0% of them were housewives. And 56.7% of them were living in urban areas. Concerning family monthly income, 68.3% of the studied women had insufficient income for their basic family demand.

Table (2): Shows distribution of the cardiacpregnant women according to their practices regarding laboratory investigations. This table indicates that all studied women did the complete blood count, blood group, and Rh factor as well as cardiac radiology, ultrasonography and fetal heart rate. Meanwhile, only 16.7% of them did not do cardiac enzyme.

Table (3): Distribution of The Studied Cardiac Pregnant Women According to Their Practices Regarding Treatment Regimen Used (n=180)

Treatment used	No=180	%
Anti-rheumatic drugs	114	63.3
Iron supplements / multivitamins	120	66.7
Anti-arrhythmic drugs	66	36.7
Anti-hypertensive drugs	63	35.0
Anti-coagulant drugs	9	5.0

Table (4): Distribution of the Studied Cardiac Pregnant Women According to Their Daily Self-Reported Practices (n=180)

Reported practice items	No=180	%
Personnel hygiene		
- Satisfactory	15	8.3
- Unsatisfactory	165	91.7
Teeth care		
- Satisfactory	33	18.3
- Unsatisfactory	147	81.7
Foot care		
- Satisfactory	0	0.0
- Unsatisfactory	180	100.0
Nutrition		
- Satisfactory	6	3.3
- Unsatisfactory	174	96.7
Practicing exercise		
- Satisfactory	0	0.0
- Unsatisfactory	180	100.0
Rest & Sleep		
- Satisfactory	57	31.7
- Unsatisfactory	123	68.3
Treatment & Follow up		
- Satisfactory	36	20.0
- Unsatisfactory	144	80.0
Total reported self-practices score level		
- Satisfactory	0	0.0
- Unsatisfactory	180	100.0

Table (3): Indicates 63.3% of the studied cardiac pregnant women used anti-rheumatic drugs. Also, 66.7% of the women used iron supplements and multivitamins. Meanwhile, 36.7% & 35.0% respectively of the studied women used anti-arrhythmic & antihypertensive drugs.

Table (4): indicates (100.0%, 100.0%, 96.7%, 91.7%, 81.7%, 80.0%) respectively of the studied cardiac pregnant women had unsatisfactory score levels of reported practices related to practicing

exercise, foot care, nutrition, personnel hygiene, teeth care, as well as treatment and follow up of pregnancy.

Table (5): indicates that, 86.7% of the studied cardiac pregnant women had full term pregnancy. Also, the same percentage 86.7% of the women had conducted normal vaginal delivery, followed by 11.6% had conducted cesarean section. On the other hand, 90.0% of the women had normal live baby. Only (6.7% & 3.3%) of them respectively had congenital anomalies and stillbirth.

Regarding the newborn problems, the table shows that, the highest percentages of the problems were tachycardia and blue skin color, followed by poor feeding and sucking, then weak muscle tone, and finally poor crying (27.6%,27.6%, 24.1%, 15.5% & 5.2%) respectively. According to Apgar score (18.3%) of the studied women had delivered babies with unhealthy Apgar score less than 7.

Table (6): clarifies that, 84.5% of the newborn birth weight ranged from (2500-3500 gm) that lied within normal range of the weight. Also, 72.4% of the women had babies length ranged between (48-53 cm).

Figure (1): illustrates that, about 76.7% of the studied women received antenatal care. On the other

hand, 71.7% of them maintain regular follow up. Also, this figure displays that 56.7% of the studied cardiac pregnant women reported that doctors are suitable persons for providing health education. Meanwhile, 38.3% of them reported that nurses provided health education.

Figure (2): demonstrates that, (56.7% & 63.3%) of the studied women had high level of psychological and social stressors. Concerning the total coping level of psychological and social stressors, this figure elaborates that only 3.3% of the women had high total coping level in this study.

Table (5): Distribution of The Studied Cardiac Pregnant Women According to Newborn & Labor Outcome (n=180).

Outcome	No=180	%
Gestational weeks at delivery		
Full term	156	86.7
Pre term	24	13.3
Mode of delivery		
Normal vaginal	156	86.7
Caesarean section	21	11.6
Others (Vento's & forceps)	3	1.7
Newborn condition		
Normal live	162	90.0
Congenital anomalies	12	6.7
Still birth	6	3.3
Apgar score		
- 3-6	33	18.3
- 7-10	147	81.7
Problems of newborn (no=174)		
Tachycardia	48	27.6
Blue skin color	48	27.6
Poor feeding & sucking	42	24.1
Weak muscles tone	27	15.5
Poor crying	9	5.2

Table (6): Distribution of The Studied Cardiac Pregnant Women Regarding Weight and Length of Their Newborn (n=174).

Weight/Length	No=(174)	%
Less than 2500 gm	27	15.5
2500-3500 gm	147	84.5
Less than 48 cm	42	24.1
48-53 cm	126	72.4
More than 53 cm	6	3.5




Figure (1): Distribution of The Studied Cardiac Pregnant Women According to Their Practices Regarding Receiving Services and Follow up (no=180)




Figure (2): Distribution of the Studied Cardiac Pregnant Women According to Their Psychological and Social Stressors and Their Coping Pattern (no=180)

4. Discussion:

Self-care practices in pregnant cardiac women are a key concept in health promotion that involves the activities affecting the health of mothers, fetus and newborns. Health practices are important for the result of pregnancy and should be determined and gained during prenatal care (Miyake et al., 2014).

This study was intended to identify self-care practices utilized by pregnant cardiac women in Sharkie Governorate, and its associated pregnancy outcome, also conducting a mother classes and provide a handout booklets for vulnerable women. The study was conducted in antenatal high risk ward and outpatient clinic, in obstetric department, at Zagazig University Hospital.

Women socio-demographic characteristics play a great role in practicing healthy behavior and they affect their health either positively or negatively Aziz and Maqsood (2016). The present study finding revealed that, the highest percentage of the pregnant cardiac women situated in the age group (25 - 31) year, which was (51.7%) (table 1). This may be due to most women with heart problems were married in the middle age of years and have late pregnancy as well as fear from any complications that occurred in young age. This finding is in agreement with a study conducted in Brazil, about quality of life in pregnant women with heart diseases by Meneguín and Xavier (2013) who reported that the mean age was 28.6 ± 5.63 years.

In relation to educational level, income and residence the present study findings revealed that more than half of the studied women had secondary education, had insufficient income for their basic family demands and they were from urban areas. This finding is in agreement with a study about the stressors and coping strategies among cardiac pregnant women, Faculty of Nursing, Ain-Shams University, Egypt by Farag (2005) who, reported that more than half of the studied women had intermediate education and more

than half of them were living in urban areas and they had insufficient income.

The present study results showed that more than half of the studied women were housewives; it may be due to the health condition of cardiac pregnant women not allow doing over or hard activities. This finding is in agreement with Farag (2005), who found that most of the cardiac pregnant women were housewives. While, this finding comes incongruence with Meneguín and Xavier (2013), who showed that the majority of participants had a formal job.

As regard medications used (table 3), the present study findings showed that more than two thirds of the studied women used (long acting penicillin) as anti-rheumatic drugs, and more than one third of them used anti- arrhythmic, anti-hypertensive drugs. On the other hand, more than two thirds used iron supplements and multivitamins. This could be explained that cardiac pregnant women compliance with drug regimens during pregnancy to improve maternal and fetal circulation under medical supervision. This result was in disagreement with the study conducted by Makino et al. (2012) in Japan about risk factors associated with preterm delivery in women with cardiac disease, who reported that the cardiac medication used by pregnant cardiac women constituted 3.2% diuretic, 2.2% digoxin, 1.2% antihypertensive, and 1.0% warfarin.

In relation to laboratory investigations (table 2), the present study findings indicated that all studied women did complete blood count, blood group and Rh factor as well as noninvasive test (cardiac radiology, ultra- sonography, and fetal heart rate). This may reflect to availability of the antenatal care in this setting of the study, and increase awareness of studying women about the importance of follow up for antenatal care. This finding was in accordance with Mohy-Eldine (2014) who mentioned that, most pregnant women made laboratory investigations as blood group, hemoglobin, Rh factor, urine and blood sugar.

As regards follow up, the present study findings revealed that the majority of the studied women had regular follow up (figure 1). Because the cardiac disease is a danger to pregnancy and most of cardiac pregnant women maintain regular visits and take care. However, close follow up of women with cardiac diseases will improve maternal tolerance of the cardiovascular burden imposed by pregnancy, promote fetal growth and neonatal survival, and enable adjustments to cardiovascular changes in the post-partum period. This finding was in agreement with the study conducted by Zahra (2012) in Qalubiya governorate, about quality improvement of maternal health care in maternal and child health center, who emphasized that most of the studied cardiac pregnant

women regularly, attended the antenatal clinic for follow up.

As for the suitable person for providing health education about cardiac disease the present study findings revealed that, more than one half of the sample reported doctors, and more than one third of them reported nurses. This finding was in disagreement with **Ranimah et al. (2012)**, who reported that slightly more than one quarter of cardiac women obtained the cardiovascular disease information from health care workers.

Regarding the reported total practices score levels of the studied women about cardiac diseases, the present study results revealed that, 100.0% of the studied sample had unsatisfactory total reported practices score level (table 4). This finding was in agreement with **Rosediani et al. (2012)**, who clarified that the self-reported practice on cardiovascular disease on healthy lifestyle was very poor among women in the North-East coast of Malaysia.

As regards psychological and social stressors of cardiac pregnant women (figure 2), the present study findings revealed that, more than half of the studied women had high levels of psychological and social stressors. This could be explained that the studied women expresses about the internal feeling during the educational sessions and feel assured about the psychological, social condition as well as the pregnancy outcome in the future.

Concerning the total coping level of psychological and social stressors, finding showed that most of the studied women had low total coping level. This may be due to poor knowledge and reported practices among the studied women. This finding was in agreement with **Farag (2005)**, who asserted that the majority of the studied women could not highly cope with stressors of cardiac disease during pregnancy.

According to the newborn outcome, the present study findings revealed that, the majority of the studied women had a full term and normal live babies (table 5). This finding is in agreement with the study conducted in India about clinical profile and obstetric outcome in pregnancies complicated by heart disease by **Bagde et al. (2013)** who reported that, the majority of gestational ages were full term at time of delivery, were between 37-42 weeks. Also, **Diao et al. (2011)** who found that most of fetal outcome for pregnant women with heart disease were normal live births, and few of them had preterm. This agreement of the two previous studies may reflect the advanced worldwide in management of the pregnant women with cardiac disease includes the medication effect and new technology for the treatment of cardiac disease in pregnant women.

As regards mode of delivery, the present study results revealed that the majority of the women had

delivered by normal vaginal mode. This may be due to most pregnant women with cardiac disease preferred delivery by vaginal mode with episiotomy to shorten the second stage of labor compared to caesarean section as well as the previous delivery of them was through the same mode. This finding is in the same stream with **Bagde et al. (2013)** who found that more than two thirds of the women had spontaneous vaginal delivery and less than one third of them had delivered by caesarean section.

In relation to Apgar score the present study findings revealed that, the majority of the studied women born babies with good Apgar score between (7-10). This finding is in the same line with a study conducted on pregnant cardiac women at Faculty of Nursing Zagazig University, Egypt, by **Elsabagh and Zaiton (2013)** who stated that the mean of Apgar score at 5 minutes was 8.04 ± 1.65 . This may be explained that the improved fetal outcome due to the presence of specialists during labor as obstetrician, cardiologist, anesthetists, hematologist, nurses, and neonatologist.

Concerning the newborn birth weight the present study findings showed that, the majority of birth weight were within the normal weight of the newborn (table 6). This finding is in agreement with **Taha et al. (2013)** who found that the mean birth weight was 2513.7 ± 922.2 Kg. On the other hand, **Elsabagh and Zaiton (2013)** who reported that the mean of birth weight was 2897 ± 838 Kg among patients with cardiac disease in pregnancy.

Conclusion:

The study showed that the studied cardiac pregnant women had unsatisfactory score levels of reported daily practices related to practicing exercise, foot care, nutrition, personnel hygiene, teeth care, as well as treatment and follow up of pregnancy, and only few percentages of the women had high total coping level toward psychological and social stressors.

Recommendations:

- Carrying out educational program about cardiac disease during pregnancy in MCH centers and outpatient clinics to provide the women with necessary knowledge and daily practices.
- Regular check and follow up visits in the proper settings during pregnancy.
- The educational booklet should be distributed and implemented in the antenatal high risk ward & outpatient clinic, in the hospital and family health centers.
- An also further research is needed to investigate the long-term effect of such educational intervention on improvement of the mothers and her babies.

References:

1. Aziz KF and Maqsood SHS (2016): Self-Management of Pregnant Women Regarding Minor Discomforts in Primary Health Care Centers in Erbil City,13(2),284-293.
2. Bagde, D., Bagde, N., Shivkumar, P and Tayade, S., (2013): Clinical Profile and Obstetric Outcome in Pregnancies Complicated by Heart Disease: A five Year Indian Rural Experience. Research Article International Journal of Reproduction, Contraception, Obstetrics and Gynecology Int J Report Contracept Obstet Gynecol. Mar; 2 (1) Pp: 52-57.
3. Barlow J (2012): Self-care: International Encyclopedia of rehabilitation, (2012), Coventry University, UK, A valaible: j.barlow@coventry.ac.uk, accessed at 20/4/2016.
4. Bowater, S. E and Thorne, S. A. (2016): Management of pregnancy in women with acquired and congenital heart disease post graduate medical Journal, 86 (1012), pp: 100-105.
5. Crawley H (2013): Eating well in pregnancy a practical guide to support teenagers. London: First Steps Nutrition Trust.
6. Cunningham F.G Leveno K.G Bloom S.L., Hauth J.C., Rouse D.J., and Spong C.Y., (2010): Cardiovascular Disease. United States of America Williams Obstetrics. 23rd ed.: McGraw-Hill Companies.
7. Diao, M., Kane, A., Ndiaye, M., Mbaye, A., Bodian, M., Dia, M., Sarr, M., Kane, A., Monsuez, J and Abdou, S., (2011): Pregnancy in Women with Heart Disease in Sub-Saharan Africa, CLINICAL RESEARCH, Archives of Cardiovascular Disease,© 2011 Elsevier Masson SAS.104, Pp:370-374.
8. Elsabagh, EM., and Zaiton, HI., (2013): Impact of Intervention Program on Nurse's Knowledge and Practice regarding Nursing Care of Pregnant Cardiac Patients during Labor, Faculty of Nursing, Zagazig University, Egypt, Med. J. Cairo Univ., Vol. 81, No. 2, (12), Pp: 141-153.
9. Farag A. F., (2005): Stressors and Coping Strategies among Cardiac Pregnant Women. Unpublished Master Thesis in Maternity and Newborn Nursing, Faculty of Nursing, Ain shams University, Egypt, Pp: 64 – 65.
10. Finster M., And Wood M., (2015): The Apgar score has survived the test of time. Anesthesiology 102 (4): p-p. 855-857.
11. Fraster L., (2003): Principle and Practices of Psychiatric Nursing, 7th ed., Mosby, U.S.A., PP: 60-78.
12. Gelson, E., And Johnson, M., (2016): Effect of Maternal Heart Disease on Pregnancy Outcomes. Vol.5, No. (5) PP: 617- 650.
13. Hanem, A., (2015): Effect of educational program on pregnant women with cardiac disease to improve health and outcome, Unpublished thesis, PHD Thesis, Fayoom University, Faculty of nursing.
14. Harris, I., (2017): Management of pregnancy in patient with congenital heart disease. Progress in cardiovascular disease, 53 (4), P.p 305-311.
15. Johns, J., (2013): Cardiac disease. O & G Magazine; 15(1) pp: 30-2.
16. Kurijak, A., and Chervenak, F. (2012): Donald School Manual of Pracial Problems in Obstetrics, 1st ed., Medical diseases and pregnancy, cardiac disease during pregnancy, Japee Brothers Medical publisher, New Delhi, Panama city, London, Pp 3-4.
17. Leonard, D., and Lerry, S., (2007): Maternity and Women's Health Care, Medical-Surgical Problems in Pregnancy, Cardiovascular Disorders, 9ed., Mosby, Elsevier, San Francisco, California. Pp. 849-60.
18. Lewis G., (2017): The Confidential Enquiry into Maternal and Child Health. Saving Mothers' Lives: Reviewing Maternal Deaths to Make Motherhood Safer 2003-2005. The seventh Report on Confidential Enquires into Maternal Deaths in the United Kingdom, London.
19. Makino, Y., Matsuda, Y., Mitani, M., Shinohara, Tand Matsui H., (2012): Risk Factors Associated with Preterm Delivery in Women with Cardiac Disease. Japanese College of Cardiology. Published by Elsevier Ltd. journal of Cardiology.59. 27 March, P:p: 291-298.
20. Meneguín, S and Xavier, C., (2013): Quality of Life of Pregnant Women with Heart Disease.Text Context Nursing, Florianopolis, Jul-Set; 22(3) Pp: 811-8.
21. Miyake Y, Tanakan K, Okubo H, Sasaki S and Arakawa M (2014): Alcohol consumption during pregnancy and birth outcomes: the Kyushu Okinawa maternal and child health study. BMC Pregnancy and Childbirth; 14:79. Doi: 10.1186/1471-2393-14-79.
22. Mohy-Eldine, HF. (2014): The impact of health educational intervention for rural health nurses on their quality of home visit for women during maternity cycle, Unpublished thesis, PHD in Community Health Nursing Department, Benha University, Faculty of nursing.
23. Murray, S., and Mc Kinney, S., (2006): Concurrent disorders during pregnancy families at risk during childbearing period. Foundation of

- Maternal – Newborn Nursing, 4th ed., Saunders, Elsevier, California. Pp 672:76.
24. Ranimah, Y., Rosediani, M and Harmy, MY., (2012): Association between Knowledge, Attitude and Practice on Cardiovascular Disease among Women in Kelantan, Malaysia, International Journal of Collaborative Research on Internal Medicine and Public Health; No. 8(1) Pp: 1507-23.
25. Regitz, Z. (2011): ESC Guidelines on the management of cardiovascular diseases during pregnancy. European Heart Journal European Society Cardiology, (32) Pp; 3147-3197.
26. Rosediani, M., Ranimah, Y and Harmy MY., (2012): Knowledge, Attitude and Practice on Cardiovascular Disease among Women in North-East coast Malaysia, International Journal of Collaborative Research on Internal Medicine & Public Health; No. 4(1), Pp:85-98.
27. Simpson, S., (2013): Cardiac Disease in Pregnancy and Promoting the Health of Women and Newborn, Perinatal Nursing, 4th ed., Lippincott Williams & Wilkins, Philadelphia, London, Pp 224:243.
28. Taha, N., Mahmoud, K., Eisa, M and Darder, A., (2013): Structural Heart Disease in Pregnancy in El-Minia Localities. The Egyptian Heart Journal Volume 65, Issue 2, June, Pp. 99–109.
29. Titia, p., Ruys, M., Jerome, M., Jolien, W and Roos, H., (2013): Review pregnancy and delivery in cardiac disease. The Netherlands Journal of Cardiology Volume 61, Issue 2, February. Pp: 107-112. Received 15 August 2012. Accepted 10 November 2012. Available online 2 January. Lincense: Http://Www. Elsevier.Con / Open-Access/User license.
30. Zahra, A.N. (2012): Quality Improvement of Maternal Health Care in Maternal and Child Center Qalyubia Governorate, Unpublished thesis, PHD Thesis, Benha University, Faculty of nursing.

Electronic website:

- <http://www.Americanheart.Org/Presenter.Jhtml?Identifier=4688>. Accessed March 31, 2011.Pregnancy and Heart Disease.American Heart Association.
- <http://www.heartdiseaseandpregnancy.Com>. Copy right 2011. Heart Disease and Pregnancy.
- <http://www.athersstream.com/2012.Presentation/OsmanAhmed90-488126-HeartDiseaseDuringPregnancy>.
- <http://www.google.com.eg/search?q=picture+of+pregnant+women+with+cardiac+disease>

Appendix

نموذج إرشادي حول ممارسات العناية الذاتية أثناء فترة الحمل للسيدات الحوامل المصابات بمرض القلب

القلب مرض عن عامة معلومات أولا

- ✓ الشرايين تصلب: هي القلب بأمراض المقصود الجلطة، القلب عضلة ضعف، الدم ضغط ارتفاع القلبية.
- ✓ الشرايين أمراض: في تتمثل القلب أمراض القلب أمراض، القلب عضلة قصور، التاجية ارتفاع عن الناتج القلب مرض، والدموية والأوعية مرض، الحمل أثناء الدم ضغط ارتفاع، الدم ضغط القلب صمامات مرض وأيضا، الإلتهابي القلب.
- ✓ التاريخ: هي الخطر وعوامل القلب أمراض أسباب أن بمرض الإصاية، الدم ضغط ارتفاع، العائلي، الحركي النشاط قلة، الوزن زيادة، السكر الإكتئاب وأيضا، السليبي والتدخين التدخين.
- ✓ باختلاف تختلف: القلب أمراض وعلامات أعراض أو اليمني اليدفي ألم تكون وقد القلب مرض نوع، الرقبة اوردة انتفاخ، الصدر في ألم، الظهر بالغثيان الشعور، إغماء، (الجلد زرقة) إزرقاق أو الإستلقاء عند التنفيس صعوبة، والدوار النبض في خلل وأيضا، النوم من الصحو.

- ✓ الحامل السيدة لها تتعرض التي المضاعفات الإحتقاني القلب فشل هي:بالقلب المصابة توقف أو القلبية النوبة، القلب في حاد وهبوط القلبية والسكتة الدماغية السكتة، فجأة القلب بالقلب المحيطة الشرايين أمراض، المفاجئة الشريان إنسداد، الرئوي الشريان إنسداد، صدمة، الشرياني الدم ضغط ارتفاع، الميترالي ميكرة ولادة أو إجهاد.
- ✓ الجنين لها يتعرض أن ممكن التي المضاعفات: الميعاد بعد طفل ولادة، الميعاد قبل طفل ولادة نمو تخلف، خلقية عيوب، الوزن ناقص طفل أو، الرحم الجنين داخل موت أو، الرحم داخل الجنين في مشاكل أيضا، المشيمة انفصال، الولادة بعد التنفسي الجهاز.
- ✓ أثناء القلب أمراض لتشخيص المهمة الفحوصات ضغط قياس، للقلب صورة، القلب رسم: الحمل، الدموية والأوعية للقلب البدني الفحص، الدم الموجات، الدم فحوصات، بالمجهود القلب رسم أشعة، القلب قسطرة، القلب علي الصوتية

- أخذ، المغناطيسي بالرنين التصوير، مقطعية الضرورة عند القلب من عينة
- ✓ الجسدية الراحة: **في تتلخص أمراض وعلاج**، الرياضية التمرينات، الغذائي النظام، والعقلية لضربات منظم جهاز تركيب، بالأدوية العلاج الحمل أثناء للقلب عملية، (اسميكرالب) القلب
 - ✓ وقف: **القلب أمراض من السبعة بالمنجيات واليكسي** ساعة نصف، الوزن علي المحافظة، فوراً التدخين 75 أطفال أسبوعين قرص، أسبوعياً مرات 5 مشي 2، يومياً ه فيتأمين كبسولة، يومياً ملجرام تناول، يومياً الرقيق علي كبسولة أو ثوم فص يومياً تقاحة
 - ✓ **لخفض البديل الطب في العلاجات بعض هناك**، البيضاوي الحمل لسان نبات مثل: **الكولستيرول** الأحماض، الشوفان ونخالة الشوفان، الكتان بذور لأخضرا الشاي، الدهنية
 - ✓ عليك **ينبغي: القلب في مرض أي لديك كان إذا** قبل الصحية حالتك وتقييم الطبيب إستشارة له والتخطيط القادم الحمل
 - ✓ **الشائعة القلب مرض أعراض علي التغلب كيفية** **:فوراً عنها والإبلاغ الحمل أثناء**
 - ووضع نصفي وضع في الجلوس: التنفيس ضيق الظهر خلف وسادة
 - والاتصال فوراً العلاج أخذ: الصدر في ألم بالطبيب
 - الرأس رفع وعدم الضغط قياس: الأنف نزيغ إلي والوصول الدم يتوقف حتي لأعلي المستشفى
 - وإبلاغ مريح وضع في الجلوس: والإغماء الدوار الإسعاف
 - الطبيب وإستشارة وصوفامال العلاج أخذ: الصداع أوية، ساخنة سوائل شرب: والخناق يليال السعال، جلسة نصف وضع في كوني، للسعال مهدئة بالطبيب الإتصال
 - تنظيم علاج أخذ: **القلبو الخفقان ضربات زيادة** الطبيب وإستشارة القلب ضربات
 - ✓ **بعد تحدث التي المنذرة والأعراض المخاطر أهم** ضيق: **القلب بمرض المصابة للام الولاد** بعد الغزير النزيف، بالصدر ألم، التنفيس الحرارة درجة إرتفاع، الساقين جلطة، الولادة الدم ضغط إرتفاع، ورعشة
 - ✓ **ملاحظة، تدفئة: كالاتي الجديد بمولودك الإهتمام** الطبقة ملاحظة، القلب وضربات التنفيس العناية، عليها والحفاظ الجلد علي البيضاء حمام، المنعكسة الأفعال ردود ملاحظة، بالسرة بدء، الطفل لنوم المناسب الجو توفير، الطفل يمكن إن الطبيعي الرضاعة

المصابة الحامل للام اليومية الصحية الممارسات ثانيا **:الحمل فترة أثناء القلب بمرض**

والعلاج المتابعة:

- العلاج أخذ بمواعيد متاز الإل
- لكي توصف التي فيتامينات خذاً
- إسبوعياً الضغط قياس متابعة
- الحمل ومتابعة الزيارة مواعيد علي الحرص
- القلب رسم وعمل المطلوبة التحاليل بعمل القيام

مشكلة بأي الشعور عند الطبيب الي الرجوع صحية

:التغذية:

- العناصر جميع علي يحتوي صحي طعام تتاولي الغذائية
- وأوقاتهما في يةأئ الغذ الوجبات تناول في الإنتظام ومتعددة خفيفة وجبات تناول مع
- ، الصحية الحالة حسب الموصوف الطعام تتاولي والحرق المسبك وعدم المسلوقة ويفضل هناك كان إذا خصوصاً الطعام في الملح من قللي الدم ضغط في إرتفاع
- لأنها التونة مثل البيضاء السمك لحوم من الإكثار للقلب مفيدة
- من الإكثار، والفاكهة الخضروات تناول من الإكثار الخضراء السلطة
- والبقوليات الحبوب مثل الكربوهيدرات تتاولي وجبات علي وتوزع الأسمر والخبز
- الدسم عالي والحليب والحلويات السكريات تقليل و المشبعة الحيوانية الدهون عن الإمتناع والذرة الزيتون زيت مثل نباتية بدهون التعويض الشمس ودوار
- وجبات 3، إسبوعياً حمراء لحوم واحدة وجبة تناول 3، الجلد منزوعة ومشوية مسلوقة وطيور دجاج سمك وخصوصاً إسبوعياً الأسماك من وجبات البيض وتناول، والسردين والتونة السالمون صفار بدون
- الإمكان بقدر المقلي الطعام تجنب والإستاكوزا الجمبري من المتأولة الكميات قليل الكوليسترول عالية والكابورياو
- مرتين من أكثر الطعام قللي في الزيت إستخدام عدم ودوار والذرة الزيتون زيت استخدام ويفضل القللي في الشمس
- يومياً المكسرات من صغيرة حفنة تناول والحديد وحمض بالكالسيوم الغنية الأظعمة تناول الفولييك
- والعصائر والسوائل الماء من كبيرة كمية تناول بجوي، الغازية السوائل شرب تجنبي، الطبيعية أو الكلي أو بالقلب هبوط وجود عند الكمية تحديد آخري موانع أي
- الجهاز حماية علي يساعد والمشروم الثوم تناول كذلك والمخ العام المزاج تحسين، والقلب الدوري المفاصل إتهابات من يقلل

:والملايس الشخصية النظافة:

- صيفا الإسبوع أيام طوال لإستحمام
- شتاء الإسبوع أيام بعض جزني حمام عمل
- ومصنوعة للحمل مناسبة فضفاضة ملايس إرتداء القطن من
- استخدام عند يومياً العانة منطقة بنظافة القيام الحمام
- أي وجود عند والمطهرات الدافئ الماء إستخدام طبيعية غير إفرزات أو إتهابات

:بالأسنان العناية:

- ✓ بحركة يومياً السواك أو بالفرشاة الأسنان غسل لأعل أسفل ومن لأسفل أعلى من دائرية
- ✓ مشاكل أي وعلاج الأسنان لفحص للطبيب الذهاب بالفم

بالقدمين العناية:

- ✓ مستمرة بصفة القدم فحص
- ✓ والصابون الدافئ بالماء القدم سلغ
- ✓ للقدم المرطب الكريم استخدام
- ✓ باستمرار القدم أظافر تقليم
- ✓ الصحية وحالتك للحمل ومريح مناسب حذاء ارتدي
- ✓ في مشاكل وأي التهابات أي لعلاج للطبيب الذهاب القدم

الرياضة:

- ✓ يوميا ساعة نصف لمدة المشي ممارسة
- ✓ للرجلين مساج
- ✓ العنيفة الرياضة ممارسة تجنب

والنوم الراحة:

- ✓ 6 و ،الظهيرة وقت 2 ،يوميا ساعات 8 الأقل علي الإجهاد وعدم الراحة مع ،الليل أثناء ساعات

النفسية الضغوط مع والتكيف التغلب كيفية ثالثا والإجتماعية

- ✓ الراحة من كافية أوقات خذي
- ✓ والأقارب الأهل من الآخرين مع تحدثي

- ✓ أخري بمهام نفسك إشغلي
- ✓ بالمرض نفسك تشغلي ولا الله الي تقربي
- ✓ والإستفادة للآخرين المفيدة بالنصائح العمل مشاكلك حل في منهم
- ✓ البسيطة الرياضية التمرينات بعض عمل
- ✓ والحمل القلب مرض عن تفيدك معلومات عن البحث
- ✓ القلق علي للتغلب حولك من الأشياء تغير
- ✓ التلفزيون شاهدي
- ✓ الحمل ومتابعة العلاج أخذ في انتظمي
- ✓ بنفسك مشاكلك تحلي أن حاولي
- ✓ للبكاء اللجوء
- ✓ التنفيس تمارين مارسي
- ✓ أخذ الأغاني سماع مثل الإسترخاء وسائل إستخدمي شاور
- ✓ أسرتك مع تحدثي
- ✓ التسوق متعة استخدمي
- ✓ الجديد الطفل إستقبال في نفسك اشغلي واحتياجاته لبسة وتجهيز

12/25/2018