

**Solomon Press**  
**Core Mathematics C3**  
**Paper J**  
**(Question Paper)**

**All exam papers are issued free to students for education purpose only.  
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)  
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations  
Advanced Subsidiary

## Core Mathematics C3

Paper J

Time: 1 hour 30 minutes

### *Instructions and Information*

---

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration.

Full marks may be obtained for answers to ALL questions.

Mathematical formulae and statistical tables are available.

This paper has seven questions.

### *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner.  
Answers without working may gain no credit.



*Written by Shaun Armstrong*

© Solomon Press

*These sheets may be copied for use solely by the purchaser's institute.*

1. (a) Given that  $\cos x = \sqrt{3} - 1$ , find the value of  $\cos 2x$  in the form  $a + b\sqrt{3}$ , where  $a$  and  $b$  are integers. (3)

(b) Given that

$$2 \cos (y + 30)^\circ = \sqrt{3} \sin (y - 30)^\circ,$$

find the value of  $\tan y$  in the form  $k\sqrt{3}$  where  $k$  is a rational constant. (5)

---

2. The functions  $f$  and  $g$  are defined by

$$f(x) \equiv x^2 - 3x + 7, \quad x \in \mathbb{R},$$

$$g(x) \equiv 2x - 1, \quad x \in \mathbb{R}.$$

(a) Find the range of  $f$ . (3)

(b) Evaluate  $gf(-1)$ . (2)

(c) Solve the equation

$$fg(x) = 17. \quad (4)$$

---

3. 
$$f(x) = \frac{x^4 + x^3 - 13x^2 + 26x - 17}{x^2 - 3x + 3}, \quad x \in \mathbb{R}.$$

(a) Find the values of the constants  $A$ ,  $B$ ,  $C$  and  $D$  such that

$$f(x) = x^2 + Ax + B + \frac{Cx + D}{x^2 - 3x + 3}. \quad (4)$$

The point  $P$  on the curve  $y = f(x)$  has  $x$ -coordinate 1.

(b) Show that the normal to the curve  $y = f(x)$  at  $P$  has the equation

$$x + 5y + 9 = 0. \quad (6)$$

---

4. (a) Given that

$$x = \sec \frac{y}{2}, \quad 0 \leq y < \pi,$$

show that

$$\frac{dy}{dx} = \frac{2}{x\sqrt{x^2-1}}. \quad (5)$$

(b) Find an equation for the tangent to the curve  $y = \sqrt{3+2\cos x}$  at the point where  $x = \frac{\pi}{3}$ . (6)

---

5.  $f(x) = 5 + e^{2x-3}, \quad x \in \mathbb{R}.$

(a) State the range of  $f$ . (1)

(b) Find an expression for  $f^{-1}(x)$  and state its domain. (4)

(c) Solve the equation  $f(x) = 7$ . (2)

(d) Find an equation for the tangent to the curve  $y = f(x)$  at the point where  $y = 7$ . (4)

---

6. (a) Prove the identity

$$2 \cot 2x + \tan x \equiv \cot x, \quad x \neq \frac{n}{2}\pi, \quad n \in \mathbb{Z}. \quad (5)$$

(b) Solve, for  $0 \leq x < \pi$ , the equation

$$2 \cot 2x + \tan x = \operatorname{cosec}^2 x - 7,$$

giving your answers to 2 decimal places. (6)

---

*Turn over*

7. The functions  $f$  and  $g$  are defined by

$$f : x \rightarrow |2x - 5|, \quad x \in \mathbb{R},$$

$$g : x \rightarrow \ln(x + 3), \quad x \in \mathbb{R}, \quad x > -3.$$

(a) State the range of  $f$ . (1)

(b) Evaluate  $fg(-2)$ . (2)

(c) Solve the equation

$$fg(x) = 3,$$

giving your answers in exact form. (5)

(d) Show that the equation

$$f(x) = g(x)$$

has a root,  $\alpha$ , in the interval  $[3, 4]$ . (2)

(e) Use the iteration formula

$$x_{n+1} = \frac{1}{2} [5 + \ln(x_n + 3)],$$

with  $x_0 = 3$ , to find  $x_1, x_2, x_3$  and  $x_4$ , giving your answers to 4 significant figures. (3)

(f) Show that your answer for  $x_4$  is the value of  $\alpha$  correct to 4 significant figures. (2)

---

**END**