


## Emergency Care

THIRTEENTH EDITION




### CHAPTER 25

#### Bleeding and Shock

Daniel Limmer  
 Michael F. O'Keefe  
 Medical Editor: Edward T. Dickinson, MD, FACP

ALWAYS LEARNING | Emergency Care, 13e | Daniel Limmer | Michael F. O'Keefe | Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON



## Main Components

- PUMP - Heart
- PIPES - Blood vessels
- FLUID - Blood

continued on next slide

ALWAYS LEARNING | Emergency Care, 13e | Daniel Limmer | Michael F. O'Keefe | Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Main Components

- Arteries
  - Carry oxygen-rich blood away from the heart
  - Comprised of thick, muscular walls that enable dilation and constriction
- Veins
  - Carry oxygen-depleted blood rich in carbon dioxide back to the heart
  - Contain one-way valves to prevent back flow of blood

continued on next slide


ALWAYS LEARNING | Emergency Care, 13e | Daniel Limmer | Michael F. O'Keefe | Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Main Components

- Capillaries
  - Microscopic blood vessels
  - Vital exchange site
    - Oxygen, nutrients passed through capillary walls in exchange for carbon dioxide from cells

continued on next slide

ALWAYS LEARNING | Emergency Care, 13e | Daniel Limmer | Michael F. O'Keefe | Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON



## Main Components

- Functions of blood
  - Transportation of oxygen and CO<sub>2</sub>
  - Circulates nutrients
  - Excretion of waste products
  - Protection (WBC's, platelets)
  - Regulation

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Main Components

- Perfusion
  - Adequate circulation of blood throughout body
- Hypoperfusion
  - Inadequate perfusion of body's tissues and organs

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Bleeding

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Types of Bleeding

- Hemorrhage is severe bleeding.
  - Major cause of shock (hypoperfusion) in trauma
- External
- Internal

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## External Bleeding



Three types of external bleeding. Left to right, spurting to steady.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## External Bleeding

- Occurs outside of body after force penetrates skin and lacerates or destroys underlying blood vessels
- Typically visible on surface of the skin
- How much a person bleeds determined by:
  - Size and severity of wound
  - Size and pressure of ruptured vessel
  - Individual's ability to clot

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Massive Hemorrhage

- Arterial bleeding
  - Bright red color
  - Spurting with heartbeat
  - Oxygen rich
  - Most difficult to control

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Massive Hemorrhage

- Venous bleeding
  - Darker in color than arterial bleeding
  - Less pressure than arterial bleeding
  - Volume of blood carried by some veins can create immediately life-threatening hemorrhage

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Other External Hemorrhage

- Capillary bleeding
  - Caused by superficial wounds to surface of skin
  - Slow and oozing
  - Stops spontaneously

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Other External Hemorrhage

- Bleeding can be accelerated by underlying conditions.
  - Prescription medications (ASA, Coumadin, Plavix, Xarelto) designed to limit body's natural ability to form blood clots
- Hypothermia affects body's ability to clot.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Think About It

- How severe is the bleeding? Is it exsanguinating hemorrhage? If so, how does that affect the priorities of treatment?

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Assessment and Care of External Hemorrhage

- Must use Standard Precautions
- Ensure open airway.
- Ensure adequate breathing.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Assessment and Care of External Hemorrhage

- Control bleeding only after assessing and treating prior elements.
  - Be aware of signs or symptoms of shock.
  - Use direct pressure, elevation, hemostatic agent, or a tourniquet to control bleeding.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Direct pressure
  - Apply firm pressure to wound with gloved hand and gauze bandage.
  - Hold pressure until bleeding is controlled.
  - If necessary, add dressings when lower ones are saturated.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Direct pressure
  - Once bleeding is controlled, bandage a dressing firmly in place to form a pressure dressing.
  - Never remove bandages—even when bleeding is controlled.
  - When controlled, check for pulse distal to wound to make sure dressing has not been applied too tightly.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Direct pressure
  - Pressure dressing
    - Place several gauze pads on wound.
    - Hold dressings in place with self-adhering roller bandage wrapped tightly over dressings and above and below wound site.
    - Create enough pressure to control bleeding.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Elevation
  - Elevate injured extremity above level of the heart while applying direct pressure.
  - Do not elevate if musculoskeletal injury, impaled objects in extremity, or spine injury is suspected.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Hemostatic agents
  - Designed to enhance direct pressure's ability to control bleeding
  - Work by applying a material design to absorb liquid portion of blood and leave larger formed elements to clot
  - Originated as powders, but now may include dressings and gauze bandages
  - Manual pressure is always necessary.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding



Hemostatic bandage. © Edward T. Dickinson, MD

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Tourniquet
  - Use if bleeding is uncontrollable by direct pressure.
  - Use only on extremity injuries.
  - Always apply between the wound and the heart.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Tourniquet



The Mechanical Advantage Tourniquet (MAT).

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- Tourniquet
  - Follow manufacturer's instructions.
  - Once applied, do not remove or loosen.
  - Attach notation to patient alerting other providers tourniquet has been applied.



ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Think About It

- Is the current method of bleeding control working? Do you need to move on to a more aggressive step? How would you evaluate this?

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Hemorrhage Control Video



Maintaining the airway and administering high flow oxygen; to r. Prepare the patient for transport and monitor vital signs. ket.

Click on the screenshot to view a video on the subject of controlling bleeding.

[Back to Directory](#)

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Controlling External Bleeding

- A systematic approach to treat uncontrolled external hemorrhage
  - Recommendations from American College of Surgeons
  - Begin with direct pressure.
  - If not controlled, apply tourniquet.
  - If ineffective and wound on trunk or head, apply hemostatic dressing or bandage with direct pressure.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Other Methods of Bleeding Control

- Splinting
  - Stabilizing sharp ends of broken bones
  - Inflatable (air) splints
- Cold application
  - Minimizes swelling, constricts blood vessels, and reduces pain
  - Use in conjunction with other manual techniques.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Special Situations Involving Bleeding

- Head injury
  - From increased intracranial pressure, not direct trauma to ears or nose
  - Stopping bleeding only increases intracranial pressure.
  - Allow drainage to flow freely, using gauze pad to collect it.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Special Situations Involving Bleeding

- Nosebleed (epistaxis)
  - Have patient sit and lean forward.
  - Apply direct pressure to fleshy portion of nostrils.
  - Keep patient calm and quiet.
  - Do not let patient lean back.
  - If patient becomes unconscious, place patient in recovery position and be prepared to suction.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Internal Bleeding

- Damage to internal organs and large blood vessels can result in loss of a large quantity of blood in short time.
- Blood loss commonly cannot be seen.
- Severe blood loss can even result from injuries to extremities.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment

- Blunt Trauma
  - Falls
  - Motor-vehicle or motorcycle crashes
  - Auto-pedestrian collisions
  - Blast injuries

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment

- Penetrating Trauma
  - Gunshot wounds
  - Stab wounds
  - Impaled objects

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment

- Signs of internal bleeding
  - Injuries to surface of body
  - Bruising, swelling, or pain over vital organs
  - Painful, swollen, or deformed extremities
  - Bleeding from mouth, rectum, or vagina

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment

- Signs of internal bleeding
  - Tender, rigid, or distended abdomen
  - Vomiting coffee-grounds like substance or bright red vomitus
  - Dark, tarry stools or bright red blood in stool
  - Signs and symptoms of shock

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment



Bruising is one sign of internal bleeding. © Edward T. Dickinson, MD

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Care

- Maintain ABCs.
- Administer high-concentration oxygen by nonrebreather mask.
- Control any external bleeding.
- Preserve body temperature.
- Provide prompt transport to appropriate medical facility.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Shock (Hypoperfusion)

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Shock (Hypoperfusion)

- Inadequate tissue perfusion
- It also causes inadequate removal of waste products from cells.


ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Causes of Shock

- Failure of any component of circulatory system
  - Heart
    - Loses ability to pump
  - Blood vessels
    - Dilate, making too large a "container" to fill
  - Blood
    - Loses volume from bleeding

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Causes of Shock



## Severity of Shock

- Compensated shock
  - Body senses the decrease in perfusion and attempts to compensate for it.
  - Early signs of shock
- Decompensated shock
  - Begins when the body can no longer compensate for low blood volume or lack of perfusion
  - Late signs of shock

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Types of Shock

- Hypovolemic shock
  - Decreased volume of circulating blood and plasma
  - Called hemorrhagic shock if caused by uncontrolled bleeding (internal or external)
  - Can be caused by burns or crush injuries or severe dehydration

continued on next slide

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Types of Shock

- Cardiogenic shock
  - Patients suffering myocardial infarction
  - Results from inadequate pumping of blood by heart, decreasing strength of contractions
  - Heart's electrical system may malfunction, causing heartbeat that is too slow, too fast, or irregular.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON



## Types of Shock

- Neurogenic shock
  - Uncontrolled dilation of blood vessels because of nerve paralysis
  - No blood loss, but vessels dilated so much that blood volume cannot fill them
  - Rarely seen in the field

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Pediatric Note

- Infants and children
  - Efficient compensating mechanisms maintain blood pressure until half of volume is depleted
- Potential for shock must be recognized and treated before tell-tale signs appear

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Assessment

- Progression of the signs and symptoms of shock
  - Altered mental status
  - Pale, cool, clammy skin
  - Nausea and vomiting
  - Vital sign changes
  - Late signs of shock include thirst, dilated pupils, and sometimes cyanosis around lips and nail beds.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Emergency Care for Shock

- Transportation is an intervention.
  - Every minute between the time of injury and the patient's getting to an operating suite is, in fact, like gold to the patient—and to his chances of survival.
- Goal is *platinum 10 minutes* at the scene.
- Prevent heat loss, coagulopathy, and further blood loss.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Care

- Maintain open airway and assess respiratory rate
  - Address inadequate breathing immediately and aggressively.
  - If patient is breathing adequately, apply high-concentration oxygen by nonrebreather mask.

continued on next slide

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Care

- Control any external bleeding.
- If pelvic fracture is suspected, use pelvic binding device.



ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Patient Care

- Splint any suspected bone or joint injuries.
- Prevent loss of body heat.
- Transport patient immediately.
- Speak calmly and reassure throughout assessment and care.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Shock Animation



Click on the screenshot to view an animation on the subject of shock.

[Back to Directory](#)

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Chapter Review

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Chapter Review

- Almost all external bleeding can be controlled by direct pressure and elevation. When these do not work, apply a tourniquet if bleeding is on an extremity or a hemostatic dressing if the bleeding is from the head or torso.
- Emergency care for internal bleeding is based on prevention and treatment of shock.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Chapter Review

- Early signs of shock are often restlessness, anxiety, pale skin, rapid pulse, and respirations.
- If shock is uncontrolled, the patient's blood pressure falls, late sign of shock.
- Signs and symptoms may not be evident early in the call, so treatment based on MOI may be lifesaving.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Chapter Review

- Treat shock by maintaining the airway, administering high-concentration oxygen, controlling bleeding, and keeping the patient warm. One of most important treatments is early recognition of shock and immediate transport to a hospital.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Remember

- The circulatory system is designed to ensure adequate perfusion of body tissues.
- The classification of hemorrhage is directly related to the type of vessel ruptured and the pressure within that vessel.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Remember

- Treatment of external hemorrhage includes progression through the following steps: direct pressure, elevation, tourniquet application, use of hemostatic agents.
- Internal bleeding is impossible to evaluate. The most appropriate treatment must be rapid transport to an appropriate facility.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Remember

- Shock develops if the heart fails, blood volume is lost, or blood vessels dilate, resulting in inadequate perfusion.
- Signs of shock reflect the body's attempts at compensating for inadequate perfusion.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Remember

- The most significant treatment for the shock patient is early recognition and prompt transport to a hospital where the patient will receive definitive care.

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Questions to Consider

- What can I use for a tourniquet that will control bleeding but not damage tissue?
- When treating a patient with shock, what should I do at the scene and what should I do en route to the hospital?

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Questions to Consider

- Is a patient with pale, cool skin, tachycardia, and rapid, shallow respirations in shock or just under stress? How will continuing assessment help in making that decision?

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Critical Thinking

- A patient has been involved in a motor-vehicle collision. There is considerable damage to the vehicle. The steering column and wheel are badly deformed. The patient complains of a "sore chest." You note no external bleeding.

*continued on next slide*

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON

## Critical Thinking

- The patient's vital signs are pulse 116, respirations 20, blood pressure 106/70. How would you proceed to assess and care for this patient?

ALWAYS LEARNING Emergency Care, 13e Daniel Limmer | Michael F. O'Keefe Copyright © 2016, 2012, 2009 by Pearson Education, Inc. All Rights Reserved. PEARSON