

Calculus 3 - Volumes

In Calculus 1 we considered the area problem. Find the area under the curve $y = f(x)$ on the interval $[a, b]$. To do this, we broke the interval up into smaller segments, approximated the area on each segment with a rectangle, added the rectangles up, and then took the limit as the number of rectangles went to infinity and the thickness of each rectangle went to zero.




Figure 1: $y = f(x)$

Mathematically: We subdivide the interval

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < \cdots < x_n = b. \quad (1)$$

Let

$$\Delta x_i = x_{i-1} - x_i. \quad (2)$$

Pick x_i^* so that

$$x_i^* \in [x_{i-1}, x_i]. \quad (3)$$

Height of the i^{th} rectangle

$$h_i = f(x_i^*). \quad (4)$$

Area of this rectangle

$$A_i = f(x_i^*)\Delta x_i. \quad (5)$$

Add up the rectangles

$$\sum_{i=1}^n A_i = \sum_{i=1}^n f(x_i^*)\Delta x_i. \quad (6)$$

Then take the limit so

$$A = \lim_{\substack{n \rightarrow \infty \\ \Delta x_i \rightarrow 0}} \sum_{i=1}^n f(x_i^*)\Delta x_i \quad (7)$$

and we gave this Riemann sum a name - a definite integral

$$A = \int_a^b f(x) dx = \lim_{\substack{n \rightarrow \infty \\ \Delta x_i \rightarrow 0}} \sum_{i=1}^n f(x_i^*)\Delta x_i. \quad (8)$$

So now we consider the volume problem. Find the volume under the surface $z = f(x, y)$ on the interval $[a, b] \times [c, d]$. The process is the same thing as in the area problem. We approximate the volume with small rectangular boxes.

Mathematically: Subdivide the interval

$$\begin{aligned} a &= x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < \cdots < x_m = b \\ c &= y_0 < y_1 < y_2 < \cdots < y_{j-1} < y_j < \cdots < y_n = d. \end{aligned} \quad (9)$$

Let

$$\Delta x_i = x_{i-1} - x_i, \quad \Delta y_j = y_{j-1} - y_j, \quad (10)$$




Figure 2: Grid

Pick (x_i^*, y_j^*) so that

$$(x_i^*, y_j^*) \in [x_{i-1}, x_i] \times [y_{j-1}, y_j]. \quad (11)$$

Height of the the rectangle box

$$h_{ij} = f(x_i^*, y_j^*). \quad (12)$$

The volume of this rectangle box is

$$V_{ij} = f(x_i^*, y_j^*) \Delta x_i \Delta y_j. \quad (13)$$

Add up the boxes

$$\sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \Delta x_i \Delta y_j. \quad (14)$$

Then take the limit so


$$\lim_{\substack{m,n \rightarrow \infty \\ \Delta x_i, \Delta y_j \rightarrow 0}} \sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \Delta x_i \Delta y_j \quad (15)$$

and we give this Riemann sum a name - a double integral

$$\int_c^d \int_a^b f(x, y) dx dy = \lim_{\substack{m,n \rightarrow \infty \\ \Delta x_i, \Delta y_j \rightarrow 0}} \sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \Delta x_i \Delta y_j \quad (16)$$

Example 1.

Find the volume under $z = 1$ for $[0, 2] \times [0, 3]$. Well, we see right away that the volume is 6. Let us use the double integral. So here



$$V = \int_0^3 \int_0^2 1 dx dy. \quad (17)$$

As with partial derivatives, when integrating one variable, we hold the

other constant, we do the same with double integral.

$$\begin{aligned} V &= \int_0^3 \left(\int_0^2 1 dx \right) dy \\ &= \int_0^3 \left(x \Big|_0^2 \right) dy \\ &= \int_0^3 2 dy \\ &= 2y \Big|_0^3 \\ &= 6. \end{aligned} \tag{18}$$

Switching the Order of Limits

We could also have done

$$\begin{aligned} V &= \int_0^2 \left(\int_0^3 1 dy \right) dx \\ &= \int_0^2 \left(y \Big|_0^3 \right) dx \\ &= \int_0^2 3 dx \\ &= 3x \Big|_0^2 \\ &= 6. \end{aligned} \tag{19}$$

Integrating when f is not constant

Consider

$$\begin{aligned} V &= \int_0^3 \int_{-1}^1 12x^2y \, dx dy \\ &= \int_0^3 4x^3y \Big|_{x=-1}^{x=1} dy \\ &= \int_0^3 8y \, dy \\ &= 4y^2 \Big|_{y=0}^{y=3} \\ &= 36. \end{aligned} \tag{20}$$

Switching limits

$$\begin{aligned} V &= \int_{-1}^1 \int_0^3 12x^2y \, dy dx \\ &= \int_{-1}^1 6x^2y^2 \Big|_0^3 dy \\ &= \int_{-1}^1 54x^2 \, dy \\ &= 18x^3 \Big|_{-1}^1 \\ &= 36. \end{aligned} \tag{21}$$

In fact, if f is continuous on $[a, b] \times [c, d]$ then


$$\int_a^b \int_c^d f(x, y) \, dy dx = \int_c^d \int_a^b f(x, y) \, dx dy \tag{22}$$

Integration over non constant regions


Suppose we wish to set up the double integral

$$\iint_R f(x,y) dydx \quad (23)$$

where R is the region below (the lines $y = 0$, $y = x$ and $x = 1$)



To get an idea on how to do this let us first consider the problem when the region is a rectangular box.




So we have

$$\int_a^b \int_c^d f(x, y) dy dx = \int_a^b \left(\int_c^d f(x, y) dy \right) dx \quad (24)$$

In the round bracket x is fixed and y moves from $y = c$ to $y = d$. Now in the triangular region when x is fixed, then y moves from $y = 0$ to $y = x$ and so the limits of integration are

$$\int_a^b \left(\int_0^x f(x, y) dy \right) dx \quad (25)$$



Now as the rectangle moves, it moves from $x = 0$ to $x = 1$ and these are the outside limits and so

$$\int_0^1 \left(\int_0^x f(x, y) dy \right) dx \quad (26)$$

or simply

$$\int_0^1 \int_0^x f(x, y) dy dx \quad (27)$$

Example 2. Evaluate

$$\int_0^1 \int_0^x 6y^2 - x^3y \, dydx \quad (28)$$

Soln. We first integrate wrt y holding x fixed. So

$$\int_0^1 2y^3 - \frac{x^3y^2}{2} \Big|_{y=0}^{y=x} dx \quad (29)$$

Then substitute in the limits

$$\int_0^1 \left(2x^3 - \frac{x^3x^2}{2} \right) - \left(2 \cdot 0^3 - \frac{x^3 \cdot 0^2}{2} \right) dx \quad (30)$$


Then integrate one more time

$$\frac{x^4}{2} - \frac{x^6}{12} \Big|_{x=0}^{x=1} = \frac{1}{2} - \frac{1}{12} = \frac{5}{12}. \quad (31)$$

In general


In general we have

$$\int_a^b \int_{g(x)}^{h(x)} f(x,y) \, dydx \quad (32)$$



and

$$\int_c^d \int_{G(y)}^{H(y)} f(x, y) dx dy \quad (33)$$



Example 1. Find the volume under the plane $2x + y + z = 2$ for $x, y, z \geq 0$.

The volume is shown in fig. 1 in the first octant.

so the volume is

$$V = \int_0^1 \int_0^{2-2x} (2 - 2x - y) dy dx \quad (34)$$

or

$$V = \int_0^2 \int_0^{\frac{2-y}{2}} (2 - 2x - y) dx dy \quad (35)$$




Figure 3: Surface and region of integration

and we integrate so

$$\begin{aligned}
 V &= \int_0^1 \int_0^{2-2x} (2 - 2x - y) dy dx \\
 &= \int_0^1 \left(2y - 2xy - \frac{1}{2}y^2 \right) \Big|_0^{2-2x} dx \\
 &= \int_0^1 2(2 - 2x) - 2x(2 - 2x) - \frac{1}{2}(2 - 2x)^2 dx \quad (36) \\
 &= \int_0^1 2 - 4x - 2x^2 dx \\
 &= 2x - 2x^2 - \frac{2}{3}x^3 \Big|_0^1 = \frac{2}{3}.
 \end{aligned}$$

Example 2. pg 987, #26 Find the volume under the parabolic cylinder $z = 4 - y^2$ on the region bound by $y = x$, $x = 0$ and $y = 2$




Figure 4: Volume and region of integration

$$V = \int_0^2 \int_x^2 (4 - y^2) dy dx \quad (37)$$

or

$$V = \int_0^2 \int_0^y (4 - y^2) dx dy \quad (38)$$

Volume

$$\begin{aligned} V &= \int_0^2 \int_0^y (4 - y^2) dx dy \\ &= \int_0^2 4x - xy^2 \Big|_0^y dy \\ &= \int_0^2 4y - y^3 \Big|_0^y dy \\ &= 2y^2 - \frac{1}{4}y^4 \Big|_0^2 = 8 - 4 = 4 \end{aligned} \quad (39)$$