

VI. References

1. Anthony Gauthier, MD, et al. Relative Efficacy of Sildenafil Compared to Other Treatment Options for Erectile Dysfunction [South Med J 93(10):962-965, 2000. © 2000 Southern Medical Association]
2. Urban RJ, Bodenburg YH, Gilkison C, Foxworth J, Coggan A.R.Wolfe RR, Ferrando A. Testosterone administration to elderly men increases skeletal muscle strength and protein synthesis. Am J Physiol 1995 Nov; 269(5 Pt 1): E820-6
3. Brown WA, Monti PM, Corriveau DP . Serum testosterone and sexual activity and interest in men. Arch Sex Behav 1978 Mar; 7(2): 97-103
4. Gooren LJ. Androgen levels and sex functions in testosterone-treated hypogonadal men. Arch Sex Behav 1987 Dec;16(6):463-73
5. Lugg JA, Rajfer J, Gonzalez-Cadavid NF. Dihydrotestosterone is the active androgen in the maintenance of nitric oxide-mediated penile erection in the rat. Endocrinology 1995 Apr;136(4):1495-501
6. Anderson RA, Bancroft J, Wu FC. The effects of exogenous testosterone on sexuality and mood of normal men. J Clin Endocrinol Metab 1992 Dec;75(6): 1503-7
7. Korenman SG, Viosca S, Garza D, Guralnik M, Place V, Campbell P, Davis SS . Androgen therapy of hypogonadal men with transscrotal testosterone systems. American Journal of Medicine 1987 Sep;83(3):471-8
8. Kryger, AH. The Effect of Administration of a Testosterone Cream in Hypogonadal men. (unpublished). Study available on www.TestoCreme.com .