





PATIENT ASSESSMENT/MANAGEMENT - MEDICAL

Skill Sheet 1 TESTED

Possible Points	Candidate:			Examiner:_			
Takes or varbalizes appropriate PPE precautions SCENE SIZE-UP Determines the scene/situation is safe Determines the mechanism of finjury/nature of lilness 1 Determines the number of patients 1 Determines the number of patients 1 1 Requests additional EMS assistance if necessary 1 1 Considers stabilization of the spine 1 1 PRIMARY SURVEY/RESUSCITATION Verbalizes the general impression of the patient Determines responsiveness/level of consciousness (AVPU) 1 Determines responsiveness/level of consciousness (AVPU) Determines responsiveness/level of consciousness (AVPU) 1 Determines responsiveness/level of consciousness (AVPU) 3 Assessess affine (lementing of point) -Assessesses/mortine and makes treatment/transport decision 1 HISTORY TAKING HISTORY TAKING HISTORY TAKING HISTORY TAKING HISTORY TAKING Past medical history -Past pertinent history (1 point) -Past medical history -Past pertinent history (1 point) -Past medical history -Past pertinent history (1 point) -Past pertinent history	Date:			Signature:_			
SCENE SIZE-UP Determines the scene/situation is safe 1	Actual Time Started:	_					
SCENE SIZE-UP Determines the scene/situation is safe 1	Takes or verbalizes appropriate F	PE precautions				1	
Determines the mechanism of injury/nature of illness Determines the number of patients Requests additional EMS assistance if necessary 1 Considers stabilization of the spine PRIMARY SURVEY/RESUSCITATION Verbalizes the general impression of the patient Determines chief compliant/apparent life-threats Assesses airway and breathing -Assesses airway and breathing -Assesses airway and breathing -Assesses airway and breathing -Assesses circulation Assesses circulation -Assesses circulation -Assesses sinculation -Assesses sinculation -Assessess sinculation -Assessess sinculation -Assessess sinculation -Assessess sinculation -Assesses sinculation -Provocation (1 point) -Provocation (1 point							
Determines the mechanism of injury/nature of illness Determines the number of patients Requests additional EMS assistance if necessary 1 Considers stabilization of the spine PRIMARY SURVEY/RESUSCITATION Verbalizes the general impression of the patient Determines chief compliant/apparent life-threats Assesses airway and breathing -Assesses airway and breathing -Assesses airway and breathing -Assesses airway and breathing -Assesses circulation Assesses circulation -Assesses circulation -Assesses sinculation -Assesses sinculation -Assessess sinculation -Assessess sinculation -Assessess sinculation -Assessess sinculation -Assesses sinculation -Provocation (1 point) -Provocation (1 point	Determines the scene/situation is	safe				1	
Determines the number of patients 1	Determines the mechanism of inj	ury/nature of illness				1	
Requests additional EMS assistance if necessary Considers stabilization of the spine PRIMARY SURVEY/RESUSCITATION Verbalizes the general impression of the patient Determines responsiveness/level of consciousness (AVPU) Determines chief complaint/apparent life-threats 1 Assesses airway and breathing -Assesses airway and breathing -Assesses circulation -Assesses circulation -Assesses (I (point) -Assesses (1	
PRIMARY SURVEY/RESUSCITATION						1	
Verbalizes the general impression of the patient 1 Determines responsiveness/level of consciousness (AVPU) 1 Determines responsiveness/level of consciousness (AVPU) 1 Determines responsiveness/level of consciousness (AVPU) 1 Determines chief complaint/apparent life-threats 1 Assesses circulation Assesses circulati						1	
Determines responsiveness/level of consciousness (AVPU) 1 1 1 1 1 1 1 1 1	PRIMARY SURVEY/RESUSCITA	ATION					
Determines responsiveness/level of consciousness (AVPU) 1 1 1 1 1 1 1 1 1	Verbalizes the general impression	n of the patient				1	
Determines chief complaint/apparent life-threats Assesses airway and breathing -Assesses airway and breathing -Assesses circulation -Assesses (roulation -Assesses (roulation) -Assesses skin [either skin color, temperature or condition] (1 point) Identifies patient priority and makes treatment/transport decision HISTORY TAKING HISTORY TAKING HISTORY TAKING HISTORY TAKING -Quality (1 point) -Radiation (1 point) -Radiation (1 point) -Radiation (1 point) -Provocation (1 point) -Provocation (1 point) -Past pertinent history (1 point) -Medications of associated signs and symptoms related to OPQRST (2 points) -Allergies (1 point) -Medications (1 point) -Last oral intake (1 point) -Severity (1 point) -Events leading to present illness (1 point) -Medications (1 point) -Last oral intake (1 point) -SecondDARY ASSESSMENT -Cardiovascular -Neurological -Neurological -Integumentary -Reproductive 5 -Pulmonary -Nusculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 pointeach) 4 States field impression of patient Interventions (verbalizes proper interventions/treatment) REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit Actual Time Ended: TOTAL 42 CRITICAL CRITERIA Failure to to take or verbalize appropriate PPE precautions Failure to texical and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy Failure to voice and ultimately provide appropriate oxygen therapy			VPU)			1	
-Assessment (1 point) -Assures adequate ventilation (1 point) -Initiates appropriate oxygen therapy (1 point) 3 Assesses circulationChecks pulse (1 point) -Checks pulse (1 point) 3 -Assesses skin [either skin color, temperature or condition] (1 point) -Checks pulse (1 point) 1 Identifies patient priority and makes treatment/transport decision 1 1 IHISTORY TAKING	Determines chief complaint/appa	rent life-threats				1	
Assesses circulation Assesses kin [either skin color, temperature or condition] (1 point) Identifies patient priority and makes treatment/transport decision IISTORY TAKING History of the present illness Onset (1 point) Provocation (1 point) Provocation (1 point) Provocation (1 point) Assesses as a service (1 point) Past medical history Allergies (1 point) Assesses affected body part/system Allergies (1 point) Assesses affected body part/system Cardiovascular Neurological Pulmonary Ausculoskeletal Fulmonary Ausculoskeletal Interventions (1 point) Provides and value (1 point) Assesses affected body part/system Bodo pressure (1 point) Pulse (1 point) Prulse (1 point) Assesses affected body part/system Cardiovascular Neurological Fulmonary Ausculoskeletal Fulmonary Ausculoskeletal Fulmonary Ausculoskeletal	Assesses airway and breathing						
-Assesses/controls major bleeding (1 point) -Checks pulse (1 point) 3 -Assesses skin [either skin color, temperature or condition] (1 point) Identifies patient priority and makes treatment/transport decision HISTORY TAKING History of the present illness -Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) -SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -Gi/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 Interventions [verbalizes proper interventions/treatment] 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to voice and ultimately provide appropriate oxygen therapy Failure to soice and ultimately provide appropriate oxygen therapy Failure to soice and ultimately provide appropriate oxygen therapy Failure to sesses/provide adequate ventilation	-Assessment (1 point) -A	ssures adequate vent	tilation (1 point)	-Initiates a	appropriate oxygen therapy (1 point)	3	
-Assesses skin [either skin color, temperature or condition] (1 point) Identifies patient priority and makes treatment/transport decision HISTORY TAKING History of the present illness -Onset (1 point) -Quality (1 point) -Time (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICAL CRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions -Failure to initiate or call for transport of the patient within 15 minute time limit Failure to voice and ultimately provide appropriate oxygen therapy -Failure to sesses/provide adequate ventilation	Assesses circulation						
Identifies patient priority and makes treatment/transport decision HISTORY TAKING History of the present illness -Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to intitate or call for transport of the patient within 15 minute time limit Failure to voice and ultimately provide appropriate oxygen therapy Failure to sesses/provide adequate ventilation	-Assesses/controls major bleedin	g (1 point)		-Checks p	ulse (1 point)	3	
HISTORY TAKING History of the present illness -Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) 8 -Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Allergies (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 1 Interventions (verbalizes proper interventions/treatment) 1 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 1 Provides accurate verbal report to arriving EMS unit 1 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to voice and ultimately provide appropriate oxygen therapy Failure to sesses/provide adequate ventilation	-Assesses skin [either skin color,	temperature or condi-	tion] (1 point)				
History of the present illness -Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) -Medications (1 point) -Last oral intake (1 point) -EsconDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions (verbalizes proper interventions/treatment) 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to take or verbalize appropriate PPE precautions Failure to voice and ultimately provide appropriate oxygen therapy Failure to sessess/provide adequate ventilation	Identifies patient priority and mak	es treatment/transpor	t decision			1	
-Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Time (1 point) 8 -Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended:	HISTORY TAKING						
-Provocation (1 point) -Radiation (1 point) -Time (1 point) -Secrifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 1 Interventions [verbalizes proper interventions/treatment] 1 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 1 Provides accurate verbal report to arriving EMS unit 1 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation							
-Clarifying questions of associated signs and symptoms related to OPQRST (2 points) Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA - Failure to initiate or call for transport of the patient within 15 minute time limit 5 - Failure to determine scene safety before approaching patient 5 - Failure to voice and ultimately provide appropriate oxygen therapy - Failure to sassess/provide adequate ventilation							
Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended:						8	
-Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) 5 -Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICAL CRITERIA - Failure to initiate or call for transport of the patient within 15 minute time limit - Failure to determine scene safety before approaching patient - Failure to voice and ultimately provide appropriate oxygen therapy - Failure to assess/provide adequate ventilation		d signs and symptom	s related to OPQ	RST (2 point	s)		
-Medications (1 point) -Last oral intake (1 point) SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to intitate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation						_	
SECONDARY ASSESSMENT Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICAL CRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation				-Events lea	iding to present illness (1 point)	5	
Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICAL CRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation		-Last oral intake (1 point)				
-Cardiovascular -Neurological -Integumentary -Reproductive 5 -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation							
Pulmonary -Musculoskeletal -GI/GU -Psychological/Social VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation			•			_	
VITAL SIGNS -Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation	The state of the s	,		itary		5	
-Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each) 4 States field impression of patient 1 Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended:		lusculoskeletai	-GI/GU		-Psychological/Social		
States field impression of patient Interventions [verbalizes proper interventions/treatment] REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition Provides accurate verbal report to arriving EMS unit Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation		Dulas (d. naint)		Despisato	number and suplified (1 point apply)	1	
Interventions [verbalizes proper interventions/treatment] 1 REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition 1 Provides accurate verbal report to arriving EMS unit 1 Actual Time Ended:				-Respirator	y rate and quality (1 point each)		
REASSESSMENT Demonstrates how and when to reassess the patient to determine changes in condition Provides accurate verbal report to arriving EMS unit Actual Time Ended:			41				
Demonstrates how and when to reassess the patient to determine changes in condition Provides accurate verbal report to arriving EMS unit Actual Time Ended:		iterventions/treatmen	tj				
Provides accurate verbal report to arriving EMS unit Actual Time Ended:			a datamaina ahaa	andit	ina	1	
Actual Time Ended: TOTAL 42 CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation			determine char	iges in condit	1011	_	
CRITICALCRITERIA Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation					TOTAL		
Failure to initiate or call for transport of the patient within 15 minute time limit Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation						*******	
Failure to take or verbalize appropriate PPE precautions Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation		and of the metions with	in 15 minute time	line it			
Failure to determine scene safety before approaching patient Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation				IIIIII			
Failure to voice and ultimately provide appropriate oxygen therapy Failure to assess/provide adequate ventilation							
Failure to assess/provide adequate ventilation		•	•				
			gen merapy				
	· (1985)		ciated with airway	breathing he	morrhage or shock		

__ Failure to differentiate patient's need for immediate transportation versus continued assessment or treatment at the same

Performs secondary examination before assessing and treating threats to airway, breathing and circulation

Orders a dangerous or inappropriate intervention
Failure to provide accurate report to arriving EMS unit
Failure to manage the patient as a competent EMT

Exhibits unacceptable affect with patient or other personnel
 Uses or orders a dangerous or inappropriate intervention



PATIENT ASSESSMENT/MANAGEMENT – MEDICAL Oral Glucose Administration

Skill Sheet Supplement 1 TESTED/VERIFIED

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Perform blood glucose check			
Prepare glucometer and supplies		1	
Cleanse site		1	
Lance site		1	
		1	
Apply direct pressure to site		1	
Read and interpret results		1 1	
Determine appropriate indications for glucose administration			
Level of consciousness		1	
Pertinent past medical history		1	
Contact Medical Command if patient condition indicates		1	
Confirm expiration date of oral glucose		1	
Confirm the rights of drug administration: Right Patient ☐ Right Route ☐ Right Drug ☐ Right Dose ☐	Right Time	1 point each	
Explain the procedure to the patient		1	
Place oral glucose between cheek and gum		1	
Recheck patient's blood glucose level within 5 minutes of administrati	on	1	
If no improvement contact medical command and request ALS back u	ıp	1	
Document the procedure		1	
Actual Time Ended:	TOTAL	. 21	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to consult medical command at appropriate times Failure to confirm at least three of the "Rights" of medication add Failure to determine blood glucose level prior to, or following, or Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	ministration al glucose administration	,	



PATIENT ASSESSMENT/MANAGEMENT – MEDICAL Nitroglycerin Administration

Skill Sheet Supplement 2 TESTED/VERIFIED

Candidate:	Examiner:			
Date:	Signature:			
Actual Time Started:			Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions			1	
Determine appropriate indications for glucose administration				
Confirm patient allergies			1	
Determine if the patient has self-administered nitroglycerine price	or to EMS arrival		1	
Confirm patient's blood pressure is ≥ 100 systolic			1	
Contact Medical Command			1	
Confirm expiration date of nitroglycerine			1	
Confirm the rights of drug administration: Right Patient ☐ Right Route ☐ Right Drug ☐ Right Do	ose Right Time	'n	1 point each	
Explain the procedure and possible side effects to the patient			1	
Place the patient in a comfortable position			1	
Place one nitroglycerine tablet or administer one pump of liquid	nitroglycerine under the tongue		1	
Instruct the patient to allow the medication to absorb			1	
Recheck the patients blood pressure within 3 – 5 minutes of ad	ministration		1	
If no improvement contact medical command and repeat proces	dure as directed		1	
Document the procedure			1	
Actual Time Ended:		TOTAL	18	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to consult medical command at appropriate times Failure to confirm at least three of the "Rights" of medication Failure to determine patient's blood pressure prior to, or foll Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention				



PATIENT ASSESSMENT/MANAGEMENT – MEDICAL Nebulized Medication Administration

Skill Sheet Supplement 3 TESTED/VERIFIED

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	(4)
Properly determines the need for nebulized medications		1	
Assess the patient's ability to utilize a nebulizer		1	
Confirm patient allergies		1	
Confirm patient's heart rate is ≤ 130 for adults and ≤ 150 in pediatrics		1	
Confirm expiration date of medication		1	
Confirm the rights of drug administration: Right Patient Right Route Right Drug Right Dose	Right Time	1 point each	
Prepare the Nebulizer			
Assemble Nebulizer		1	
Add appropriate medication		1	
Connect the mouthpiece		1	
Attach oxygen to the nebulizer flowing at 8 – 10 liters per minute		1	
Explain the procedure and possible side effects to the patient		1	
Place the patient in a sitting up position		1	
Administer Medication			
Instruct the patient to hold the nebulizer with lips sealed around the mo	uthpiece	1	
Instruct the patient to breath as deeply as possible at a normal rate		1	
Continue administration until all medication has been utilized		1	
Monitor patient's condition and vital signs after administration		1	
If no improvement contact medical command for additional treatment a	s directed	1	
Document the procedure		1	
Actual Time Ended:	TOTAL	23	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to consult medical command at appropriate times Failure to confirm at least three of the "Rights" of medication adr Failure to administer all medication Failure to monitor the patient's condition and vital signs Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	ninistration		



PATIENT ASSESSMENT/MANAGEMENT – MEDICAL Epinephrine Auto-Injector Administration

Skill Sheet Supplement 4 TESTED/VERIFIED

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Properly determines the need for medication		1	
Consults with Medical Command		1	
Confirm patient allergies		1	
Confirm expiration date of medication		1	
Confirm the rights of drug administration: Right Patient ☐ Right Route ☐ Right Drug ☐ Right Dose ☐	Right Time □	1 point each	
Explain the procedure and possible side effects to the patient		1	
Remove the cap from the Auto-Injector		1	
Expose the thigh area (may verbalize)		1	
Cleanse the area		1	
In a smooth, firm fashion push the auto injector into the thigh until a c	ick is heard	1	
Hold the auto injector against the thigh for 10 seconds		1	
Properly dispose of the auto injector in a sharps container		1	
Monitor patient's condition and vital signs after administration		1	
If no improvement, contact medical command for additional treatment	as directed	1	
Document the procedure		1	
Actual Time Ended:	TOTAL	20	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to consult medical command at appropriate times Failure to confirm at least three of the "Rights" of medication ac Failure to monitor the patient's condition and vital signs Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	dministration		



PATIENT ASSESSMENT/MANAGEMENT – MEDICAL Epinephrine 1:1000 Ampule Administration

Skill Sheet Supplement 5 TESTED/VERIFIED

Candidate:	_ Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Properly determines the need for medication		1	
Consults with Medical Command for orders		1	
Confirm patient allergies		1	
Confirm the rights of drug administration: Right Patient ☐ Right Route ☐ Right Drug ☐ Right Dos	e	1 point each	
Explain the procedure and possible side effects to the patient		1	
Selects proper medication and concentration		1	
Checks medication for cloudiness or discoloration		1	
Selects proper needle and syringe		1	
Confirm expiration date of medication		1	
Cleans the neck of the ampule		1	
Opens ampule properly snapping it at the break line while directir	ng the action away from the patient and others	1	
Withdraw the medication utilizing the prepared syringe and needl	e	1	
Verify the correct dosage of medication once its withdrawn from t	he ampule	1	
Tap the barrel of the syringe to remove excess air bubbles		1	
Select and cleanse the appropriate administration site		1	
Penetrates the muscle at a 90° angle		1	
Aspirated for blood return prior to injection		1	
Injects medication and removes needle in the same 90° motion		1	
Properly disposes of needle in a sharps container		1	
Monitor patient's condition and vital signs after administration		1	
If no improvement contact medical command for additional treatm	nent as directed	1	
Document the procedure		1	
Actual Time Ended:	TOTAL	_ 27	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to consult medical command at appropriate times Failure to appropriate medication and concentration Failure to confirm at least three of the "Rights" of medication a Failure to select appropriate needle and syringe Failure to properly cleanse injection site Failure to aspirate for blood return prior to medication adminis Failure to monitor the patient's condition and vital signs Failure to properly dispose of needle Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention			







Skill Sheet 2 **TESTED**

PATIENT ASSESSMENT/MANAGEMENT - TRAUMA

Candidate: Examiner:		
Date:Signature:		
Actual Time Started:Note: Areas denoted by "**" may be integrated within sequence of Primary Survey/Resuscitation	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
SCENE SIZE-UP		
Determines the scene/situation is safe	1	
Determines the mechanism of injury/nature of illness	1	
Determines the number of patients	1	
Requests additional EMS assistance if necessary	1	
Considers stabilization of the spine	1	
PRIMARY SURVEY/RESUSCITATION		
Verbalizes general impression of the patient	1	
Determines responsiveness/level of consciousness	1	
Determines chief complaint/apparent life-threats	1	
Airway	200	
-Opens and assesses airway (1 point) -Inserts adjunct as indicated (1 point)	2	
Breathing		
-Assures adequate ventilation (1 point) -Initiates appropriate oxygen therapy (1 point) -Manages any injury which may compromise breathing/ventilation (1 point)	4	
Circulation -Checks pulse (1point) -Assess skin [either skin color, temperature or condition] (1 point) -Assesses for and controls major bleeding if present (1 point) -Initiates shock management [positions patient properly, conserves body heat] (1 point)	4	
Identifies patient priority and makes treatment/transport decision (based upon calculated GCS)	1	_
HISTORY TAKING		
Obtains baseline vital signs [must include BP, P and R] (1 point)	1	
Attempts to obtain SAMPLE history	1	
SECONDARY ASSESSMENT		
Head -Inspects and palpates scalp and ears (1 point) ** -Inspects mouth**, nose** and assesses facial area (1 point)	3	
Neck** -Checks position of trachea (1 point) -Checks jugular veins (1 point) -Palpates cervical spine (1 point)	3	
Chest** -Inspects chest (1 point) -Palpates chest (1 point) -Auscultates chest (1 point)	3	
Abdomen/pelvis** -Inspects and palpates abdomen (1 point) -Verbalizes assessment of genitalia/perineum as needed (1 point)	3	
Lower extremities** -Inspects, palpates and assesses motor, sensory and distal circulatory functions (1 point/leg)	2	
Upper extremities -Inspects, palpates and assesses motor, sensory and distal circulatory functions (1 point/arm)	2	
Posterior thorax, lumbar and buttocks** -Inspects and palpates posterior thorax (1 point) -Inspects and palpates lumbar and buttocks areas (1 point)	2	
Manages secondary injuries and wounds appropriately	1	
REASSESSMENT		
Demonstrates how and when to reassess the patient	1	
Actual Time Ended: TOTAL	42	
CRITICAL CRITERIA Failure to initiate or call for transport of the natient within 10 minute time limit		

Failure to initiate of	r call for transport of th	the patient within 10 minute time limit
------------------------	----------------------------	---

Failure to take or verbalize appropriate PPE precautions

Failure to determine scene safety

Failure to assess for and provide spinal protection when indicated

Failure to voice and ultimately provide high concentration oxygen

Failure to assess/provide adequate ventilation

Failure to find or appropriately manage problems associated with airway, breathing, hemorrhage or shock

Failure to differentiate patient's need for immediate transportation versus continued assessment/treatment at the scene

Performs other assessment before assessing/treating threats to airway, breathing and circulation

Failure to manage the patient as a competent EMT

Exhibits unacceptable affect with patient or other personnel

Uses or orders a dangerous or inappropriate intervention



Skill Sheet 3 TESTED

BLEEDING CONTROL/SHOCK MANAGEMENT

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Applies direct pressure to the wound		1	
NOTE: The examiner must now inform candidate that the would	nd continues to bleed.		
Applies tourniquet		1	
NOTE: The examiner must now inform candidate that the patie	ent is exhibiting signs and symptoms of hyp	operfusion.	
Properly positions the patient		1	
Administers high concentration oxygen		1	
Initiates steps to prevent heat loss from the patient		1	
Indicates the need for immediate transportation		1	
Actual Time Ended:	TOTAL	7	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to administer high concentration oxygen Failure to control hemorrhage using correct procedures in a ti Failure to indicate the need for immediate transportation Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	mely manner		



Skill Sheet 4
TESTED

AIRWAY MANAGEMENT

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Checks responsiveness		1	
Requests additional EMS assistance		1	
Checks breathing and pulse simultaneously		1	
NOTE: After checking responsiveness, then checking breathing		miner infori	ns
candidate, "The patient is unresponsive, apneic and has a weak	pulse of 60."		
Opens airway properly		1	
Ventilates the patient at a proper volume and rate via BVM		1	
Directs assistant to take over BVM ventilation and pre-oxygenate pat	ient	1	
Selects appropriate size Supraglottic Airway		1	
Inspects and prepares Supraglottic Airway for insertion		1	
Positions head properly		1	
Displace the tongue and jaw		1	
Advance the Supraglottic Airway to the proper depth recommended by	by the manufacturer	1	
Insert orogastric tube if required		1	
Secure tube in place		1	
Confirm placement via auscultation and secondary detection method		1	
Reassess patient		1	
Document procedure		1	
Actual Time Ended:	TOTAL	17	
CRITICAL CRITERIA			
Failure to take or verbalize appropriate PPE			
Failure to check responsiveness, then check breathing and pulse	e simultaneously for no more than 10 seconds		
Failure to ventilate the patient at the appropriate rate			
Failure to select proper size Supraglottic Airway			
Failure to secure tube			
Failure to confirm placement			
Failure to manage the patient as a competent EMT			
Exhibits unacceptable affect with patient or other personnel			
Uses or orders a dangerous or inappropriate intervention			



Skill Sheet 5
VERIFIED

CARDIAC ARREST MANAGEMENT / AED

Examiner		
Signature:		
	Possible Points	Points Awarded
	1	
	1	
	1	
	1	
	1	
	1	
and pulse for no more than 10 seconds ic and pulseless."	,	
llows the chest to recoil completely]	1	
point)	5	
d second rescuer resumes compression	ns while	
	1	
	1	
rhythm analysis	1	
from AED	1	
	1	
TOTAL	17	
selessness is confirmed CPR		
	and pulse for no more than 10 seconds ic and pulseless." lows the chest to recoil completely] point) d second rescuer resumes compression rhythm analysis from AED TOTAL e simultaneously for no more than 10 second selessness is confirmed CPR	Possible Points 1



Skill Sheet 6
VERIFIED

BASELINE VITAL SIGNS

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Blood Pressure (Palpation)			
Apply BP cuff approximately 1" above the antecubital space Not over clothing Snug fit Center bladder over artery		1	
Palpate radial and brachial artery		1	
Inflate cuff to a minimum of 20 mmHg above the point that the palp	able pulse was lost	1	
Slowly deflate the cuff		1	
Record/report the palpable systolic blood pressure when the pulse	returns (margin +/- 4mmHg)	1	
Blood Pressure (Auscultation)			
Apply BP cuff approximately 1" above the antecubital space Not over clothing Snug fit Center bladder over artery		1	
Palpate brachial artery		1	
Place diaphragm of stethoscope over brachial artery		1	
Inflate cuff to a minimum of 20 mmHg above the point that the palp	able pulse was lost	1	
Slowly deflate the cuff	·	1	
Record/report the palpable blood pressure (margin +/- 4mmHg)		1	
Pulse			
Palpate with two (2) fingers (index and middle) over the radial arter	У	1	
Count the palpated pulse for 30 seconds and multiply X 2		1	
Asses the following: Rate Rhythm (Regular/Irregular) Quality (Strong/Weak)		1 point each	
Record/Report pulse findings (margin +/- 4 bpm)		1	
Respirations			
Observe rise and fall of the chest or abdomen		1	
Count respirations for 30 seconds and X 2		1	
Asses the following: Rate Rhythm (Regular/Irregular) Quality (Strong/Weak)		1 point each	
Record/Report pulse findings (margin +/- 4 bpm)		1	
Skin			
Skin Color: Normal, Cyanotic, Jaundice, Ashen, Pale, Flushed		1	
Skin Temperature: Normal, warm, cool, hot		1	
Skin Condition: Normal, moist, diaphoretic		1	
Actual Time Ended:	TOTAL	27	

Failure to take or verbalize appropriate PPE precautions
Failure to manage the patient as a competent EMT
Exhibits unacceptable affect with patient or other personnel
Uses or orders a dangerous or inappropriate intervention
Van annual francolle de coment commente de la feu de africa annual the above anticol itema an the recomme aide of this forms

CRITICAL CRITERIA



Skill Sheet 7
VERIFIED

SPINAL IMMOBILIZATION (SEATED PATIENT)

Candidate:	Examiner:		
Date:			
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Directs assistant to place/maintain head in the neutral, in-line	position	1	
Directs assistant to maintain manual stabilization of the head	l	1	
Reassesses motor, sensory and circulatory functions in each	n extremity	1	
Applies appropriately sized extrication collar		1	
Positions the immobilization device behind the patient		1	
Secures the device to the patient's torso		1	
Evaluates torso fixation and adjusts as necessary		1	
Evaluates and pads behind the patient's head as necessary		1	
Secures the patient's head to the device	-	1	
Verbalizes moving the patient to a long backboard		1	
Reassesses motor, sensory and circulatory function in each	extremity	1	
Actual Time Ended:	TOTAL	12	
CRITICAL CRITERIA Failure to immediately direct or take manual stabilization Failure to properly apply appropriately sized cervical co Released or ordered release of manual stabilization be Manipulated or moved patient excessively causing pote Head immobilized to the device before device sufficient Device moves excessively up, down, left or right on the Head immobilization allows for excessive movement Torso fixation inhibits chest rise, resulting in respiratory Upon completion of immobilization, head is not in a new Failure to reassess motor, sensory and circulatory funct Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other perso	Illar before ordering release of manual stabilization fore it was maintained mechanically ential spinal compromise of the secured to the torso expatient's torso of compromise of compromise of the secured to the secured to the torso of compromise of the secured to the secured	to the long back	kboard
Uses or orders a dangerous or inappropriate intervention			



Skill Sheet 8 VERIFIED

SPINAL IMMOBILIZATION (SUPINE PATIENT)

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Directs assistant to place/maintain head in the neutral, in-line position	1	1	
Directs assistant to maintain manual stabilization of the head		1	
Reassesses motor, sensory and circulatory function in each extremity	/	1	
Applies appropriately sized extrication collar		1	
Positions the immobilization device appropriately		1	
Directs movement of the patient onto the device without compromising	g the integrity of the spine	1	
Applies padding to void between the torso and the device as necessar	ary	1	
Immobilizes the patient's torso to the device		1	
Evaluates and pads behind the patient's head as necessary		1	
Immobilizes the patient's head to the device		1	
Secures the patient's legs to the device		1	
Secures the patient's arms to the device		1	
Reassesses motor, sensory and circulatory function in each extremity	y	1	
Actual Time Ended:	TOTAL	. 14	
CRITICAL CRITERIA Failure to immediately direct or take manual stabilization of the last part of properly apply appropriately sized cervical collar before Released or ordered release of manual stabilization before it was Manipulated or moved the patient excessively causing potential Head immobilized to the device before device sufficiently secured Patient moves excessively up, down, left or right on the device Head immobilization allows for excessive movement Upon completion of immobilization, head is not in a neutral, in-life Failure to reassess motor, sensory and circulatory functions in experience to manage the patient as a competent EMT	re ordering release of manual stabilization as maintained mechanically spinal compromise red to the torso	the device	

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.

Uses or orders a dangerous or inappropriate intervention



LONG BONE IMMOBILIZATION

Skill Sheet 9 VERIFIED

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Directs application of manual stabilization of the injury		1	
Assesses distal motor, sensory and circulatory functions in the inju	red extremity	1	
NOTE: The examiner acknowledges, "Motor, sensory and circ	ulatory functions are present and normal	."	
Measures the splint		1	
Applies the splint		1	
Immobilizes the joint above the injury site		1	
Immobilizes the joint below the injury site		1	
Secures the entire injured extremity		1	
Immobilizes the hand/foot in the position of function		1	
Reassesses distal motor, sensory and circulatory functions in the i	njured extremity	1	
NOTE: The examiner acknowledges, "Motor, sensory and circ	ulatory functions are present and normal	."	
Actual Time Ended:	TOTAL	10	
Critical Criteria Failure to immediately stabilize the extremity manually Grossly moves the injured extremity Failure to immobilize the joint above and the joint below the in Failure to immobilize the hand or foot in a position of function Failure to reassess distal motor, sensory and circulatory function Failure to manage the patient as a competent EMT		splinting	

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.

Exhibits unacceptable affect with patient or other personnel
Uses or orders a dangerous or inappropriate intervention



JOINT IMMOBILIZATION

Skill Sheet 10 VERIFIED

Candidate:	_ Examiner:		
Date:	_ Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Directs application of manual stabilization of the injury		1	
Assesses distal motor, sensory and circulatory functions in the ir	njured extremity	1	
NOTE: The examiner acknowledges, "Motor, sensory and ci	rculatory functions are present and normal.	•	
Selects the proper splinting material		1	
Immobilizes the site of the injury		1	
Immobilizes the bone above the injury site		1	
Immobilizes the bone below the injury site		1	
Secures the entire injured extremity		1	
Reassesses distal motor, sensory and circulatory functions in the	e injured extremity	1	
NOTE: The examiner acknowledges, "Motor, sensory and ci	rculatory functions are present and normal.	13	
Actual Time Ended:	TOTAL	9	
Critical Criteria Failure to immediately stabilize the extremity manually Grossly moves the injured extremity Failure to immobilize the bone above and below the injurys Failure to reassess distal motor, sensory and circulatory fur Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personne Uses or orders a dangerous or inappropriate intervention	nctions in the injured extremity before and after	splinting	



12 LEAD EKG ACQUISITION

Skill Sheet 11	
VERIFIED	

Candidate:	Examiner:		
Date:	Signature:		111
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Identifies Indications for 12 Lead EKG acquisition		1	
Prepares monitor and connects electrodes to the patient cable		1	
Explains procedure to patient		1	
Exposes patient's chest and preps as necessary		1	
Properly applies chest leads (V1, V2, V3, V4, V5, V6, and limb leads V1: Right 4 th intercostal space beside sternum V2: Left 4 th intercostal space beside sternum V4: Left 5 th intercostal space, midclavicular V3: Halfway between V2 and V4 V5: Horizontal to V4, anterior to axillary line V6: Horizontal to V5, Mid-axillary line)	1 point each	
Properly applies Limb Leads (RA, LA, LA, LL)		1	
Instructs patient to remain as still as possible		1	
Acquires 12 lead EKG per manufacturer's instructions		1	
Transmits EKG to receiving facility		1	
Reassess patient		1	
Confirm transmission of 12 lead has completed		1	
Document the procedure		1	
Actual Time Ended:	TOTAL	. 18	
Critical Criteria Failure to take or verbalize appropriate PPE precautions Failure to identify the need for 12 lead EKG acquisition Failure to appropriately apply leads Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention			



CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP)

Skill Sheet 12 VERIFIED

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Performs initial assessment		1	
Applies initial high flow oxygen		1	
Identifies indications for CPAP utilization		1	
Identifies any contraindications for CPAP		1	
Explains the procedure to the patient		1	
Assembles CPAP correctly per manufacturer's directions		1	
Sets device parameters per protocol		1	
Applies device to patient obtaining a good face seal		1	
Adjusts pressure as required		1	
Comforts/coaches patient through the use of CPAP		1	
Reassess patient		1	
If no improvement contact medical command and request ALS back	up	1	
Document the procedure		1	
Actual Time Ended:	TOTAL	14	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to assemble device appropriately Failure to confirm a good face seal Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention			



Skill Sheet 13 VERIFIED

NALOXONE ADMINISTRATION

Candidate:	Examiner		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Identify the need for administration of Naloxone based on PPMH, vital,	signs and symptoms, and presentation	1	
Support respirations as needed		1	
Assess blood glucose level		1	
Verbalize signs of opioid use		1	
Select the proper medication and check concentration, color, and clarit	у	1	
Selects the appropriate syringe and draw up medication if not prefilled		1	
Confirm expiration date of medication		1	
Confirm the rights of drug administration: Right Patient Right Route Right Drug Right Dose	Right Time	1 point each	
Place a nebulizer on the end of the syringe	-	1	
Place atomizer against nostril and administer 1mg (0.5 dose) of medical		1	
Repeat the procedure delivering the remainder of the medication in the	opposite nostrii	1	
Reassess patient		1	
If no improvement, contact medical command and request ALS back u	p		
Document the procedure		1	
Actual Time Ended:	TOTAL	19	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to confirm at least three of the "Rights" of medication adm Failure to select appropriate medication or concentration of medication as needed Failure to support respirations as needed Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention			



Skill Sheet 14 VERIFIED

TETRACAINE OPHTHALMIC ADMINISTRATION / MORGAN LENS IRRIGATION

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Identifies indications for use of Morgan Lens		1	
Determines no contraindications for use of Morgan Lens		1	
Confirm patient allergies		1	
Confirm expiration date of medication		1	
Confirm the rights of drug administration: Right Patient ☐ Right Route ☐ Right Drug ☐ Right Dose [☐ Right Time ☐	1 point each	
Explains the procedure to patient		1	
Administers two (2) drops of tetracaine per eye being irrigated		1	
Attached macro-drop IV tubing to IV Bag		1	
Attach Morgan Lens delivery set to IV tubing and confirm fluid flowing	g through device	1	
With patient looking downward, retract upper eye lid and insert More	gan Lens under upper eye lid	1	
Have patient look upward, retract lower eye lid and place Morgan Le	ens	1	
Adjust flow to irrigate the eye		1	
Completes irrigations and removes Morgan Lens by retracting lower	eye lid and sliding the lens out	1	
Terminates IV Flow		1	
Reassess Patient		1	
Document the procedure		1	
Actual Time Ended:	TOTAL	21	
CRITICAL CRITERIA Failure to take or verbalize appropriate PPE precautions Failure to confirm expiration date of the medication Failure to confirm at least three of the "Rights" of medication adr Failure to provide continuous flow while irrigating patient's eye Failure to manage the patient as a competent EMT Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	ninistration		





OXYGEN ADMINISTRATION BY NON-REBREATHER MASK

Skill Sheet 15 VERIFIED

Candidate: Ex	caminer:		
Date: Si	gnature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Gathers appropriate equipment		1	
Cracks valve on the oxygen tank		1	
Assembles the regulator to the oxygen tank		1	
Opens the oxygen tank valve		1	
Checks oxygen tank pressure		1	
Checks for leaks		1	
Attaches non-rebreather mask to correct port of regulator		1	
Turns on oxygen flow to prefill reservoir bag		1	
Adjusts regulator to assure oxygen flow rate of at least 10 L/minute		1	
Attaches mask to patient's face and adjusts to fit snugly		1	
Actual Time Ended:	TOTAL	11	
CRITICAL CRITERIA			
Failure to take or verbalize appropriate PPE precautions			
Failure to assemble the oxygen tank and regulator without leaks			
Failure to prefill the reservoir bag			
Failure to adjust the oxygen flow rate to the non-rebreather mask of a	t lease 10 L/minute		
Failure to ensure a tight mask seal to patient's face			
Failure to manage the patient as a competent EMT			
Exhibits unacceptable affect with patient or other personnel			
Uses or orders a dangerous or inappropriate intervention			



Skill Sheet 16 VERIFIED

BVM VENTILATION OF AN APNEIC ADULT PATIENT

Date:	Candidate:	Examiner:		
Actual Time Started:		Signature:		
Takes or verbalizes appropriate PPE precautions Takes or verbalizes appropriate PPE precautions 1 Checks responsiveness 1 Checks responsiveness 1 Checks presponsiveness 1 Checks breathing and pulse simultaneously NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and has a weak pulse of 60." Opens airway properly 1 Checks breathing and pulse simultaneously NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction and oropharynx Turns on power to suction and or	<u> </u>	Olgridiano.		
Takes or verbalizes appropriate PPE precautions Takes or verbalizes appropriate PPE precautions 1 Checks responsiveness 1 Checks responsiveness 1 Checks presponsiveness 1 Checks breathing and pulse simultaneously NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and has a weak pulse of 60." Opens airway properly 1 Checks breathing and pulse simultaneously NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction and oropharynx Turns on power to suction and or				
Takes or verbalizes appropriate PPE precautions Takes or verbalizes appropriate PPE precautions 1 Checks responsiveness 1 Checks responsiveness 1 Checks presponsiveness 1 Checks breathing and pulse simultaneously NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and has a weak pulse of 60." Opens airway properly 1 Checks breathing and pulse simultaneously NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction device or retrieves manual suction device 1 Turns on power to suction and oropharynx 2 Turns on power to suction and oropharynx 3 Turns on power to suction and oropharynx 3 Turns on power to suction and oropharynx 3 Turns on power to suction and oropharynx 4 Turns on power to suction and oropharynx 4 Turns on power to suction and oropharynx 5 Turns on power to suction and transfer to suction and transfer to suction and transfer to suct			Possible	Points
Checks responsiveness Requests additional EMS assistance Requests additional EMS assistance 1 1 1 1 1 1 1 1 1 1	Actual Time Started:			
Checks responsiveness Requests additional EMS assistance Requests additional EMS assistance 1 1 1 1 1 1 1 1 1 1				
Requests additional EMS assistance Checks breathing and pulse simultaneously 1 Cypens alway properly 1 Cypens in a cypens of suction catheter 1 Cypens alway properly 1 Cypens to suction device or retrieves manual suction device 1 Cypens the alway part of the candidate, "The mouth is full of secretions and vomitus." Cypens the alway manually	Takes or verbalizes appropriate PPE precautions		1	
Checks breathing and pulse simultaneously NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is surresponsive, apneic and has a weak pulse of 60." Opens airway properly Opens airway properly NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter Turns on power to suction device or retrieves manual suction device 1 Inserts rigid suction catheter without applying suction 1 Inserts rigid suction catheter without applying suction NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually 1 Inserts oropharyngeal airway 2 Inserts the airway manually 3 Inserts oropharyngeal airway 4 Inserts oropharyngeal airway 1 Inserts oropharyngeal airway 2 Inserts oropharyngeal airway 3 Inserts oropharyngeal airway 4 Inserts oropharyngeal airway 4 Inserts oropharyngeal airway 5 Inserts oropharyngeal airway 5 Inserts oropharyngeal airway 6 Inserts oropharyngeal airway 1 Inserts oropharyngeal airway 2 Inserts oropharyngeal airway 3 Inserts oropharyneau clear." 1 Inserts oropharyneau clear." 1 Inserts oropharyneau clear." 1 Inserts oropharyneau clear." 1 Inserts oropha	Checks responsiveness		1	
NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and has a weak pulse of 60." NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares figid suction catheter	Requests additional EMS assistance		1	
Opens alrway properly Opens alrway properly NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter Turns on power to suction device or retrieves manual suction device Inserts rigid suction catheter without applying suction Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen ["*Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:	Checks breathing and pulse simultaneously		1	
Opens airway properly NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepreser sligid suction catheter Turns on power to suction device or retrieves manual suction device 1 Inserts rigid suction catheter without applying suction Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." **Ventilates the patient immediately using a BVM device unattached to oxygen [**The examiner must now inform the candidate that ventilation is being properly performed without difficulty. **Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: TOTAL 16 **CRITICAL CRITERIA* After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to buscion airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to to vice and ultimately provide high oxygen concentration [at least 65%] Failure to voice and ultimately provide high oxygen concentration [at least 65%] Failure to voice and ultimately provide high oxygen concentration [at least 65%] Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to provide adequate volumes per breath [maximum 2 errors/min	NOTE: After checking responsiveness, then checking breathing a	nd pulse for no more than 10 seconds, exa	miner infor	ms
NOTE: The examiner must now inform the candidate, "The mouth is full of secretions and vomitus." Prepares rigid suction catheter 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	candidate, "The patient is unresponsive, apneic and has a weak p	ulse of 60."		
Prepares rigid suction catheter Turns on power to suction device or retrieves manual suction device 1 1 Inserts rigid suction catheter without applying suction Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen [""Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as 1 first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately - Proper volume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper rolume to cause visible chest rise (1 point) - Proper volume to cause visible chest ris			1	
Turns on power to suction device or retrieves manual suction device Inserts rigid suction catheter without applying suction Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen ["Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 (Ventilates the patient adequately Proper volume to cause visible chest rise (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:	NOTE: The examiner must now inform the candidate, "The mouth	is full of secretions and vomitus."		
Inserts rigid suction catheter without applying suction Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen ["*Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rolume to cause visible chest rise (1 point) -Proper rolume to cause visible chest rise (1 point) -Proper rate [10 - 12/minute (1 ventilation every 5 - 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: TOTAL 16 CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to vortilate the patient at a rate of 10 - 12/minute (1 ventilation every 5 - 6 seconds) Failure to vortilate the patient at a rate of 10 - 12/minute (1 ventilation every 5 - 6 seconds) Failure to manage the patient as a competent EMT			1	
Suctions the mouth and oropharynx NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen ("Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as Infirst ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Indicates the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	Turns on power to suction device or retrieves manual suction device		1	
NOTE: The examiner must now inform the candidate, "The mouth and oropharynx are clear." Opens the airway manually Inserts oropharyngeal airway 1 Inserts oropharyngeal airway 1 NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." **Ventilates the patient immediately using a BVM device unattached to oxygen [**Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as 1 first ventilation is delivered within 30 seconds.] **NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds 1 Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately 1 -Proper volume to cause visible chest rise (1 point) 2 -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) **Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" **Actual Time Ended:	Inserts rigid suction catheter without applying suction		1	
Inserts oropharyngeal airway 1			1	
Inserts oropharyngeal airway NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." "Ventilates the patient immediately using a BVM device unattached to oxygen ["Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:		and oropharynx are clear."		
NOTE: The examiner must now inform the candidate, "No gag reflex is present and the patient accepts the airway adjunct." ***Ventilates the patient immediately using a BVM device unattached to oxygen [***Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] **NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. **Re-checks pulse for no more than 10 seconds 1 Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) 2 -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) **Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" **Actual Time Ended: TOTAL 16 **CRITICAL CRITERIA** After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT Failure to manage the patient as a competent EMT	Opens the airway manually		121	
Ventilates the patient immediately using a BVM device unattached to oxygen [Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] **NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. **Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) **Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" **Actual Time Ended:				
Award this point if candidate elects to ventilate initially with BVM attached to reservoir and oxygen so long as first ventilation is delivered within 30 seconds.] **NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. **Re-checks pulse for no more than 10 seconds			irway adjun	ct."
first ventilation is delivered within 30 seconds.] NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 4 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to manage the patient as a competent EMT		T. T.		
NOTE: The examiner must now inform the candidate that ventilation is being properly performed without difficulty. Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	IV5	ached to reservoir and oxygen so long as	1	
Re-checks pulse for no more than 10 seconds Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT				
Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/minute] 1 Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) 2 -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: TOTAL 16 CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT		on is being properly performed without dif	ficulty.	
Ventilates the patient adequately -Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:				
-Proper volume to cause visible chest rise (1 point) -Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:	Attaches the BVM assembly [mask, bag, reservoir] to oxygen [15 L/min	nute]	1	
-Proper rate [10 – 12/minute (1 ventilation every 5 – 6 seconds)] (1 point) Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended: TOTAL 16 CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	**************************************			
Note: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?" Actual Time Ended:			2	
Actual Time Ended: TOTAL 16 CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT				
Actual Time Ended: TOTAL 16 CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	Note: The examiner must now ask the candidate, "How would you	know if you are delivering appropriate vol	umes with	
CRITICAL CRITERIA After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	each ventilation?"			
After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	Actual Time Ended:	TOTAL	16	
After suctioning the patient, failure to initiate ventilations within 30 seconds or interrupts ventilations for greater than 30 seconds at any time Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	CRITICAL CRITERIA			
Failure to take or verbalize appropriate PPE precautions Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT		seconds or interrupts ventilations for greater that	an 30 second	s at any time
Failure to suction airway before ventilating the patient Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT		3		
Suctions the patient for an excessive and prolonged time Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	- C			
Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT				
Failure to voice and ultimately provide high oxygen concentration [at least 85%] Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT		simultaneously for no more than 10 seconds		
 Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT 				
Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible] Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilate	ation every 5 – 6 seconds)		
Insertion or use of any adjunct in a manner dangerous to the patient Failure to manage the patient as a competent EMT	Failure to provide adequate volumes per breath [maximum 2 erro	rs/minute permissible1		
Failure to manage the patient as a competent EMT	Insertion or use of any adjunct in a manner dangerous to the nation	ent		
The state of the s				
Uses or orders a dangerous or inappropriate intervention				