EXTRACTIONS

POST-OPERATION INSTRUCTIONS

When to call us:

It is normal to experience some discomfort for several days after a tooth extraction, but call us right away if you have:

- · Heavy or increased bleeding
- Pain or swelling that increases or continues beyond 2-3 days
- A bad taste or odor in your mouth
- A reaction to the medication

During the first 24 hours:

It is important that a blood clot forms on the extraction site to stop bleeding, reduce pain, and speed healing. To protect the clot and avoid the pain of a dry socket:

- Bite on a gauze pad firmly for 30-60 minutes. Blood and saliva mix in the mouth and make it look like there is more bleeding than there really is. Some oozing is normal; however after 1 hour, repeat with a clean gauze pad if oozing is profuse. The site could ooze for as long as 24 hours.
- Do not spit, and do not suck on candies or through a straw.
- Do not rinse your mouth, and do not brush or floss next to the site.
- Do not smoke or use tobacco. Avoid tobacco for at least 72 hours because it slows healing.
- Do not sneeze or cough, so have sinus or allergy medication on hand if necessary.
- Limit yourself to calm activities and elevate your head with pillows when you lie down to reduce bleeding.
- Do not drink hot, carbonated, or alcoholic drinks, and avoid hot or spicy foods.
- To control discomfort, take pain medication before the anesthetic has worn off or as recommended.
- To keep swelling to a minimum, use an ice bag over the area, 20 minutes on and 20 minutes off.
- When the numbness has worn off completely, drink lots of fluids and eat only soft, nutritious foods, chewing on the opposite side.

After the first 24 hours:

- Begin to eat normally as soon as it is comfortable.
- Resume brushing and flossing, but clean gently around the site for about a week.
- If antibiotics were prescribed, continue to take them for the indicated length of time, even if all symptoms and signs of infection are gone.
- Reduce soreness or swelling by applying moist heat. Swelling usually starts to go down after 48 hours.
- Further reduce swelling by rinsing your mouth very gently with warm salt water. Use about one teaspoon of salt per glass of warm water. Rinse 2-3 times a day for the week following the extraction.
- If you have any problems or questions, please do not hesitate to call our office (303) 988-6860.