

Please join us for a discussion on



On behalf of Daiichi Sankyo, Inc. and AstraZeneca, we cordially invite you to join us for an informative, expert-led discussion.

#### This program will cover:

- An FDA-approved therapy for adult patients with HER2+ unresectable or metastatic breast cancer after two or more prior anti-HER2-based regimens in the metastatic setting\*
- Efficacy and safety results from the Phase 2 clinical trial, DESTINY-Breast01
- Preparation, dosage, and administration of ENHERTU
- Counseling considerations for patients who are being treated with ENHERTU

#### Program Details:

**09/11/2021**

7:00 PM Pacific  
The Palm - L.A. Downtown  
1100 South Flower Street  
Los Angeles, CA 90015

#### Faculty:

**Linda Buck, MSN, AOCNP**  
City of Hope  
Mission Viejo, CA

\*This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

#### RSVP: Noemi Sklar

Representative Program Host

[nsklar@dsi.com](mailto:nsklar@dsi.com)

(917) 526-2202

By: 09/04/2021

This program is being sponsored by Daiichi Sankyo, Inc. and AstraZeneca. The speaker is being compensated for the presentation. The program is not CME accredited and may not be used for CME accreditation. In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs. Please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you may receive may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity that limits or prohibits meals from pharmaceutical companies, please identify yourself so that compliance with such requirements can be ensured.

### Important Safety Information

#### Indication

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received two or more prior anti-HER2-based regimens in the metastatic setting.

This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

#### WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY

- **Interstitial lung disease (ILD) and pneumonitis, including fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**
- **Exposure to ENHERTU during pregnancy can cause embryo-fetal harm. Advise patients of these risks and the need for effective contraception.**

#### Contraindications

None.

#### Warnings and Precautions

##### Interstitial Lung Disease / Pneumonitis

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU. In clinical studies, of the 234 patients with unresectable or metastatic HER2-positive breast cancer treated with ENHERTU 5.4 mg/kg, ILD occurred in 9% of patients. Fatal outcomes due to ILD and/or pneumonitis occurred in 2.6% of patients treated with ENHERTU. Median time to first onset was 4.1 months (range: 1.2 to 8.3).

Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in  $\leq 28$  days from date of onset, maintain dose. If resolved in  $> 28$  days from date of onset, reduce dose one level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 0.5$  mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 1$  mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

## Important Safety Information (continued)

### Neutropenia

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU. In clinical studies, of the 234 patients with unresectable or metastatic HER2-positive breast cancer who received ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 62% of patients. Sixteen percent had Grade 3 or 4 decrease in neutrophil count. Median time to first onset of decreased neutrophil count was 23 days (range: 6 to 547). Febrile neutropenia was reported in 1.7% of patients.

Monitor complete blood counts prior to initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC]  $<1.0$  to  $0.5 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC  $<0.5 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 2 or less. Reduce dose by one level. For febrile neutropenia (ANC  $<1.0 \times 10^9/L$  and temperature  $>38.3^\circ C$  or a sustained temperature of  $\geq 38^\circ C$  for more than 1 hour), interrupt ENHERTU until resolved. Reduce dose by one level.

### Left Ventricular Dysfunction

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular ejection fraction (LVEF) decrease has been observed with anti-HER2 therapies, including ENHERTU. In the 234 patients with unresectable or metastatic HER2-positive breast cancer who received ENHERTU, two cases (0.9%) of asymptomatic LVEF decrease were reported. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF  $<50\%$  prior to initiation of treatment.

Assess LVEF prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. When LVEF is  $>45\%$  and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is  $<10\%$ , continue treatment with ENHERTU and repeat LVEF assessment within 3 weeks. When LVEF is 40-45% and absolute decrease from baseline is 10-20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF has not recovered to within 10% from baseline, permanently discontinue ENHERTU. If LVEF recovers to within 10% from baseline, resume treatment with ENHERTU at the same dose. When LVEF is  $<40\%$  or absolute decrease from baseline is  $>20\%$ , interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of  $<40\%$  or absolute decrease from baseline of  $>20\%$  is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure.

### Embryo-Fetal Toxicity

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for at least 7 months following the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 4 months after the last dose of ENHERTU.

### Additional Dose Modifications

#### Thrombocytopenia

For Grade 3 thrombocytopenia (platelets  $<50$  to  $25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets  $<25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less. Reduce dose by one level.

#### Adverse Reactions

The safety of ENHERTU was evaluated in a pooled analysis of 234 patients with unresectable or metastatic HER2-positive breast cancer who received at least one dose of ENHERTU 5.4 mg/kg in DESTINY-Breast01 and Study DS8201-A-J101. ENHERTU was administered by intravenous infusion once every three weeks. The median duration of treatment was 7 months (range: 0.7 to 31).

Serious adverse reactions occurred in 20% of patients receiving ENHERTU. Serious adverse reactions in  $>1\%$  of patients who received ENHERTU were interstitial lung disease, pneumonia, vomiting, nausea, cellulitis, hypokalemia, and intestinal obstruction. Fatalities due to adverse reactions occurred in 4.3% of patients including interstitial lung disease (2.6%), and the following events occurred in one patient each (0.4%): acute hepatic failure/acute kidney injury, general physical health deterioration, pneumonia, and hemorrhagic shock.

ENHERTU was permanently discontinued in 9% of patients, of which ILD accounted for 6%. Dose interruptions due to adverse reactions occurred in 33% of patients treated with ENHERTU. The most frequent adverse reactions ( $>2\%$ ) associated with dose interruption were neutropenia, anemia, thrombocytopenia, leukopenia, upper respiratory tract infection, fatigue, nausea, and ILD. Dose reductions occurred in 18% of patients treated with ENHERTU. The most frequent adverse reactions ( $>2\%$ ) associated with dose reduction were fatigue, nausea, and neutropenia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (79%), white blood cell count decreased (70%), hemoglobin decreased (70%), neutrophil count decreased (62%), fatigue (59%), vomiting (47%), alopecia (46%), aspartate aminotransferase increased (41%), alanine aminotransferase increased (38%), platelet count decreased (37%), constipation (35%), decreased appetite (32%), anemia (31%), diarrhea (29%), hypokalemia (26%), and cough (20%).

### Use in Specific Populations

- **Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months following the last dose of ENHERTU.
- **Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- **Females and Males of Reproductive Potential:** Pregnancy testing: Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. Contraception: **Females:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 7 months following the last dose. **Males:** Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 4 months following the last dose. Infertility: ENHERTU may impair male reproductive function and fertility.
- **Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- **Geriatric Use:** Of the 234 patients with HER2-positive breast cancer treated with ENHERTU 5.4 mg/kg, 26% were  $\geq 65$  years and 5% were  $\geq 75$  years. No overall differences in efficacy were observed between patients  $\geq 65$  years of age compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged  $\geq 65$  years (53%) as compared to younger patients (42%).
- **Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, closely monitor for increased toxicities related to the topoisomerase inhibitor.

To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or [fda.gov/medwatch](http://fda.gov/medwatch).

Please [click here](#) for full Prescribing Information, including Boxed WARNINGS, and [click here](#) for Medication Guide.

