

Solomon Press
Core Mathematics C1
Paper B
(Mark Scheme)

All exam papers are issued free to students for education purpose only.
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com

GCE Examinations
Advanced Subsidiary

Core Mathematics C1

Paper B

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.




Written by Shaun Armstrong

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

C1 Paper B – Marking Guide

- | | |
|--|---|
| <p>1. $f(x) = x + 6\sqrt{x} + 9 + 1 - 6\sqrt{x} + 9x$
 $= 10x + 10, \quad a = 10, b = 10$</p> | <p>M1 A1
 A1 (3)</p> |
| <p>2. quadratic, coeff of $x^2 = 1$, minimum $(-2, 5)$
 $\therefore y = (x + 2)^2 + 5$
 $= x^2 + 4x + 9, \quad a = 4, b = 9$</p> | <p>M1 A1
 M1 A1 (4)</p> |
| <p>3. (a) $u_1 = 2 + k$
 $u_3 = 8 + 3k$
 $u_1 = u_3 \therefore 2 + k = 8 + 3k$
 $k = -3$</p> <p>(b) $u_5 = 2^5 - 3(5) = 32 - 15 = 17$</p> | <p>B1
 M1
 A1
 M1 A1 (5)</p> |
| <p>4. $y = \int (2x^3 + 1) dx$
 $y = \frac{1}{2}x^4 + x + c$
 $x = 0, y = 3 \therefore c = 3$
 $y = \frac{1}{2}x^4 + x + 3$
 when $x = 2, y = 8 + 2 + 3 = 13$</p> | <p>M1 A2
 B1
 M1 A1 (6)</p> |
| <p>5. (a) $= x(4 - 3x - x^2)$
 $= x(1 - x)(4 + x)$</p> <p>(b) </p> | <p>M1
 M1 A1

 B3

 (6)</p> |
| <p>6. $x = 0 \Rightarrow y = -6 \quad \therefore (0, -6)$
 $y = 0 \Rightarrow x = 12 \quad \therefore (12, 0)$
 mid-point $= (\frac{0+12}{2}, \frac{-6+0}{2}) = (6, -3)$
 dist. from $O = \sqrt{6^2 + (-3)^2} = \sqrt{36+9} = \sqrt{45}$
 $= \sqrt{9 \times 5} = 3\sqrt{5}$</p> | <p>B1
 M1 A1
 M1
 M1 A1 (6)</p> |
| <p>7. (a) (i) $2^{x+2} = 2^2 \times 2^x = 4y$
 (ii) $2^{3-x} = \frac{2^3}{2^x} = \frac{8}{y}$</p> <p>(b) $2^{x+2} + 2^{3-x} = 33 \Rightarrow 4y + \frac{8}{y} = 33$
 $4y^2 + 8 = 33y$
 $4y^2 - 33y + 8 = 0$</p> <p>(c) $(4y - 1)(y - 8) = 0$
 $y = \frac{1}{4}, 8$
 $2^x = \frac{1}{4}, 8$
 $x = -2, 3$</p> | <p>M1 A1
 M1 A1

 M1
 A1
 M1
 A1

 A2 (10)</p> |

8.	(a)	$\frac{dy}{dx} = 3x^{\frac{1}{2}}$	M1 A1
		$\frac{d^2y}{dx^2} = \frac{3}{2}x^{-\frac{1}{2}}$	A1
	(b)	LHS = $4x^2(\frac{3}{2}x^{-\frac{1}{2}}) - 3(2x^{\frac{3}{2}} - 1)$ $= 6x^{\frac{3}{2}} - 6x^{\frac{3}{2}} + 3$ $= 3$ [$k = 3$]	M1 A1
	(c)	$= \int (2x^{\frac{3}{2}} - 1)^2 dx$ $= \int (4x^3 - 4x^{\frac{3}{2}} + 1) dx$ $= x^4 - \frac{8}{5}x^{\frac{5}{2}} + x + c$	M1 A1 M1 A3 (11)

9.	(a)	$a + d = 26$ $a + 4d = 41$ subtracting, $3d = 15$ $d = 5$	M1 A1 M1 A1
	(b)	$a = 21$ $u_{12} = 21 + (11 \times 5) = 76$	B1 M1 A1
	(c)	$\frac{n}{2} [42 + 5(n - 1)] = \frac{n}{2} [-24 + 7(n - 1)]$ $n(5n + 37) = n(7n - 31)$ $2n(n - 34) = 0$ $n > 0 \therefore n = 34$	M1 A1 M1 A1 (11)

10.	(a)	$x^2 - 3x + 5 = 2x + 1$ $x^2 - 5x + 4 = 0$ $(x - 1)(x - 4) = 0$ $x = 1, 4$ when $x = 1, y = 2(1) + 1 = 3$ $\therefore P(1, 3), Q(4, 9)$	M1 M1 A1 A1
	(b)	$\frac{dy}{dx} = 2x - 3$ grad = -1 $\therefore y - 3 = -(x - 1)$ [$y = 4 - x$]	M1 A1 M1 A1
	(c)	grad = 5 $\therefore y - 9 = 5(x - 4)$ $y - 9 = 5x - 20$ $y = 5x - 11$	M1 A1
	(d)	$4 - x = 5x - 11$ $x = \frac{5}{2}$ $\therefore (\frac{5}{2}, \frac{3}{2})$	M1 A1 A1 (13)

Total (75)