


TRIBUTE®

KALM 'N EZ® TEXTURED

A textured low-starch feed formulated for all classes of mature horses.

PRODUCT CODE: T920EZ

FEATURES AND BENEFITS:

- ✦ **No Corn! No Molasses!** Keeps sugar and starch levels low.
- ✦ **Whole oats** – the lowest starch grain.
- ✦ **Higher fat, highly digestible fiber and lower sugar and starch** for calm performances without loss of energy or condition.
- ✦ Optimal balance of **Essential** amino acids for muscle maintenance.
- ✦ **Essential organic minerals** support normal bone and tissue maintenance.
- ✦ **Essential fatty acids, including Omega 3 and 6**, required for the support of normal tissue function and healthy skin and haircoat.
- ✦ Optimum levels of antioxidants Vitamin E, Vitamin C and Selenium reduce damage from aging and exercise. Also boosts immunity.
- ✦ Excellent for hard-keepers and horses requiring less sugar and starch in their diets, like those who are hyperactive or have metabolic conditions such as insulin resistance, laminitis, ulcers, obesity or tying-up issues.
- ✦ With  microencapsulated active dry yeast; acts as both a pre-and probiotic, increases total diet digestibility and improves overall gut health.

FEEDING DIRECTIONS:

Training Level	T920EZ (Lbs. per day)	Lbs. Hay per day
Mature Weight: 660-880 lbs		
Idle	3-5	6 - 10
Light Training	5-7	8 - 12
Moderate Training	6-8	8 - 12
Intense Training	8-10	9 - 14
Mature Weight: 880-1,100 lbs		
Idle	4-8	9 - 14
Light Training	6-10	10 - 16
Moderate Training	8-12	10 - 16
Intense Training	10-14	12 - 18
Mature Weight: 1,100-1,320 lbs		
Idle	5-9	12 - 18
Light Training	7-11	14 - 20
Moderate Training	10-14	14 - 20
Intense Training	12-16	16 - 22

- Kalm 'N EZ® Textured can be used as a complete feed when hay is in short supply or unavailable.

GUARANTEED ANALYSIS:

Crude Protein (Min.)	14%
Crude Fat (Min.)	6.0%
Crude Fiber (Max.)	20.0%
Calcium (Ca) (Min.)	0.8%
Calcium (Ca) (Max.)	1.3%
Phosphorus (P) (Min.)	0.6%
Lysine (Min.)	0.9%
Methionine (Min.)	0.3%
Methionine + Cysteine (Min.)	0.5%
Threonine (Min.)	0.6%
Copper (Cu) (Min.)	65 ppm
Zinc (Zn) (Min.)	200 ppm
Selenium (Se) (Min.)	0.6 ppm
Potassium (K) (Min.)	1.0%
Magnesium (Mg) (Min.)	0.35%
Manganese (Mn) (Min.)	160 ppm
Cobalt (Co) (Min.)	1 ppm
Iron (Fe) (Min.)	390 ppm
Iodine (I) (Min.)	0.8 ppm
Salt (NaCl) (Min.)	0.5%
Salt (NaCl) (Max.)	1.0%
Vitamin A (Min.)	5,500 IU/lb.
Vitamin D (Min.)	500 IU/lb.
Vitamin E (Min.)	200 IU/lb.
Vitamin C (Min.)	25 mg/lb.
Biotin (Min.)	0.65 mg/lb.
Thiamine (Min.)	8 mg/lb.
Riboflavin (Min.)	4.2 mg/lb.
Omega 6 fatty acids (Min.)	2.8%
Omega 3 fatty acids (Min.)	0.3%
Saccharomyces cerevisiae (Min.)	350 million CFU/lb.
Direct-Fed Microorganisms (Min.)	520 million CFU/lb.
(Saccharomyces cerevisiae, Lactobacillus acidophilus, Bacillus subtilis, Bacillus licheniformis, Bacillus coagulans, Enterococcus faecium, Bifidobacterium thermophilum, and Bifidobacterium longum)	
Cellulase (Aspergillus Oryzae) (Min.)	9.6 Enzyme Units
Protease (Aspergillus Oryzae) (Min.)	12 Enzyme Units
Lipase (Aspergillus Oryzae) (Min.)	3.6 Enzyme Units
Hemicellulase (Aspergillus Niger) (Min.)	10.8 Enzyme Units
Phytase (Trichoderma reesei) (Min.)	55 FTU/lb.

*An Enzyme Unit is defined as milligrams of substrate liberated/minute/lb. of feed.

*A Phytase Unit (FTU) is defined as the quantity of enzyme which liberates one micromole of inorganic phosphate per minute from sodium phytate at 37°C, 5.5 pH.

This feed contains a dry source of cellulase that breaks down cellulose, a dry source of protease that hydrolyzes proteins and increases the digestibility of protein in soybean meal based diets, a dry source of lipase that hydrolyzes triglycerides, a dry source of hemicellulase that breaks down hemicellulose, and a dry source of phytase which hydrolyzes phytate and increases the digestibility of phytin-bound phosphorus.