

# MUSTELID & VIVERRID CONSERVATION




The Newsletter of the IUCN/SSC  
Mustelid & Viverrid Specialist Group




Number 4

April 1991



Owston's palm civet (*Chrotogale owstoni*).

This rare palm civet was photographed for the first time in Vietnam. Photo by H. J. Adler.



This number was produced with the support of  
the "Royal Zoological Society of Antwerp", Antwerp, Belgium,  
and the "Metropolitan Toronto Zoo", West Hill, Ontario, Canada.



# **Mustelid & Viverrid Conservation**

The Newsletter of the IUCN/SSC Mustelid & Viverrid Specialist Group

**Editor-in-chief:** Harry Van Rompaey, Edegem, Belgium

**Editors:** Michael Riffel, Karlsdorf, Germany  
Arnd Schreiber, Heidelberg, Germany  
Roland Wirth, München, Germany

The views expressed in this Newsletter are those of the authors and do not necessarily reflect those of the IUCN, nor the IUCN/SSC Mustelid & Viverrid Specialist Group.

We are particularly grateful to Walter Rasmussen for reading the manuscripts and improving the English style.

The aim of this newsletter is to offer the members of the IUCN/SSC M&VSG, and those who are concerned with mustelids or/and viverrids, brief papers, news items, abstracts, and titles of recent literature. All readers are invited to send material to:

**Mustelid & Viverrid Conservation**  
**c/o Dr. H. Van Rompaey**  
**Jan Verbertlei, 15**  
**2650 Edegem**  
**Belgium**

# Aspects of the biology of the Small grey mongoose *Galerella pulverulenta*

C.T. STUART

## Abstract

The small grey mongoose *Galerella pulverulenta* is a common, diurnal predator throughout much of its limited range. In the present study 34 individuals were marked and released within a 200 ha area, with the home ranges of four animals being between 5 ha and 36 ha in extent. The estimated density was 1 mongoose/12.5 ha and there was considerable range overlap. Ninety two percent of the 163 sightings were of solitary animals. Food consists mainly of small murids and insects. Comparative notes are presented on the Yellow mongoose *Cynictis penicillata*.

## Introduction

The Small grey mongoose *Galerella pulverulenta* (Fig. 1) is a small, generally solitary, diurnal viverrid, restricted in its distribution to the Cape Province south of the Orange River, Orange Free State, and marginally in Natal, Lesotho and Namibia (Lynch, 1981; Stuart, 1981).

Despite its abundance, particularly in the south-western part of its range, little attention has been given to this mongoose by research workers. Lynch (1981) has discussed the taxonomic status of this species and limited aspects of its biology have been briefly examined by Crawford *et al.* (1983) and Stuart (1981). This paper reports the results of a home range, behaviour and diet study of the small grey mongoose on the Vrolijkheid Nature Conservation Station.



Fig. 1. Small grey mongoose (*Galerella pulverulenta*, also known as *Herpestes pulverulentus*)

## Study area

The study area was situated in the Vrolijkheid Nature Conservation Station in the extreme eastern portion of the Robertson Karoo, part of the Little Karoo system. Of the 200 ha study area some 40 ha consisted of cultivated and unused farmland. The southern boundary of the area was formed by the Keisers' River, behind which rises a steep and rocky hillside. A tarred road divides the study area and extensive vineyards and orchards bound the southern sector. The study area consisted of low karroid scrub and succulent vegetation on broken, rocky outcrops. The eastern and northern boundaries abutted on to steep, rocky hillslopes. The southern area consisted of open, eroded and overgrazed scrubland. Five small earth-walled dams were located in the western sector of the study area and one in the east. Only two of the dams held water throughout the year. The faunal and floral components of the area have been covered in some detail by Stuart (1974).

## Methods

Between April 1978 and July 1980 a live-trapping programme was undertaken to determine the small carnivore population of the study area, with special emphasis on the small grey mongoose. Trapping was undertaken at regular intervals, with a total of 7260 trap days (20 traps for 363 days), with single-door live-traps (75x36x36 cm). Eight traplines were marked out and each line consisted of 20 equally spaced markers (50 m between markers). Trapping was undertaken in each line for three days every 65 days. Six traplines ran from east to west and two from north to south (Fig. 2). Traps were baited with a blood-based scent lure and raw meat. In order to reduce disturbance traps were fixed in the open position between trapping periods.

All animals trapped were sexed, aged (juvenile, sub-adult or adult), measured, weighed, marked and released. Each mongoose was marked with numbered aluminium poultry wing-tags placed close to the base of each ear. Direct observations of free-ranging animals were made on an irregular basis. Scats of the small grey mongoose were collected and prey fragments identified in the laboratory as described by Stuart (1981).

## Results

### Home range and movement

A total of 34 individual small grey mongooses were trapped (Fig. 2), marked and released, of which 18 were male and 16 female. Thirteen (38%) of the animals were caught in the first two months of trapping and a total of 20 (59%) had been taken by the end of the seventh month. Only 16 animals were recaptured after marking, with one of the males being trapped 22 times and a female 12 times. Sufficient recaptures for estimating home range were only available for two males and two females (Fig. 3).

The established home range sizes were calculated using the "minimum home range" method (Mohr, 1947). These ranged from 5 ha (female B) to 36 ha (male C). From the limited information obtained it would seem that there is considerable overlap in home ranges within the study area.




Fig. 2. Spatial distribution of first captures of Small grey mongoose within the study area. Triangles: males; circles: females. The nine traplines and topographical features are also indicated. Dams and water courses are indicated by stippled overlay.

The longest linear movement recorded between captures was by male C, of 1800 m. Most other recorded movements between captures were, however, considerably less than this, with a mean of 350 m.

#### Social organisation and behaviour

A total of 163 sightings were made during this study, of which 151 were of solitary animals, seven were of pairs and five were of groups of three individuals. Three of the pairs and four of the groups of three consisted of an adult accompanied by one, or two sub-adults.

Food searching appears to be of an opportunistic nature and this is borne out by the diet analysis. Both sight and smell appear to play a significant role in food searching (Pers. obs.). Most attention is concentrated on searching the area at the bases of bushes and rocks.

Only on one occasion was an aggressive encounter ob-

served. Two adult mongooses were observed walking towards each other on the same path. Both animals (A & B) were involved in food searching, each unaware of the presence of the other. Animal A stopped, defecated and was about to continue when B came within sight of A. Animal A raised the hair on its tail, holding the tail low to the ground and ran at B, with B turning and A chasing it for approx. 10 m.

#### Density

With 34 individual mongooses having been marked within the study area, the density was approximately one mongoose per 6 ha. Eighteen animals were trapped once only and may have been non-resident within the study area, or they learned to avoid the traps. If these animals are excluded from the calculations the resident population would have been 16 animals and therefore a possible density of one mongoose per 12.5 ha. If one takes into account that 40 ha of the area consisted of marginally utilised habitat, the actual density would have been greater.

#### Food

Small grey mongoose scats were deposited singly, or in small, scattered accumulations along pathways and on dam walls. The content of 316 scats is summarised in Table 1. Rodent remains occurred in all ten monthly samples, ranging from 65% to 100% of the content. Of the 43 individual rodents identified to species level only two, *Rhabdomys pumilio* and *Otomys unisulcatus* occurred in significant numbers. The remaining four rodent species identified were much more limited in their distribution within the study area.

#### Discussion

There are few data with which to compare the findings of this study, with only one previous estimate of home range size for this mongoose having been made (Crawford *et al.*, 1983). This was based on casual observations of a single animal believed to range over 75 ha.

The only other comparable solitary and diurnal viverrid that has been studied, the slender mongoose, *Galerella sanguinea*, which had estimated home range sizes of 100 ha in the Serengeti National Park in Tanzania (Rood & Waser, 1978). Kingdon (1977) is, however, of the opinion that in some areas home ranges of the slender mongoose are considerably smaller.

Gorman (1979) working on the closely related Small Indian mongoose *Herpestes auropunctatus* in Fiji found that males had a mean home range of approximately 39 ha and females 22 ha. In a similar study in Puerto Rico Pimentel (1955) found that the same mongoose had home ranges of 0.5 ha and 1.2 ha resp., whereas Tomich (1969) found that the small Indian mongoose in Hawaii had home ranges of between 50 ha and 200 ha. These studies serve to demonstrate the variability of home range size within the same species in different areas.

In the current study the small grey mongoose home ranges were estimated from between 5 ha and 36 ha and with a density probably greater than one mongoose per 12.5 ha.

Solitary viverrids have two advantages: exploitation of small rodents as a food source requires hunting by stealth and is best accomplished when the individual hunts alone.

Approximately 70%, by mass, of the prey taken by small grey mongoose in the study area were vertebrates. In addition, the individual gains an intimate knowledge of its foraging range and




Fig. 3. Estimated home ranges of two male (solid and closed squares) and two females (solid and open circles) Small grey mongoose within the study area.

Table 1. Percentage occurrence of different food categories from Small grey mongoose scats through the months of the year.

	J	F	M	A	M	J	J	A	S	O	N	D
No. of scats	22	10	96	20	12	-	4	-	48	34	46	24
Rodents	100	100	73	70	83	-	(4)	-	92	88	65	92
Amphibians	-	-	4	10	-	-	-	-	-	-	-	-
Birds	9	-	6	20	-	-	-	-	6	12	26	8
Reptiles	9	-	2	10	-	-	-	-	-	6	17	8
Insects	73	80	94	100	67	-	(4)	-	100	94	22	67
Arachnids	-	-	2	10	-	-	(2)	-	-	6	13	-
Myriapods	18	20	-	40	-	-	(2)	-	-	6	13	-
Molluscs	18	-	-	40	-	-	-	-	-	-	9	-
Plants	45	20	17	10	-	-	-	-	-	6	4	42

Note: No percentages were calculated for the small July sample.

this is particularly important in a species which exploits invertebrates as a food source. Although invertebrates are abundant they generally have an aggregated and local distribution (Gorman, 1979). Insects were by far the most important invertebrate group represented in the present study, followed by myriapods and arachnids.

Small grey mongooses were rarely observed in the cultivated areas, or fallow land, but these were heavily utilised by the diurnal Yellow mongoose, *Cynictis penicillata*. Of the eight species of carnivores occurring within the study area only the small grey mongoose and the yellow mongoose are diurnal, solitary foragers. Two colonies of yellow mongoose, with an estimated total of 12 individuals, were located within the study area. There were no observed interactions between the two mongoose species during the present study. The yellow mongoose is an open area feeder but usually forages in fairly close proximity to its burrow systems, whereas the small grey mongoose shows a marked preference for feeding amongst, or close to, bush and scrub cover. In the present study it was found that the diet of the small grey mongoose was dominated by small murids and insects but that the diet of the yellow mongoose consisted mainly of insects and other invertebrates (Table 2). It is suggested, therefore, that the small grey mongoose and the yellow mongoose avoid competition by utilising largely different habitats and feeding niches.

Table 2. Yellow mongoose scat contents from Vrolijkheid Nature Conservation station. A=abundant; P=present.

	Jan.	Feb.	Apr.	May	Jul.	Sep.	Nov.
Sample size	38	42	15	22	25	19	2
Coleoptera	A	P	A	A	A	A	A
Orthoptera	A	A	A	A	A	-	P
Isoptera	P	-	-	P	P	-	P
Myriapoda	-	-	A	A	P	P	-
Scorpiones	P	P	-	-	P	-	-
Aves	-	-	P	-	-	P	-
Rodentia	-	-	-	-	-	P	-

Solitary viverrids have two advantages: exploitation of small rodents as a food source requires hunting by stealth and is best accomplished when the individual hunts alone.

Approximately 70%, by mass, of the prey taken by small grey mongoose in the study area were vertebrates. In addition, the individual gains an intimate knowledge of its foraging range and

Table 1. Percentage occurrence of different food categories from Small grey mongoose scats through the months of the year.

	J	F	M	A	M	J	J	A	S	O	N	D
No. of scats	22	10	96	20	12	-	4	-	48	34	46	24
Rodents	100	100	73	70	83	-	(4)	-	92	88	65	92
Amphibians	-	-	4	10	-	-	-	-	-	-	-	-
Birds	9	-	6	20	-	-	-	-	6	12	26	8
Reptiles	9	-	2	10	-	-	-	-	-	6	17	8
Insects	73	80	94	100	67	-	(4)	-	100	94	22	67
Arachnids	-	-	2	10	-	-	(2)	-	-	6	13	-
Myriapods	18	20	-	40	-	-	(2)	-	-	6	13	-
Molluscs	18	-	-	40	-	-	-	-	-	-	9	-
Plants	45	20	17	10	-	-	-	-	-	6	4	42

Note: No percentages were calculated for the small July sample.

this is particularly important in a species which exploits invertebrates as a food source. Although invertebrates are abundant they generally have an aggregated and local distribution (Gorman, 1979). Insects were by far the most important invertebrate group represented in the present study, followed by myriapods and arachnids.

Small grey mongooses were rarely observed in the cultivated areas, or fallow land, but these were heavily utilised by the diurnal Yellow mongoose, *Cynictis penicillata*. Of the eight species of carnivores occurring within the study area only the small grey mongoose and the yellow mongoose are diurnal, solitary foragers. Two colonies of yellow mongoose, with an estimated total of 12 individuals, were located within the study area. There were no observed interactions between the two mongoose species during the present study. The yellow mongoose is an open area feeder but usually forages in fairly close proximity to its burrow systems, whereas the small grey mongoose shows a marked preference for feeding amongst, or close to, bush and scrub cover. In the present study it was found that the diet of the small grey mongoose was dominated by small murids and insects but that the diet of the yellow mongoose consisted mainly of insects and other invertebrates (Table 2). It is suggested, therefore, that the small grey mongoose and the yellow mongoose avoid competition by utilising largely different habitats and feeding niches.

Table 2. Yellow mongoose scat contents from Vrolijkheid Nature Conservation station. A=abundant; P=present.

	Jan.	Feb.	Apr.	May	Jul.	Sep.	Nov.
Sample size	38	42	15	22	25	19	2
Coleoptera	A	P	A	A	A	A	A
Orthoptera	A	A	A	A	A	-	P
Isoptera	P	-	-	P	P	-	P
Myriapoda	-	-	A	A	P	P	-
Scorpiones	P	P	-	-	P	-	-
Aves	-	-	P	-	-	P	-
Rodentia	-	-	-	-	-	P	-




Fig. 3. Estimated home ranges of two male (solid and closed squares) and two females (solid and open circles) Small grey mongoose within the study area.

## Acknowledgements

Financial and logistical support was provided by the Directorate of Nature and Environmental Conservation, Cape Province, South Africa. B. Munnik, J. Adendorf, and F. Williams provided technical assistance and P. Norton commented on an early draft of this paper. Tilde Stuart is thanked for drawing the figures.

## Footnote

There are indications that *Galerella pulverulenta* is extending its distribution range, particularly in the eastern sector of South Africa. Bronner (1990) has recorded the capture of a female near Wakkerstroom, Transvaal, some 200 km north-east of records in Natal and Lesotho. This area was extensively surveyed by Roberts (1951) but no trace of this mongoose was found at the time. Local farmers were not familiar with *Galerella*. There are also unconfirmed sightings of this species in areas north of the Orange River (Cape Province) where they would overlap with *Galerella sanguinea*, another small diurnal predator with similar dietary patterns. We will, hopefully, be able to investigate these apparent range extensions as part of the African Carnivore Survey.

## References

- Crawford, P. B., Crawford, S. A. H. & Crawford, R. J. M. 1983. Some observations on Cape grey mongoose *Herpestes pulverulentus* in the Tsitsikamma National Park. *S. Afr. J. Wildl. Res.*, 13:35-40.
- Gorman, M. L. 1979. Dispersion and foraging of the small Indian mongoose, *Herpestes auro-punctatus* (Carnivora: Viverridae) relative to the evolution of social viverrids. *J. Zool.*, 187:65-73.
- Kingdon, J. 1977. *East African mammals: an atlas of evolution in Africa*. Vol. 3 (Part A). Academic Press, London.
- Lynch, C. D. 1981. The status of the Cape grey mongoose, *Herpestes pulverulentus* Wagner, 1839 (Mammalia: Viverridae). *Navors. Nas. Mus. Bloemfontein* 4(5):121-168.
- Mohr, C. O. 1947. Table of equivalent populations of North American small mammals. *Amer. Midl. Nat.*, 37:223-249.
- Pimentel, D. 1955. Biology of the Indian mongoose in Puerto Rico. *J. Mammal.*, 36:62-68.
- Roberts, A. 1951. *The Mammals of South Africa*. Central News Agency, Cape Town.
- Rood, J. P. & Waser, P. M. 1978. The slender mongoose, *Herpestes sanguineus*, in the Serengeti. *Carnivore* 1(3):54-58.
- Stuart, C. T. 1974. Preliminary checklists of the flora and vertebrate fauna of Vrolijkheid Nature Conservation Station. *Investigational Report*, Dept. of Nature Conservation, Cape Province 19:1-18.
- Stuart, C. T. 1981. Notes on the mammalian carnivores of the Cape Province, South Africa. *Bontebok* 1:1-58.
- Tomich, P. Q. 1969. Movement patterns of the mongoose in Hawaii. *J. Wildl. Manage.*, 33:567-584.

Chris & Tilde Stuart, African Carnivore Survey  
P.O.Box 96, Nieuwoudtville 8180, South Africa

## Abstracts

### Small Indian mongoose in Yugoslavia.

Seven mongooses from three Adriatic islands (Mljet, Korcula, and Hvar) off the coast of Yugoslavia were examined. Comparison with material of *Herpestes auro-punctatus*, *H. edwardsi*, *H. sanguineus*, *H. javanicus*, and *H. pulverulentus* showed that they belong to *H. auro-punctatus*. The history of their introduction since 1910, present distribution, and biological remarks are given. No recent data on their presence are available on Brac Island, but they are frequent on the three above mentioned islands and on the Peninsula of Peljesac. The Adriatic population of *H. auro-punctatus* is the only one of its kind in Europe.

Tvrkovic, N. & Krystufek, B. 1990. Small Indian mongoose *Herpestes auro-punctatus* (Hodgson, 1836) on the Adriatic Islands of Yugoslavia. *Bonn. Zool. Beitr.*, 41(1):3-8.

### The mongooses of the genus *Galerella* in southern Africa.

*Galerella sanguinea* (Rüppel, 1836), also known as *Herpestes sanguineus*, was investigated for intra-population and geographical variation in southern Africa by means of univariate and multivariate methods. The analyses of intra-population variation indicated the presence of sexual dimorphism in skull size, with males being larger than females. No evidence was found to justify the recognition of subspecies in *G. sanguinea* from southern Africa. It is proposed that all of the 16 subspecies described for southern Africa be regarded as synonyms of *G. sanguinea caunii*, except for *G. swalius* which is regarded as a monotypic species, occurring sympatrically with *G. sanguinea* in the central and southern parts of Namibia.

In the genus *Galerella* following species are now recognized: *G. pulverulenta*, *G. sanguinea*, *G. nigrata*, and *G. swalius*, while the status of *G. swinnyi* is still considered uncertain on morphometric grounds.

The results of geographic variation analyses showed that:

- 1) Pelage colour is very variable geographically and correlates with substrate colour and humidity
- 2) An east-west decrease in skull size is present in *G. sanguinea*.

Watson, J. P. 1990. The taxonomic status of the Slender mongoose, *Galerella sanguinea* (Rüppel, 1836), in southern Africa. *Navors. Nas. Mus. Bloemfontein* 6(10):351-492.

### Siberian ferrets used as surrogate Black-footed ferrets in release study

Black-footed ferrets (*Mustela nigripes*) are an endangered mustelid that once spread across the Great Plains of North America. The last known population was decimated by canine distemper in 1985. Captive breeding has been successful and the next step in recovery of the species is the reintroduction of captive-raised, black-footed ferrets in the fall of 1991. Using a congeneric surrogate, the Siberian ferret (*Mustela eversmanni*), development of predator avoidance abilities, hunting effectiveness, and release techniques were tested. Even though the captive-raised Siberian ferrets often exhibited the correct behavioral responses, they did not always perform them as efficiently as wild-raised animals. Mortality will probably be heavy with any release of captive-raised animals. For a captive-raised carnivore the size of a black-footed ferret, it will be no easy task to locate and kill a prey as large as a Prairie dog and not be killed by a larger predator in the process.

B. Miller. 1991. *CBSG News* (Newsletter of the IUCN/SSC Captive breeding Specialist Group) 2(1):10-11.

# Some notes on terrestrial mustelids in the central Paraguayan chaco

Dan BROOKS

There are three extant genera of terrestrial mustelids in the Paraguayan chaco today. These are the Lesser grison (*Galictis cuja*), the Tayra (*Eira barbara*), and Molina's hog-nosed skunk (*Conepatus chinga*).

Over the course of a year, random status surveys were made and data were collected on terrestrial mustelids in the central Paraguayan chaco. Current status for *G. cuja* is indeterminate. *E. barbara* status is undetermined, but may well be endangered, as none were encountered. Density studies for *C. chinga* were done along a frequently traveled route. Most mustelids in this region are endangered by hunting, human disturbance, and/or habitat destruction.

## Methods

From August 1989 to August 1990 stationed at Estancia Toledo (22°33'S, 60°30'W; 35 km west of the Mennonite Colony Filadelfia, Boqueron, Paraguay), numerous random surveys were made and data were collected on the terrestrial mustelids in this area (Fig. 1). All localities and areas were driven to by vehicle, and surveyed on foot. Density studies for *C. chinga* involved driving a specific length of measured road, determining the overall area in square km, and thus obtaining one *C. chinga*/number of square km.

## Habitat

The majority of the central Paraguayan chaco has been cleared for cattle land. Other reports and sightings indicate this to be the case throughout much of the chaco, some areas more so than others.

Most of the modified land in the central chaco is typical of capoeira savannah and to a lesser extent caatinga - many short, thorny bushes, shrubs, and cacti; with a scattering of trees up to 13 m high.

The dense forest canopy, secondarily frequent to pastureland, is mid-height, and contains thorny, impenetrable foliage. Small segments of the thick forest are sometimes left in tracts when the land is being cleared for agrarian purposes.

## The Lesser grison (*Galictis cuja*)

### Status

This species is found most frequently on savannah peripheries. Not as stable as the hog-nosed skunk, but more common than the tayra, the grison is often blamed for eating poultry, and thus is hunted.

The species is important to farmers, as its diet includes a high content of granivorous rodents. In some regions *Galictis* has been kept to control rodent populations, or for sport hunting similar to European ferreting.

### Behavior

Most likely this species forms seasonal familial groups. On May 25 (late fall) a single individual was seen. On August 1 (late winter) a group of five, including adults and juveniles was seen. The lesser grison was observed to be diurnal, as both recordings

were in morning daylight, between 0745 and 0900 hours. These animals are more inquisitive than most carnivores, but are quick to flee at the first sign of danger, scampering through the tall grass. The energetically expensive retreat involves a high metabolism, with slow progress, as the body is low to the ground, and the legs are considerably short.

*G. cuja* population density is lower than for *Conepatus chinga* (i. e. < one individual/1.4 km<sup>2</sup>).

## The Tayra (*Eira barbara*)

### Status

The generic name for this species is derived from the Guarani's Indian language name, E'ira. Not one sign of evidence was found for this species. By the same token, the locals were not overly familiar with the E'ira. One old mounted specimen resides in Filadelfia's Unger museum; thus it occurred in this area to some extent historically.

Although this species is highly adaptable in other areas of its range, it is possible that the conversion of forest to cattle land was a leading cause for its disappearance in the Paraguayan chaco. No doubt if this species was present, it would be blamed for killing livestock, and thus hunted frequently. As tayras are very energetic, they are capable of destroying fruit trees while foraging, and thus in some areas are frequently shot for this reason.

The tayra holds an important place in folklore for many cultures. In some areas, various body parts of this species are used for aphrodisiacs and other ethnic products. It would be worthwhile to initiate an intensive study on this species, which would yield interest by the local community to help conserve the tayra.

## Molina's hog-nosed skunk (*Conepatus chinga*)

### Status

The pelage of *C. chinga* in this area is very similar to the Andean dwelling *C. rex*, in that both species exhibit white along the dorsum, from the forehead to the tip of the tail.

Most concentrated in grass-savannah habitat of the central chaco, *C. chinga* is the most stable of the three mustelids, and among the most abundant of carnivores in the area. The infamous odor does not give this species a bad reputation, as is the case in North America. Road-kills are the major cause of mortality. This species is sometimes shot for eating poultry eggs or for other reasons.

This skunk is economically important, as its diet includes a high content of crop-damaging insects. Historically more than today, various parts from this species were used for ethnic medicine and pelt material.

### Behavior

Like other skunks, this species erects its large white 'tail plume' as an indicator to ward off potential predators. This species is solitary, as only monogamous individuals (more than nine) were



recorded. Skunks were observed to be active diurnally just as much as nocturnally.

On July 10, 1990 at 1240 h, one individual was seen catching and consuming a small green tegu lizard. Although lethargic locomotion is typical of *C. chinga*, the prey item was seized quickly. After a little quarrel, the dying prey was carried in the mouth to a shaded area bush, where the then dead lizard was held in the forepaws and slowly eaten. Eating lasted 23 min, whereupon the skunk slowly left the area through long grass.

#### Seasonal demography

For one year, signs of *C. chinga* were recorded at project base Estancia Toledo and a 30 km radius surrounding the area. Evidence of this species was found from July-November (mid winter-late spring), and April-May (mid-late fall). The species was seen most frequently during July and August (mid-late winter). This vicinity is among the most developed areas in the chaco.




Fig. 1. Schematic map of Paraguay indicating the survey area.

#### Density

Hog-nosed skunks were seen along a route from Estancia Toledo to Filadelfia (22°33'S, 60°12'W). All live individuals sighted within 10 m of the center of the road (70 km round trip) were counted, obtaining an area of 1.4 km<sup>2</sup>/trip. Thus estimating a density of one individual/1.4 km<sup>2</sup> in a region that is highly developed for cattle land (Estancia Toledo and Colonia Fernheim).

#### Conclusions

From a year of random surveys we may conclude that in the area of the central Paraguayan chaco, the lesser grison is indeterminate or threatened, the tayra is endangered, and Molina's hog-nosed skunk is stable at present. Average density for *Conepatus chinga* in this area is one individual/km<sup>2</sup>, while *Galictis cuja* density is considerably lower. Some behavioral observations were made on *G. cuja* and *C. chinga*, and habitat is described generally.

There are substantial threats to the terrestrial mustelids of this region. All species are blamed for some form of agricultural damage, and thus are persecuted. A local expression is 'Carry a gun, if it moves shoot it'; thus these species are also hunted for sport. The tayra is probably most threatened with habitat destruction in this area. A further potential threat to all species is human disturbance (i. e. road-kills).

All three species have historically played an important role to South American people. These species are economically important to agriculturists for crop-predator control. Various body parts have been used for ethnic medicine or for pelts. The lesser grison was once used for a hunting sport, similar to ferreting in Europe. As the tayra has played an important role in traditional folklore in many areas, it may be feasible to develop a conservation oriented program for this species.

#### Acknowledgements

Financial support was provided by Dr. Kurt Benirschke and the Foundation for Endangered Animals. Local logistics were provided by the Zoological Society of San Diego's C.R.E.S. through the support of Proyecto Tagua (the Tagua Project).

409 University, # 204-N, Lubbock, Texas 79401, USA

## Coming events

**The Biology and Management of Martens and Fishers**  
Laramie, Wyoming, USA, May 29 - June 1, 1991.

A Symposium bringing together biologists and managers from North America and Eurasia. Other conference activities will include panel discussions, a field trip, banquet, picnic, mini-tours, etc. Registration forms and information available from:

Marten-Fisher Symposium Organizing Committee  
Department of Zoology and Physiology  
University of Wyoming, Laramie, WY 82071-3166, USA  
Tel. (307)766-2357

**The 6th International Colloquium on the Ecology and Taxonomy of Small African Mammals**

Mitzpe Ramon, Israël, 11 - 16 August, 1991

Information available from:

Abraham Haim  
University of Haifa  
Oranim  
P.O. Kiryat Tivon, 36910  
Israel

# Owston's palm civet, *Chrotogale owstoni*, in captivity

Nguyen Xuan Dang, Pham Trong Anh, Ngo Bich Nhu, and Le Chan

Owston's palm civet is the sole living representative of the genus *Chrotogale*. The species is endemic in northern Vietnam, northern Laos, and a small adjoining area of China. The IUCN/SSC Mustelid & Viverrid Specialist Group has listed it as one of the highest "Conservation Priority Species" (Schreiber et al., 1989). In Vietnam, Owston's palm civets still survive in small numbers in some areas of their former range. Cuc Phuong National Park is the only reserve where the occurrence of the species can be confirmed. But, at present, considerable hunting pressure on *Chrotogale* still exists.

The "Institute of Ecology and Biological Resources" in Hanoi is now conducting a programme of conservation of Owston's palm civet in Vietnam. The first attempt of captive breeding of the species is being carried out as a cooperative Captive Breeding Project between the Institute and Hanoi Zoo.

The cages in which the animals are kept are 2 m long, 1.5 m wide, and 2.5 m high; they are made of steel and covered with fine wire-netting to prevent rats and mice from entering the cages. The roof is covered with palm leaves. Some bamboo sheets are covering the sides to avoid direct sunshine in summer and strong and cold wind in winter. Inside the cages there are some thick branches for climbing and either a wooden box (0.4 m x 0.6 m x 0.4 m) or a piece of a big tree trunk with a large central hole for nesting. The floor is cleaned every day. Each cage contains one or two animals.

The captive animals prefer fresh lean meat and sweet soft fruits, especially ripe bananas. The usual diet consists of beef meat, chicken, both in small pieces, earthworms and ripe bananas. Occasionally small birds, chicken's eggs, frogs, insects (grasshoppers, crickets, etc.) and other fruit like ripe papaya are given. The daily diet includes both fruit and meat. The animals are fed two times a day: between 9 and 10 and at 16.30. The meat is given only in the afternoon feeding. In general the animals only eat a part of the food given. Fresh drinking water is always available and is changed every morning.




Fig. 1. First live Owston's palm civets (*Chrotogale owstoni*) photographed. Two one month-old females (Vc-2 and Vc-3, see text), one of them lying on adult female (Vc-1). Photo by Nguyen Xuan Dang.

Table 1. Growth and weight gain of Owston's palm civets in captivity.

Name	Date	Body weight g	Head+ body length mm	Tail mm	Ear mm
Vc-1: adult female	20.07.1990	2.300	630	430	50
	20.08.1990	2.600	630	430	50
	21.09.1990	2.900	630	435	50
	22.10.1990	3.400	635	440	55
	28.11.1990	4.200	635	440	55
Vc-2: young female	20.07.1990	320	380	290	25
	20.08.1990	800	400	290	30
	16.09.1990	1.500	450	350	35
		Died			
Vd-1: adult male	20.07.1990	2.500	618	415	58
	20.08.1990	2.800	620	420	60
	7.09.1990	2.700	625	425	60
		Died			
Vc-3: young female	20.07.1990	310	370	285	25
		Died			
Vd-2: subadult male	28.10.1990	1.400	450	350	35
	22.12.1990	2.000	620	400	40
Vc-4: adult female	28.10.1990	3.000	630	430	50
	22.12.1990	3.800	635	440	56
Vc-5: adult female	25.12.1990	2.100	600	400	40

Table 1 shows the growth rate and weight gain of the seven kept animals. All specimens showed weight gain but unfortunately three died.

The young females Vc-2 and Vc-3 (Fig. 1) have been kept since they were about one month old. They were kept in the company of adult female Vc-1 whose young died in an accident just before captivity. The young females fed on both milk of female Vc-1 and on the given diet. After a week young Vc-3 suddenly died of an unknown disease. The animal which looked alright and was normally active suddenly became weak, was found lying down and died after two hours. Autopsy did not reveal any disease symptoms. Young Vc-2 grew well until she was three months old and then also died of unknown causes. Two days before her death the animal showed paralysis in one hind leg. The next morning the other hind leg was also paralysed. Treatment with penicilline and streptomycin gave no result and after becoming paralysed in both fore legs also in the afternoon, the animal died during the night.

Adult male Vd-1 was kept since May 1990. During its capture the animal was severely bitten and two wounds of ca. 20 mm diameter remained on each side of his pelvis. During its captivity the animal's health had slightly improved but the wounds did not heal in spite of the antibiotic treatment. On 6 Sept. 1990 the animal became very weak and died the next day. Adult female Vc-1 sometimes showed diarrhoea but each time recovered after biseptol treatment. Earlier on, Hanoi Zoo had kept some Owston's palm civets but all the animals died of unknown causes.

# Conservation program of *Chrotogale owstoni* Thomas, 1912: Some first results

H. J. ADLER

In this issue of the newsletter Nguyen Xuan Dang *et al.* report the first experiences in keeping Owston's palm civets. Thus the status in captivity is no longer "no records" (Schreiber *et al.*, 1989). This is a first important step to inquire into the almost unknown habits of these animals.

As reported in the a. m. article, *Chrotogale* have been kept at Hanoi Zoo several times, but not always under the conditions described by N. X. Dang. When inquiring after the situation of *Chrotogale* in Hanoi in November 1990 at the suggestion of IUCN/SSC M&VSG chairman R. Wirth, I found three animals at the zoo, two of them being kept under provisional conditions. My conclusion was that a more severe infestation with endoparasites caused by stress may be an important cause of death. The responsible colleagues at the Institute of Ecology & Biological Resources and at the zoo, however, endeavoured to improve the conditions for keeping the animals. Another reason for the high losses may be an incorrect diet. As described, the *Chrotogale*'s diet included a high percentage of fruit, but it is supposed that the diet should consist mainly of insects, worms and perhaps small vertebrates. This hypothesis is supported by the fact that the labels of two collected specimens carry the information that the stomachs contained earthworms (as mentioned by Nowak & Paradiso, 1983). Pham Chong Ahn (1980) also stated: "Chiefly insects, worms, frogs, and less than 20% fruit".

All these facts as well as the situation in the wild show that a conservation program aiming at improved conservation of wild populations and including a general status survey of the species is urgently needed. Part of the general project should be to improve conditions for a pilot captive breeding attempt to allow Vietnamese scientists to study the biology of the species. Therefore a training program for the staff being entrusted with captive breeding in Vietnam is also necessary. But all these efforts are beyond the possibilities of the Vietnamese colleagues due to the country's extremely bad economical conditions and to their low rank in the official Governmental Programs. Therefore it is very meritorious

that Dr. R. Faust, chairman of the Zoologische Gesellschaft Frankfurt von 1858 e. V., is prepared to finance, together with the Wildlife Preservation Trust, Jersey, a training program for Nguyen Xuan Dang. The director of the Institute of Ecology & Biological Resources in Hanoi has declared his great interest in future projects.

I was also shown an island about 100 km from Hanoi where a breeding project for the Vietnamese Sika deer (*Cervus nippon pseudaxis*) has already started and which is planned to be also the location for a *Chrotogale* breeding station. Under a special agreement the Zoological Gardens of Frankfurt will receive the first specimens of *Chrotogale* this year in order to support the efforts for conservation of this species in Vietnam by a research and breeding project. Moreover, the Zoologische Gesellschaft Frankfurt von 1858 e. V. will participate in financing the intended Conservation Program.

My last year's stay in Hanoi and in the Cuc Phuong National Park was financially supported by the Zoologische Gesellschaft für Arten- und Populationsschutz, Munich. The conversations and reached agreements justify the hope for the realization of concrete conservation efforts for *Chrotogale owstoni*.

## References

- Nowak, R. M. & Paradiso, J. L. (eds.) 1983. *Walker's mammals of the world*. The John Hopkins Univ. Press, Baltimore-London.
- Pham Chong Ahn. 1980. Morphology and ecology of Viverridae in Vietnam. *Zool. Zhurn.*, 59(6):905-914. (In Russian)
- Schreiber, A. *et al.* 1989. *Weasels, civets, mongooses, and their relatives. An action plan for the conservation of the mustelids and viverrids*. IUCN, Gland.

Allwetterzoo Münster, Sentruper Str., 309,  
W-4400 Münster, Germany

*Continued from page 7*

## Recommendations & Appeal for international cooperation

*Chrotogale owstoni* being a very rare and endangered animal in the wild, their losses in captivity should also be prevented as much as possible. We obtained some results in keeping this species but our losses have been great. One reason is that we have no sufficient financial support to provide our animals with better conditions. The other reason is that we have not enough experience in keeping this kind of carnivores. In view of this international collaboration in captive breeding of Owston's palm civet is of the utmost importance and a training programme in techniques of captive breeding of endangered species for our staffs is a necessity.

We are most grateful to the Directorates of IEBR and Hanoi Zoo for constant support for Project implementation. Our many thanks are due to R. Wirth, R. Ratajszczak, R. Cox, and H. Van

Rompaey of the IUCN/SSC Mustelid & Viverrid Specialist Group for their encouragement and invaluable recommendations.

## Reference

- Schreiber, A., Wirth, R., Riffel, M. & Van Rompaey, H. 1989. *Weasels, civets, mongooses, and their relatives. An Action Plan for the conservation of mustelids and viverrids*. IUCN, Gland, pp. 43, 72-75.

Nguyen Xuan Dang and Pham Trong Anh,  
Institute of Ecology and Biological Resources,  
NCSR Vietnam, Nghia do-Tu Liem-Hanoi,  
Vietnam.

Ngo Bich Nhu and Le Chan, Hanoi Zoo,  
Hanoi, Vietnam.

# Restoration of the endangered Black-footed ferret

Tim W. CLARK

The current Black-footed ferret restoration effort is now in its 10th year. Much of the progress is summarized in two annotated bibliographies (Casey *et al.*, 1986; Reading & Clark, in press).

Discovery of a small ferret population near Meeteetse, Wyoming, in 1981 brought hope for species recovery (Clark, 1989). About 300 ferrets were studied, with a peak population of 129 in 1984. In 1985, a catastrophe struck when canine distemper killed most ferrets. The species reached a low in early 1986 of about 10 individuals. The ferret is extinct in the wild today and only exists in three captive populations totalling about 186 animals. Continued increases in captive populations, reintroduction, and successful management offer an opportunity for species restoration.

Discussion about how to restore ferrets was part of the Meeteetse conservation work from the beginning. In 1983, Richardson *et al.* (1986) first formally examined recovery strategies and strongly recommended the captive-rearing/translocation option.

The 1988 Fish and Wildlife Service Recovery Plan objective is: "To ensure immediate survival of the black-footed ferret by: (1) Increasing the captive population of black-footed ferrets to a census size of 200 breeding adults by 1991; (2) establishing a prebreeding census population of 1500 free-ranging black-footed ferret breeding adults in 10 or more populations with no fewer than 30 breeding adults in any population by the year 2010; and (3) encouraging the widest possible distribution of reintroduced black-

footed ferret populations" (Fish and Wildlife Service 1988:19). Ferret numbers currently in captivity and plans for reintroduction approximate these goals.

Captive breeding efforts got off to a dismal start but have since been very promising. Eighteen wild ferrets constituted the entire founder population. In 1990, 66 young were born and at the start of 1991, 180 animals are in captivity. Ferrets are now at the Conservation and Research Center, a branch of the National Zoological Park, the Henry Doorly Zoo, and a Wyoming Game and Fish Department Facility where they have been for several years. Additional populations are now at the Louisville Zoo, Louisville, Kentucky (12 animals) and Cheyenne Mountain Zoo, Colorado Springs, Colorado (10 animals). Much of the success of the captive breeding effort is due to advice and assistance provided by the Captive Breeding Specialists Group of the IUCN/SSC.

Reintroducing the Meeteetse ferrets to several new sites was part of the first conservation and research plans for the newly discovered ferrets (Clark, 1981) and was formally recommended in 1983 and 1985, with the reports of Richardson *et al.* (1986) and Forrest *et al.* (1985). Reintroduction became formalized as the basic recovery strategy in government plans in 1987 and 1988 (Wyoming Game and Fish Department 1987, Fish and Wildlife Service 1988). A Black-footed ferret Interstate Coordinating Committee was established in 1986. The Committee meets at least once yearly and otherwise communicates through a Fish and Wildlife Service Secretariat.



Black-footed ferret (*Mustela nigripes*)

Several reintroduction sites have been located throughout the western United States. One of the biggest is in northcentral Montana, estimated to be capable of supporting about 500 ferret families. Restoring ferrets to the wild is now a matter of developing successful reintroduction techniques and subsequent management procedures.

Many details remain to be worked out before ferrets can be restored. Successful reintroduction and management of the black-footed ferret is an opportunity to restore the unique prairie dog ecosystem (Clark *et al.*, 1989; Miller *et al.*, in press). Hopefully, beginning in 1991 with the first introductions, and extending over several years, ferrets will be reestablished to the prairies and intermountain basins of North America. All involved hope ferret restoration will go smoothly and quickly.

#### References

- Casey, D.E., DuWaldt, J. & Clark, T. W. 1986. An annotated bibliography of the black-footed ferret. *Great Basin Nat. Memoirs* 8:185-208.
- Clark, T. W. 1981. *The Meeteetse black-footed ferret conservation studies: a proposal*. Box 2705, Jackson, WY. 78 pp.
- Clark, T. W. 1989. Conservation biology of the black-footed ferret, *Mustela nigripes*. *Wildlife Preservation Trust, Special Scientific Report* 3:1-175.
- Clark, T. W., Hinckley, D. & Rich, T. (eds.) 1989. The prairie dog ecosystem: managing for biological diversity. *Montana Bureau of Land Manage. Wildl. Techn. Bull.* 2:1-55.

- Forrest, S. C., Clark, T. W., Richardson, L. & Campbell III, T. M. 1985. Black-footed ferret habitat: some management and reintroduction considerations. *Wyoming Bureau of Land Manage. Wildl. Techn. Bull.* 2:1-75.
- Fish and Wildlife Service. 1988. *Black-footed ferret recovery plan*. U. S. Fish and Wildlife Service, Denver, CO., 154 pp.
- Miller, B., Wimmer, C., Biggins, D. & Reading, R. In press. A proposal to conserve black-footed ferrets and the prairie dog ecosystem. *Environmental Management*.
- Reading, R. P. & Clark, T. W. In press. Black-footed ferret annotated bibliography, 1986-1990. *Montana Bureau of Land Manage. Wildl. Techn. Bull.*, 3.
- Richardson, L., Clark, T. W., Forrest, S. C. & Campbell III, T. M. 1986. Black-footed ferret recovery: a discussion of some options and considerations. *Great Basin Nat. Memoir* 8:169-184.
- Wyoming Game and Fish Department. 1987. *A strategic plan for the management of black-footed ferrets in Wyoming*. Wyoming Game and Fish Department Report, Cheyenne WY., 70 pp.

**Chicago Zoological Society, Brookfield, IL 60513**  
**Northern Rockies Conservation Cooperative,**  
**Box 2705, Jackson, WY 83001, USA**  
**School of Forestry and Environmental Studies,**  
**Yale University, New Haven, CT 06511, USA**

## Recent publications

### Encyclopaedia of the carnivores of France

We would like to draw our readers' attention to a highly interesting series of publications in French: the "Encyclopédie des carnivores de France: espèces sauvages ou errantes, indigènes ou introduites, en métropole et dans les dom-tom". All issues are published by the "Société Française pour l'Etude et la Protection des Mammifères (S.F.E.P.M.), Bohallard, Puceul, 44390 Nort sur Erdre, France. For information write:

Mad. D. Thévenot, Grand-rue, 54200 Lagny, France.

Following issues on viverrids and mustelids have already been published:

- Henry, C., Lafontaine, L. & Mouches, A. 1988. Le blaireau (*Meles meles* Linnaeus, 1758). N° 7:35 pp.
- Labrid, M. 1986. La martre (*Martes martes* Linnaeus, 1758). N° 9:22 pp.
- Delattre, P. 1987. La belette (*Mustela nivalis* Linnaeus, 1766) et l'hermine (*Mustela erminea* Linnaeus, 1758). N° 11 & 12:73 pp.
- Camby, A. 1990. Le vison d'Europe (*Mustela lutreola* Linnaeus, 1761), Maizeret, C. 1990. Le vison d'Amérique (*Mustela vison* Schreber, 1777). N° 13 & 14:44 pp.

- Roger, M., Delattre, P. & Herrenschmidt, V. 1988. Le putois (*Mustela putorius* Linnaeus, 1758). N° 15:38 pp.
- Livet, F. & Roeder, J.-J. 1987. La genette (*Genetta genetta* Linnaeus, 1758). N° 16:33 pp.
- Charles-Dominique, P. & Moutou, F. 1987. Les carnivores des Départements et Territoires d'Outre-Mer. N° 20 & 21:27 pp.

Several issues on the canids, ursids, procyonids, and felids have been and will be published.

### Populationsökologie Marderartiger Säugetiere

The papers presented at the "Mitteleuropäisches Symposium zur Populationsökologie von Mustelidenarten" held in Spremberg, Germany from 12 to 16 April 1989 were edited by Michael Stubbe and published as "Populationsökologie Marderartiger Säugetiere" Vol. 1 & Vol. 2 (647 pp.), Kongress- und Tagungsberichte der Martin-Luther-Universität Halle-Wittenberg 1989/37 (P 39): Halle (Saale), 1989.

All the "no-otter" papers are listed separately in the "Recent literature" section of this number.