

- Hunter is a fast establishing crop, with the first grazing possible at 6-8 weeks (45-55 days).
- Providing excellent quality forage Hunter can increase liveweight gain on animals.
- Hunter was bred for tolerance to turnip mosaic virus, cauliflower mosaic virus and most clubroot strains.
- Selected for vigorous regrowth and fast recovery from grazing, Hunter has the ability to yield in the second, third and sometimes fourth regrowth cycle.
- Best suited to heavier soil conditions with periodic summer moisture or irrigation where multiple grazings can be achieved.

Hunter is a hybrid developed by crossing turnips with related Asiatic leaf vegetables of the same species. The resulting quickgrowing, leafy turnip, with minimal bulb development, is best suited to multiple grazings for summer and early-fall feed requirements.

This hybrid is excellent quality forage, providing high liveweight gains for finishing stock to target weight in summer/fall.

When given moisture, Hunter recovers very quickly from multiple grazings for excellent subsequent yields. Hunter is an integral component of "cocktail covercrops".

## **Sowing Guide**

- Seed in spring to the end of summer, with given moisture. Sow in the fall, on the west coast.
- Seed at 4 lbs/acre.
- Time until first grazing 6 to 8 weeks.
- Sow with 10 lbs/acre Crusader Italian ryegrass (2 lbs/acre Hunter) for extended grazing, from summer through fall.

## **Suitability**



## For more information please contact: pggseeds@gmail.com

Results will vary depending on all circumstances. PGG Seeds and its officers, employees, agents and licensors of intellectual property (PGG Seeds) provide no assurances, guarantees or warranties in relation to any advice, information, cultivar or product referred to or recommend, except those that by law must be provided. © PGG Seeds 2012