

AMPT™-A RU/IGR 54229PIZ

▶ **MEDICATED** ◀

For Beef Cattle on Pasture Weighing from 1050 to 1500 lbs

Growing cattle on pasture or in dry lot (stocker and feeder and dairy and beef replacement heifers): For increased rate of weight gain.

ACTIVE DRUG INGREDIENT

Monensin..... (200 mg/lb)..... 400 gm/ton

To prevent the breeding of horn flies in the manure of treated cattle only. A pesticidally active mineral feed for beef cattle containing methoprene* insect growth regulator for continuous feeding during the fly season.

ACTIVE INGREDIENT:

S-methoprene (CAS Number 65733-16-6)..... 0.014%

OTHER INGREDIENTS: 99.986%

Contains 63.64 mg of methoprene per lb. Total 100.000%

GUARANTEED ANALYSIS	INGREDIENTS
Calcium (Ca), Min. 13.80% Max. 16.35%	Magnesium Carbonate, Salt, Dicalcium Phosphate, Calcium Carbonate, Processed Grain By-Products, Magnesium-Mica, Vegetable Oil, Molasses Products, Manganese Hydroxychloride, Zinc Hydroxychloride, Iron Oxide, Zinc Oxide, Sodium Selenite, Basic Copper Chloride, Ethylenediamine Dihydriodide, Manganous Oxide, Vitamin A Supplement, Cobalt Carbonate, Lactic Acid, Vitamin E Supplement, Vitamin D ₃ Supplement, Artificial Flavor.
Phosphorus (P), Min. 4.00%	
Salt (NaCl), Min. 19.20% Max. 22.80%	
Magnesium (Mg), Min. 3.00%	
Potassium (K), Min..... none added	
Cobalt (Co), Min. 150 ppm	
Copper (Cu), Min. 1,200 ppm	
Iodine (I), Min. 200 ppm	
Manganese (Mn), Min..... 3,600 ppm	
Selenium (Se), Min. 25.0 ppm	
Zinc (Zn), Min. 4,200 ppm	
Vitamin A, Min. 200,000 IU per pound	
Vitamin D3, Min..... 5,000 IU per pound	
Vitamin E, Min..... 100 IU per pound	

KEEP OUT OF REACH OF CHILDREN CAUTION

USE AND MIXING DIRECTIONS

STORAGE AND DISPOSAL: Do not contaminate water or food by storage or disposal.

STORAGE: Store in dry area. Do not contaminate with pesticides or fertilizer.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

MIXING AND FEEDING DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product only prevents emergence of adult horn flies from the manure of treated cattle. Begin use in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity. Introduction of the product when significant numbers of adult flies are present may require the use of adulticide sprays or dusts.

This product is designed to be mixed and fed to cattle weighing from 1050 to 1500 lbs continuously at the rate of 0.25 lb per head per day. When fed at this rate, this product will provide 50 mg monensin per head per day and from 45.4 to 31.8 mgs methoprene per 100 lb body weight per month to cattle weighing from 1050 to 1500 lbs. Before feeding, this product must be mixed with non-medicated feed so that it is contained in at least 1 lb of feed. For example, blend 500 lb per ton, and feed 1 lb of this medicated mix. During the first 5 days, cattle should receive no more than 100 mg monensin per head per day. In addition, forage should be fed free choice. Provide clean, fresh water at all times.

CAUTION: Consumption of this product by sheep and goats may result in copper toxicity. Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use; do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

04117-TG

**ADM Animal Nutrition, a division of Archer Daniels Midland Company, Quincy, IL 62305-3115
Net Weight 50 lb (22.67 kg) Or As Invoiced**