
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1612A-S1 0820

OXATHIPIPROLIN	GROUP	49	FUNGICIDE
MANDIPROPAMID	GROUP	40	FUNGICIDE

Orondis® Ultra

Fungicide

This supplemental label expires on 08/17/2023 and must not be used or distributed after this date.

Active Ingredients:

Oxathiapiprolin ¹ :	2.77%
Mandipropamid ² :	23.10%
Other Ingredients:	74.13%
Total:	100.00%

¹CAS No. 1003318-67-9

²CAS No. 374726-62-2

Orondis® Ultra is formulated as a suspension concentrate and contains 0.25 lb of oxathiapiprolin and 2.08 lb of mandipropamid per gallon.

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 100-1612

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Orondis Ultra as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

Hops

Crop			
Hops			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Downy mildew (<i>Pseudoperonospora humuli</i>)	5.5 - 8.0*	Begin applications prior to disease development and continue on a 7- to 10-day interval.	Apply as a foliar application by ground, or air assisted (air blast) application. Use the higher rate when conditions are favorable for disease development, for longer application intervals, or for susceptible varieties. See Section 4.4.5 for adjuvant recommendations.
*5.5 fl oz is equivalent to 0.01 lb oxathiapiprolin and 0.09 lb mandipropamid *8 fl oz is equivalent to 0.02 lb oxathiapiprolin and 0.13 lb mandipropamid			
Integrated Pest Management			
<ul style="list-style-type: none"> Refer to Section 3.1. 			
Resistance Management:			
<ul style="list-style-type: none"> Refer to Section 3.2. Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. Do not follow soil applications of oxathiapiprolin-containing products with foliar applications of oxathiapiprolin-containing products. Use either soil applications or foliar applications but not both for disease control. Where 3 or more fungicide applications are made, use Orondis Ultra (or any other oxathiapiprolin-containing product) in no more than 33% of the applications, or a maximum of 3 applications, whichever is fewer. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table. Maximum Number of Applications per Year: DO NOT make more than 3 applications at the maximum rate per year. Minimum Application Interval: 7 days Maximum Annual Rate: 24.0 fl oz/A/year (equivalent to 0.05 lb ai/A/year oxathiapiprolin and 0.39 lb ai/A/year mandipropamid) <ol style="list-style-type: none"> DO NOT apply more than 0.09 lb ai/A/year of oxathiapiprolin-containing products. DO NOT apply more than 0.39 lb ai/A/year of mandipropamid-containing products. DO NOT use in nursery production of transplanted crops. Pre-harvest Interval (PHI): 7 days 			

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2020 Syngenta