## Subcutaneous

Prescriber's Signature:\_



Phone: (305) 221-1421

Date: \_

Cabcatariocac	Rx Internat	lional Pharmacy	Fov: (205) 224 2275
Immune Globulin	TX III CITICA	aronai i mannaoy	Fax: (305) 221-3275
Patient information		Prescriber + Shipping Informa	ation
	DOB:	Prescriber Name:	
Sex: ☐ Female ☐ Male SS #:		NPI #:	
	Wt: □ kg □ lbs Ht: □ cm □ in	Address:	
Address:		Apt/Suite: City:	
Apt/Suite: City:	State: Zip:	Contact: A	Itarnata
Phone: Al	ternate Phone:	Fax:	
Caregiver name:	Relation:	Email address:	
Local Pharmacy:	Phone:	If shipping to presciber: ☐ 1st Fill ☐ Always ☐ Never	
<b>Insurance Information (Ple</b>	ase fax a copy of front and back	of the insurance cards)	
1° Insurance Plan:	Plan ID #	Policy Holder:	Relation:
2° Insurance Plan:	Plan ID #	Policy Holder:	Relation:
<b>Clinical Information (Pleas</b>	e fax all pertinent clinical and lab	information)	
ICD-10/Diagnosis Code:			
Date of Diagnosis:		Has patient received immune globulin p	
gA deficiency:   Yes No IgA level mg/dL Date:		If yes, product information:	
IgG trough:mg/dL Date: Diabetic: □ Yes □ No Date of last infusion: Date of next infusion:			
Comorbidities: Allergies: ☐ NKDA ☐ Other:			
Prescription			
☐ Hizentra® 20%			
Immune Globulin Products			
	Weekly Sub-Q dose = IVIG Dose (g) x 1.37 / IVIG weekly interval originally given		
	☐ HyQvia® 10% Complete the Mylg source form at <a href="http://www.myigsource.com/pdf/Baxalta_Rx_Forms.pdf">http://www.myigsource.com/pdf/Baxalta_Rx_Forms.pdf</a> for orders and patient registration.		
	☐ GammaKed <sup>™</sup> 10% ☐ Gammagard liquid <sup>®</sup> 10% ☐ Gamunex-C <sup>®</sup> 10%		
	Weekly Sub-Q dose = IVIG Dose (g) x 1.53 / IVIG weekly interval originally given		
Therapy Regimen	Dose: grams		
	# Doses: Refill:		
	Administration rate and number of sites: ☐ Per manufacture guidelines, as tolerated ☐		
	☐ Acetaminophen Take mg by mouth every 4-6 hours as needed for fever and/or headache		
Other Medications	Take mg by mount every 4-6 nouts as needed for never and/or neadache		
	☐ Diphenhydramine Take mg by mouth every 4-6 hours as needed for itching		
	Drug	Strength:	Othir
	Drug: Directions:	Strength	Refills:
	Bircolloris.		Tems.
Anaphylaxis Orders and Medications	Orders:		
	<ol> <li>Stop infusion</li> <li>Call 911 and prescribing physic</li> </ol>	ian	
	<ol><li>Administer medications below a</li></ol>		
	☐ Epinephrine		
	□ EpiPEN® - Administer 0.3 mg (1:1000) subcutaneously as needed (≥ 30 Kg or ≥ 66 lbs)		
		□ EpiPEN Jr® - Administer 0.15 mg (1:2000) subcutaneously as needed (< 30 Kg or < 66 lbs)	
	Qty: Refill:		
Ancillary Supplies	☐ As needed for proper administration and disposal of medication		
0.00	To train patient/caregiver in Subcutaneous Immune Globulin administration, provide education related to disease		
Skilled Nursing Visits	state/therapy and assess general status. Typically 3 training visits required. Once trained and able to return demonstrate, patient/caregiver to self-adminster Subcutaneous Immune Globulin medication independently		
Administration procedures to be	e followed per pharmacy protocol.		
		(DANAM / 1)	
Prescription will be filled with ge	eneric (if available) unless prescriber wri	ites "DAW" (dispense as written):	

I authorize Rx International Pharmacy and its representatives to act as an agent to initiate and execute the insurance prior authorization process.