

Math 2371 Calc III

Sample Test 3 - Solns

1. (i) Is the following vector field conservative? If so, find the potential ϕ

$$\vec{F} = \langle yz + 3, xz + 4y, xy + 3z^2 \rangle.$$

Soln. Since $\nabla \times \vec{F} = 0$ then yes, the vector field is conservative. Thus f exists such that $\vec{F} = \vec{\nabla} f$ so

$$\begin{aligned} f_x = 2xy &\Rightarrow f = x^2y + A(y, z) \\ f_y = x^2 + z^2 &\Rightarrow f = x^2y + yz^2 + B(x, z) \\ f_z = 2yz &\Rightarrow f = yz^2 + C(x, y) \end{aligned}$$

Therefore we see that

$$f = x^2y + yz^2 + c.$$

(ii) Evaluate the following where c is any path from $(0, 0, 0)$ to $(1, 2, 3)$.

$$\int_c (yz + 3)dx + (xz + 4y)dy + (xy + 3z^2)dz$$

Soln.

$$\int_c (yz + 3)dx + (xz + 4y)dy + (xy + 3z^2)dz = x^2y + yz^2 \Big|_{(0,0,0)}^{(1,2,3)} = 20$$

2. Evaluate the following line integral $\int_c xy \, ds$ where c is counterclockwise direction around a circle of radius 1 from $(1, 0)$ to $(0, 1)$.

Soln. Here parameterize the circle of radius $r = 1$ with $x = \cos t, y = \sin t$. Now

$$\frac{dx}{dt} = -\sin t, \quad \frac{dy}{dt} = \cos t \tag{1.1}$$

so

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \sqrt{\sin^2 t + \cos^2 t} dt = dt \tag{1.2}$$

To evaluate the integral is to evaluate

$$\int_0^{\pi/2} \cos t \sin t dt = \frac{1}{2} \sin^2 t \Big|_0^{\pi/2} = \frac{1}{2} \tag{1.3}$$

3. Green's Theorem is

$$\int_C P dx + Q dy = \iint_{R_1} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Verify Green's Theorem where $\vec{F} = \langle y^2, x^2 + 2xy \rangle$ where R is the region bound by the curves $y = x^2$, $y = 1$ and $x = 0$ in Q1.

Soln. Again, we have three separate curves which we denote by C_1 , C_2 and C_3 .

$$C_1 : \text{ Here } y = x^2, dy = 2x dx \text{ so } \int_0^1 x^4 dx + (x^2 + 2x^3)2x dx = 3/2$$

$$C_2 : \text{ Here } y = 1, dy = 0 \text{ so } \int_1^0 dx = -1$$

$$C_3 : \text{ Here } x = 0, dx = 0 \text{ so } \int_{C_3} 0 = 0$$

$$\text{Thus } \int_C y^2 dx + (x^2 + 2xy) dy = 3/2 - 1 + 0 = 1/2.$$

Since $P = y^2$ and $Q = x^2 + 2xy$ then $Q_x - P_y = 2x + 2y - 2y = 2x$ so

$$\iint_R (Q_x - P_y) dA = \int_0^1 \int_{x^2}^1 2x dy dx = 1/2$$

4. Evaluate $\iint_S z dS$ where S is the surface of the paraboloid $z = 1 - x^2 - y^2, z \geq 0$.

Soln. Since $z = 1 - x^2 - y^2$ then $dS = \sqrt{1 + z_x^2 + z_y^2} dA = \sqrt{1 + 4x^2 + 4y^2} dA$ and so far we have $\iint_R (1 - x^2 - y^2) \sqrt{1 + 4x^2 + 4y^2} dA$ where the region of integration is the circle $x^2 + y^2 = 1$. Switching to polar gives

$$\int_0^{2\pi} \int_0^1 (1 - r^2) \sqrt{1 + 4r^2} r dr d\theta = \left(\frac{5\sqrt{5}}{24} - \frac{11}{120} \right) 2\pi$$

5. Find the flux $\iint_S \vec{F} \cdot \hat{n} dS$ of the vector field $\vec{F} = \langle 2x, 2y, 2z + 2 \rangle$ through the surface of the plane $x + y + z = 1$ in the first quadrant.

Soln. The unit normal to the surface is given by $\vec{n} = \frac{\langle 1, 1, 1 \rangle}{\sqrt{3}}$. For this surface $dS = \sqrt{1 + 1 + 1} dA$ so

$$\begin{aligned} \vec{F} \cdot \vec{n} dS &= \iint_S \langle 2x, 2y, 2z + 2 \rangle \cdot \frac{\langle 1, 1, 1 \rangle}{\sqrt{3}} \sqrt{1 + 1 + 1} dA \\ &= \iint_S (2x + 2y + 2z + 2) dA \end{aligned}$$

Bringing in the surface we obtain

$$\int_0^1 \int_0^{1-x} 4 dy dx = 2$$