

# Join Us for an Expert Discussion on LYNPARZA<sup>®</sup> (olaparib)

## LYNPARZA<sup>®</sup> (olaparib): A Targeted Treatment for Eligible Patients With gBRCAm, HER2-Negative Early or Metastatic Breast Cancer

September 28, 2024  
06:00 PM – 08:30 PM Pacific Daylight Time

### Presented by

Linda Buck  
MSN, NP-C, AOCNP

### Location

Rossblu  
1124 San Julian Street  
Los Angeles, CA 90015

### RSVP is required by

9/25/2024

To register or for more information, contact  
Ashley Hart at (310) 905-9227 [Ashley.Hart@astrazeneca.com](mailto:Ashley.Hart@astrazeneca.com)

### IMPORTANT SAFETY INFORMATION

#### CONTRAINDICATIONS

There are no contraindications for LYNPARZA.

#### WARNINGS AND PRECAUTIONS

**Myelodysplastic Syndrome/Acute Myeloid Leukemia (MDS/AML):** Occurred in approximately 1.2% of patients with various *BRCAm*, *gBRCAm*, *HRR* gene-mutated or HRD-positive cancers who received LYNPARZA as a single agent or as part of a combination regimen, consistent with the approved indications, and the majority of events had a fatal outcome. The median duration of therapy in patients who developed MDS/AML was approximately 2 years (range: <6 months to >4 years). All of these patients had previous chemotherapy with platinum agents and/or other DNA-damaging agents, including radiotherapy.

In SOLO-1, patients with newly diagnosed advanced *BRCAm* ovarian cancer, the incidence of MDS/AML was 1.9% (5/260) in patients who received LYNPARZA and 0.8% (1/130) in patients who received placebo based on an updated analysis. In PAOLA-1, of patients with newly diagnosed advanced ovarian cancer with HRD-positive status, the incidence of MDS/AML was 1.6% (4/255) in patients who received LYNPARZA and 2.3% (3/131) in the control arm.

In SOLO-2, patients with *BRCAm* platinum-sensitive relapsed ovarian cancer, the incidence of MDS/AML was 8% (15/195) in patients who received LYNPARZA and 4% (4/99) in patients who received placebo. The duration of LYNPARZA treatment prior to the diagnosis of MDS/AML ranged from 0.6 years to 4.5 years.

Do not start LYNPARZA until patients have recovered from hematological toxicity caused by previous chemotherapy (≤Grade 1). Monitor complete blood count for cytopenia at baseline and monthly thereafter for clinically significant changes during treatment. For prolonged hematological toxicities, interrupt LYNPARZA and monitor blood count weekly until recovery.

If the levels have not recovered to Grade 1 or less after 4 weeks, refer the patient to a hematologist for further investigations, including bone marrow analysis and blood sample for cytogenetics. Discontinue LYNPARZA if MDS/AML is confirmed.

**Pneumonitis:** Occurred in 0.8% of patients exposed to LYNPARZA monotherapy, and some cases were fatal. If patients present with new or worsening respiratory symptoms such as dyspnea, cough, and fever, or a radiological abnormality occurs, interrupt LYNPARZA treatment and initiate prompt investigation. Discontinue LYNPARZA if pneumonitis is confirmed and treat patient appropriately.

**Venous Thromboembolism (VTE):** Including severe or fatal pulmonary embolism (PE) occurred in patients treated with LYNPARZA. In the combined data of two randomized, placebo-controlled clinical studies (PROfound and PROpel) in patients with metastatic castration-resistant prostate cancer (N=1180), VTE occurred in 8% of patients who received LYNPARZA, including pulmonary embolism in 6%. In the control arms, VTE occurred in 2.5%, including pulmonary embolism in 1.5%. Monitor patients for signs and symptoms of venous thrombosis and pulmonary embolism, and treat as medically appropriate, which may include long-term anticoagulation as clinically indicated.

**Embryo-Fetal Toxicity:** Based on its mechanism of action and findings in animals, LYNPARZA can cause fetal harm. Verify pregnancy status in females of reproductive potential prior to initiating treatment.

#### Females

Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception during treatment and for 6 months following the last dose.

#### Males

Advise male patients with female partners of reproductive potential or who are pregnant to use effective contraception during treatment and for 3 months following the last dose of LYNPARZA and to not donate sperm during this time.

#### ADVERSE REACTIONS—First-Line Maintenance *BRCAm* Advanced Ovarian Cancer

Most common adverse reactions (Grades 1-4) in ≥10% of patients who received LYNPARZA in the first-line maintenance setting for SOLO-1 were: nausea (77%), fatigue (67%), abdominal pain (45%), vomiting (40%), anemia (38%), diarrhea (37%), constipation (28%), upper respiratory tract infection/influenza/nasopharyngitis/bronchitis (28%), dysgeusia (26%), decreased appetite (20%), dizziness (20%), neutropenia (17%), dyspepsia (17%), dyspnea (15%), leukopenia (13%), urinary tract infection (13%), thrombocytopenia (11%), and stomatitis (11%).

Most common laboratory abnormalities (Grades 1-4) in ≥25% of patients who received LYNPARZA in the first-line maintenance setting for SOLO-1 were: decrease in hemoglobin (87%), increase in mean corpuscular volume (87%), decrease in leukocytes (70%), decrease in lymphocytes (67%), decrease in absolute neutrophil count (51%), decrease in platelets (35%), and increase in serum creatinine (34%).

Please see additional Important Safety Information on the back of this card. Please see accompanying complete Prescribing Information, including Medication Guide or visit [https://www.azpicentral.com/lynparza\\_tb/lynparza\\_tb.pdf](https://www.azpicentral.com/lynparza_tb/lynparza_tb.pdf)

**Lynparza<sup>®</sup>**  
olaparib  
tablets 150 mg

## IMPORTANT SAFETY INFORMATION (Cont'd)

### ADVERSE REACTIONS—First-Line Maintenance Advanced Ovarian Cancer in Combination with Bevacizumab

Most common adverse reactions (Grades 1-4) in  $\geq 10\%$  of patients treated with LYNPARZA/bevacizumab and at a  $\geq 5\%$  frequency compared to placebo/bevacizumab in the **first-line maintenance setting for PAOLA-1** were: nausea (53%), fatigue (including asthenia) (53%), anemia (41%), lymphopenia (24%), vomiting (22%), and leukopenia (18%). In addition, the most common adverse reactions ( $\geq 10\%$ ) for patients receiving LYNPARZA/bevacizumab irrespective of the frequency compared with the placebo/bevacizumab arm were: diarrhea (18%), neutropenia (18%), urinary tract infection (15%), and headache (14%).

In addition, venous thromboembolism occurred more commonly in patients receiving LYNPARZA/bevacizumab (5%) than in those receiving placebo/bevacizumab (1.9%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients for LYNPARZA in combination with bevacizumab in the **first-line maintenance setting for PAOLA-1** were: decrease in hemoglobin (79%), decrease in lymphocytes (63%), increase in serum creatinine (61%), decrease in leukocytes (59%), decrease in absolute neutrophil count (35%), and decrease in platelets (35%).

### ADVERSE REACTIONS—Maintenance gBRCAm Recurrent Ovarian Cancer

Most common adverse reactions (Grades 1-4) in  $\geq 20\%$  of patients who received LYNPARZA in the **maintenance setting for SOLO-2** were: nausea (76%), fatigue (including asthenia) (66%), anemia (44%), vomiting (37%), nasopharyngitis/upper respiratory tract infection (URI)/influenza (36%), diarrhea (33%), arthralgia/myalgia (30%), dysgeusia (27%), headache (26%), decreased appetite (22%), and stomatitis (20%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients who received LYNPARZA in the **maintenance setting for SOLO-2** were: increase in mean corpuscular volume (89%), decrease in hemoglobin (83%), decrease in leukocytes (69%), decrease in lymphocytes (67%), decrease in absolute neutrophil count (51%), increase in serum creatinine (44%), and decrease in platelets (42%).

### ADVERSE REACTIONS—Adjuvant Treatment of gBRCAm, HER2-Negative, High-Risk Early Breast Cancer

Most common adverse reactions (Grades 1-4) in  $\geq 10\%$  of patients who received LYNPARZA in the **adjuvant setting for OlympiA** were: nausea (57%), fatigue (including asthenia) (42%), anemia (24%), vomiting (23%), headache (20%), diarrhea (18%), leukopenia (17%), neutropenia (16%), decreased appetite (13%), dysgeusia (12%), dizziness (11%), and stomatitis (10%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients who received LYNPARZA in the **adjuvant setting for OlympiA** were: decrease in lymphocytes (77%), increase in mean corpuscular volume (67%), decrease in hemoglobin (65%), decrease in leukocytes (64%), and decrease in absolute neutrophil count (39%).

### ADVERSE REACTIONS—gBRCAm, HER2-Negative Metastatic Breast Cancer

Most common adverse reactions (Grades 1-4) in  $\geq 20\%$  of patients who received LYNPARZA in the **metastatic setting for OlympiAD** were: nausea (58%), anemia (40%), fatigue (including asthenia) (37%), vomiting (30%), neutropenia (27%), respiratory tract infection (27%), leukopenia (25%), diarrhea (21%), and headache (20%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients who received LYNPARZA in the **metastatic setting for OlympiAD** were: decrease in hemoglobin (82%), decrease in lymphocytes (73%), decrease in leukocytes (71%), increase in mean corpuscular volume (71%), decrease in absolute neutrophil count (46%), and decrease in platelets (33%).

### ADVERSE REACTIONS—First-Line Maintenance gBRCAm Metastatic Pancreatic Adenocarcinoma

Most common adverse reactions (Grades 1-4) in  $\geq 10\%$  of patients who received LYNPARZA in the **first-line maintenance setting for POLO** were: fatigue (60%), nausea (45%), abdominal pain (34%), diarrhea (29%), anemia (27%), decreased appetite (25%), constipation (23%), vomiting (20%), back pain (19%), arthralgia (15%), rash (15%), thrombocytopenia (14%), dyspnea (13%), neutropenia (12%), nasopharyngitis (12%), dysgeusia (11%), and stomatitis (10%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients who received LYNPARZA in the **first-line maintenance setting for POLO** were: increase in serum creatinine (99%), decrease in hemoglobin (86%), increase in mean corpuscular volume (71%), decrease in lymphocytes (61%), decrease in platelets (56%), decrease in leukocytes (50%), and decrease in absolute neutrophil count (25%).

### ADVERSE REACTIONS—HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer

Most common adverse reactions (Grades 1-4) in  $\geq 10\%$  of patients who received LYNPARZA for **PROfound** were: anemia (46%), fatigue (including asthenia) (41%), nausea (41%), decreased appetite (30%), diarrhea (21%), vomiting (18%), thrombocytopenia (12%), cough (11%), and dyspnea (10%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 25\%$  of patients who received LYNPARZA for **PROfound** were: decrease in hemoglobin (98%), decrease in lymphocytes (62%), decrease in leukocytes (53%), and decrease in absolute neutrophil count (34%).

### ADVERSE REACTIONS—Metastatic Castration-Resistant Prostate Cancer in Combination with Abiraterone and Prednisone or Prednisolone

Most common adverse reactions (Grades 1-4) in  $\geq 10\%$  of patients who received LYNPARZA/abiraterone with a difference of  $\geq 5\%$  compared to placebo for **PROpel** were: anemia (48%), fatigue (including asthenia) (38%), nausea (30%), diarrhea (19%), decreased appetite (16%), lymphopenia (14%), dizziness (14%), and abdominal pain (13%).

Most common laboratory abnormalities (Grades 1-4) in  $\geq 20\%$  of patients who received LYNPARZA/abiraterone for **PROpel** were: decrease in hemoglobin (97%), decrease in lymphocytes (70%), decrease in platelets (23%), and decrease in absolute neutrophil count (23%).

## DRUG INTERACTIONS

**Anticancer Agents:** Clinical studies of LYNPARZA with other myelosuppressive anticancer agents, including DNA-damaging agents, indicate a potentiation and prolongation of myelosuppressive toxicity.

**CYP3A Inhibitors:** Avoid coadministration of strong or moderate CYP3A inhibitors when using LYNPARZA. If a strong or moderate CYP3A inhibitor must be coadministered, reduce the dose of LYNPARZA. Advise patients to avoid grapefruit, grapefruit juice, Seville oranges, and Seville orange juice during LYNPARZA treatment.

**CYP3A Inducers:** Avoid coadministration of strong or moderate CYP3A inducers when using LYNPARZA.

## USE IN SPECIFIC POPULATIONS

**Lactation:** No data are available regarding the presence of olaparib in human milk, its effects on the breastfed infant or on milk production. Because of the potential for serious adverse reactions in the breastfed infant, advise a lactating woman not to breastfeed during treatment with LYNPARZA and for 1 month after receiving the final dose.

**Pediatric Use:** The safety and efficacy of LYNPARZA have not been established in pediatric patients.

**Hepatic Impairment:** No adjustment to the starting dose is required in patients with mild or moderate hepatic impairment (Child-Pugh classification A and B). There are no data in patients with severe hepatic impairment (Child-Pugh classification C).

**Renal Impairment:** No dosage modification is recommended in patients with mild renal impairment (CLcr 51-80 mL/min estimated by Cockcroft-Gault). In patients with moderate renal impairment (CLcr 31-50 mL/min), reduce the dose of LYNPARZA to 200 mg twice daily. There are no data in patients with severe renal impairment or end-stage renal disease (CLcr  $\leq 30$  mL/min).

## INDICATIONS

LYNPARZA is a poly (ADP-ribose) polymerase (PARP) inhibitor indicated:

### First-Line Maintenance BRCAm Advanced Ovarian Cancer

For the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### First-Line Maintenance HRD-Positive Advanced Ovarian Cancer in Combination with Bevacizumab

In combination with bevacizumab for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD)-positive status defined by either:

- a deleterious or suspected deleterious BRCA mutation, and/or
- genomic instability

Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### Maintenance BRCA-mutated Recurrent Ovarian Cancer

For the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, who are in complete or partial response to platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### Adjuvant Treatment of gBRCAm, HER2-Negative, High-Risk Early Breast Cancer

For the adjuvant treatment of adult patients with deleterious or suspected deleterious gBRCAm, human epidermal growth factor receptor 2 (HER2)-negative, high-risk early breast cancer who have been treated with neoadjuvant or adjuvant chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### gBRCAm, HER2-Negative Metastatic Breast Cancer

For the treatment of adult patients with deleterious or suspected deleterious gBRCAm, human epidermal growth factor receptor 2 (HER2)-negative metastatic breast cancer who have been treated with chemotherapy in the neoadjuvant, adjuvant, or metastatic setting. Patients with hormone receptor (HR)-positive breast cancer should have been treated with a prior endocrine therapy or be considered inappropriate for endocrine therapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### First-Line Maintenance gBRCAm Metastatic Pancreatic Cancer

For the maintenance treatment of adult patients with deleterious or suspected deleterious gBRCAm metastatic pancreatic adenocarcinoma whose disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer

For the treatment of adult patients with deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer (mCRPC) who have progressed following prior treatment with enzalutamide or abiraterone. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

### BRCAm Metastatic Castration-Resistant Prostate Cancer in Combination with Abiraterone and Prednisone or Prednisolone

In combination with abiraterone and prednisone or prednisolone (abi/pred) for the treatment of adult patients with deleterious or suspected deleterious BRCA-mutated (BRCAm) metastatic castration-resistant prostate cancer (mCRPC). Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.

Please see accompanying complete Prescribing Information, including Medication Guide or visit [https://www.azpicentral.com/lynparza\\_tb/lynparza\\_tb.pdf](https://www.azpicentral.com/lynparza_tb/lynparza_tb.pdf)

You are encouraged to report negative side effects of AstraZeneca prescription drugs by calling 1-800-236-9933. If you prefer to report these to the FDA, call 1-800-FDA-1088.



LYNPARZA is a registered trademark of the AstraZeneca group of companies.

©2023 AstraZeneca. All rights reserved. US-81846 11/23

