Math 2471 Calculus III – Sample Final Questions

- 1. Find the unit tangent and unit normal vector for $\vec{r}(t) = \langle t, \frac{1}{2}t^2 \rangle$.
- 2. (i) Prove the limits either exist or doesn't exist

(i)
$$\lim_{(x,y) \to (0,0)} \frac{x^2 + 2y^2}{x^2 + y^2}$$
 (ii) $\lim_{(x,y) \to (0,0)} \frac{x^2y^4}{x^2 + y^2}$

3. Find the equation of the tangent plane to the given surface at the specified point

$$x^2y + xz + yz^2 = 3$$
, $P(1, 2, -1)$

- 4. Find the directional derivative of $z = x^2 + 3xy + y^2$ at (1, 1) in the direction of $\langle -3, 4 \rangle$
- 5. Classify the critical points for $z = x^2y x^2 + y^2 18y$.
- 6. Set up and evaluate an integral to calculate
 - (i) the volume bound by z = 0, $z = 1 x^2 y^2$
 - (ii) the volume inside both $x^2 + y^2 + z^2 = 2$, $x^2 + y^2 = 1$
 - (ii) the surface area of x + 2y + 3z = 6, $x, y, z \ge 0$

7. Set of the triple integral $\iiint f(x, y, z) dV$ in both cylindrical and spherical coordinates for the volume inside the cone $z = \sqrt{x^2 + y^2}$ bound by z = 1

8. Is the following vector field conservative?

$$\vec{F} = \langle y^2 + 3yz, 2xy + 3xz, 3xy \rangle$$
.

If so, determine the potential function *f* such that $\vec{F} = \vec{\nabla} f$. Use this to evaluate

$$\int_{c} (y^2 + 3yz) \, dx + (2xy + 3xz) \, dy + 3xy \, dz > .$$

where *c* is a curve that connects the points (0,0,0) and (1,2,3).

9. Evaluate the following line integrals:

- (*i*) $\int_{c} 2xy \, dx + (x+1) \, dy$ where c is the ccw direction around the square with vertices (0,0), (1,0), (1,1) and (0,1)
- (*ii*) $\int_{c} (x y) dx + (x + y) dy$ where c is clockwise direction around a circle of radius 2.

10. Verify Green's Theorem

$$\oint_{c} Pdx + Qdy = \iint_{R} \left(Q_{x} - P_{y} \right) dA$$

where $\vec{F} = \langle 3x^2y, x^3 + x \rangle$ and *c* the curve enclosing the area given by y = x and $y = x^2$.

11. Evaluate the following surface integrals

- $\iint_{S} xy \, dS \text{ where } S \text{ is the portion of the plane } 2x + y + z = 6 \text{ in the first octant.}$ $\iint_{S} (x + z) \, dS \text{ where } S \text{ is the portion of } y^2 + z^2 = 9 \text{ in the first octant between}$ *(i)*
- *(ii)* x = 0 and x = 4.
- 12. Verify the divergence theorem

$$\iint\limits_{S} \vec{F} \cdot \vec{N} \, dS = \iiint\limits_{V} \nabla \cdot \vec{F} \, dV$$

where $\vec{F} = \langle x + yz, y + xz, z + xy \rangle$ and *V* is the volume of the tetrahedron bound by x + y + z = 1 and the planes x = 0, y = 0 and z = 0.