



Supplemental Labeling

ECHO® 720
Agricultural Fungicide

ACCEPTED
 FOR REGISTRATION

3/29/11

New York State Department
 of Environmental Conservation
 Division of Materials Management
 Pesticide Product Registration

EPA Reg. No. 60063-7

Supplemental Label for brassica (head and stem), cucurbits, fruiting vegetables, horseradish, lupine and lentil, persimmon, rhubarb and yam.

Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile).....	54.0%
Other Ingredients:	46.0%
Total:	100.0%

Contains 6.0 Pounds Chlorothalonil Per Gallon (720 grams per liter)

Keep Out of Reach of Children

WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Manufactured for
 Sipcam Agro USA, Inc.
 2520 Meridian Parkway, Suite 525
 Durham, NC 27713

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner that is inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label for ECHO 720.

Application Rates

Dosage rates on this label indicate pints of ECHO 720 per acre, unless otherwise stated. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

For each listed crop, the maximum total amount of chlorothalonil active ingredient (lbs a.i./A) which may be applied per acre of that crop (or crop group) during each growing season is given in bold print within a box beneath the crop name. For each crop use situation listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased.

CROPS

CROP	PHI (DAYS)	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Brassica, Head and Stem: Cabbage Chinese (napa), Cabbage Chinese Mustard Cavalo, Broccolo, Kohlrabi 8.8 lbs a.i./A	7	Alternaria leaf spot, Downy mildew	1.5 pints	Start applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals to maintain control. Apply by air, ground or chemigation
		Ring spot (California only)	2 pints	Start applications when conditions favor disease development. Repeat at 7 to 10 day intervals to maintain control. Apply by air, ground or chemigation

<p>Cucurbits:</p> <p>Cucumber, Cantaloupe, Muskmelon, Honeydew melon, Watermelon, Squash, Pumpkin, Zucchini</p> <p>Including cultivars and/or hybrids of these.</p> <p>15.75 lbs a.i./A</p> <p>Additional crops: Chayote, Chinese waxgourd Gourds Momordica spp. (Bitter melon, Balsam apple)</p>	<p>0</p>	<p>Anthracnose, Downy mildew, Target spot</p>	<p>1.5 to 2 pints</p>	<p>Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals.</p> <p>Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply ECHO to watermelons when any of the following conditions are present:</p> <ol style="list-style-type: none"> 1. Intense heat and sunlight; 2. Drought conditions; 3. Poor vine canopy; 4. Other crop and environmental conditions which may be conducive to increased natural sunburn. <p>DO NOT combine ECHO with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p> <p>Apply by ground, air or chemigation.</p>
		<p>Cercospora leaf spot, Gummy stem blight (black rot), Alternaria leaf blight, Alternaria leaf spot, Scab, Powdery mildew (Sphaerotheca only)</p>	<p>2 to 3 pints</p>	

<p>Fruiting Vegetables: (Does not include tomatoes)</p> <p>Eggplant, Groundcherry, Okra, Pepino, Pepper (includes, bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo</p> <p>9.0 lbs a.i./A</p>	3	<p>Anthrachnose, Botrytis leaf mold, Cercospora leaf spot, Powdery mildew</p>	1.5 pints	<p>Begin applications as a foliage, flower and fruit spray when disease is anticipated. Repeat applications at 7-10 day intervals.</p> <p>Apply by ground, air or chemigation</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---	-------------------------------------------------------------------------------------------	-----------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Horseradish 18.0 lbs a.i./A	14	Ramularia stem and leafspot	3 pints	Start applications when disease threatens and repeat at 7 to 10 day intervals as necessary to maintain control. Use in sufficient water to obtain adequate coverage.
Lupine, Lentil 6.0 lbs a.i./A	14	Anthrachnose, Ascochyta blight	1 – 1.5 pints	Start applications when disease threatens and repeat at 7 to 10 day intervals as necessary to maintain control. Use in sufficient water to obtain adequate coverage.
Persimmon 4.7 lbs a.i./A	14	Cercospora leaf spot	1.25 pints	Start applications when disease first threatens and repeat at 14 day intervals as necessary to maintain control. Apply in the States of Florida and Hawaii only. Use in sufficient water to obtain adequate coverage. Aerial application requires the use of a minimum of 10 gallons per acre.
Rhubarb 13.5 lbs a.i./A	30	Ramularia leaf spot Ascochyta blight	3 pints	Start applications when disease threatens and repeat at 7 to 10 day intervals or as necessary to maintain control. Use in sufficient water to obtain adequate coverage.
Yam 11.25 lbs a.i./A	7	Anthrachnose	1 – 1.25 pints	Start applications when disease threatens and repeat at 10 -14 day intervals as necessary to maintain control. Use in sufficient water to obtain adequate coverage.

Supplemental label: 5/17/10