

Case-Mix Adjustment and Variable Scores

CLINICAL DIMENSION					
	Episode number within sequence of adjacent episodes	1 or 2	1 or 2	3+	3+
	Therapy Visits	0-13	14+	0-13	14+
	EQUATION	1	2	3	4
1	Primary or Other Diagnosis = Blindness/Low Vision				
2	Primary or Other Diagnosis = Blood disorders		6		3
3	Primary or Other Diagnosis = Cancer, selected benign neoplasms		8		8
4	Primary Diagnosis = Diabetes		8		7
5	Other Diagnosis = Diabetes	1			
6	Primary or Other Diagnosis = Dysphagia AND Primary or Other Diagnosis = Neuro 3 -Stroke	2	16	1	9
7	Primary or Other Diagnosis = Dysphagia AND M1030 (Therapy at home) = 3 (Enteral)	2	7		7
8	Primary or Other Diagnosis = Gastrointestinal disorders				
9	Primary or Other Diagnosis = Gastrointestinal disorders AND M1630 (ostomy)= 1 or 2		6		
10	Primary or Other Diagnosis = Gastrointestinal disorders AND Primary or Other Diagnosis = Neuro 1 - Brain disorders and paralysis, OR Neuro 2 - Peripheral neurological disorders, OR Neuro 3 - Stroke, OR Neuro 4- Multiple Sclerosis				
11	Primary or Other Diagnosis = Heart Disease OR Hypertension	1			
12	Primary Diagnosis = Neuro 1 - Brain disorders and paralysis	3	11	6	11
13	Primary or Other Diagnosis = Neuro 1 - Brain disorders and paralysis AND M1840 (Toilet transfer)= 2 or more				
14	Primary or Other Diagnosis = Neuro 1 - Brain disorders and paralysis OR Neuro 2 -Peripheral neurological disorders AND M1810 or M1820 (Dressing upper or lower body)=1, 2, or 3	2	7	1	7
15	Primary or Other Diagnosis = Neuro 3 - Stroke	3	10	2	
16	Primary or Other Diagnosis = Neuro 3 – Stroke AND M1810 or M1820 (Dressing upper or lower body)=1, 2, or 3		4		8
17	Primary or Other Diagnosis = Neuro 3 – Stroke AND M1860 (Ambulation) = 4 or more				
18	Primary or Other Diagnosis = Neuro 4 -Multiple Sclerosis AND AT LEAST ONE OF THE FOLLOWING: M1830 (Bathing)= 2 or more, OR M1840 (Toilet transfer)= 2 or more OR M1850 (Transferring)= 2 or more OR M1860 (Ambulation) = 4 or more	3	8	7	13
19	Primary or Other Diagnosis = Ortho 1 – Leg Disorders or Gait Disorders AND M1324 (most problematic pressure ulcer stage)= 1, 2, 3 or 4	8	1	8	4
20	Primary or Other Diagnosis = Ortho 1 -Leg OR Ortho 2 – Other orthopedic disorders AND M1030 (Therapy at home)= 1 (IV/Infusion) or 2 Parenteral)	4	3	2	
21	Primary or Other Diagnosis = Psych 1 -Affective and other psychoses, depression				
22	Primary or Other Diagnosis = Psych 2 - Degenerative and other organic psychiatric disorders				
23	Primary or Other Diagnosis = Pulmonary disorders				
24	Primary or Other Diagnosis = Pulmonary disorders AND M1860 (Ambulation) = 1 or more				
25	Primary Diagnosis = Skin 1 -Traumatic wounds, burns, and post-operative complications	4	21	8	19
26	Other Diagnosis = Skin 1 - Traumatic wounds, burns, post-operative complications	6	15	7	15
27	Primary or Other Diagnosis = Skin 1 -Traumatic wounds, burns, and post-operative complications OR Skin 2–Ulcers and other skin conditions AND M1030 (Therapy at home) = 1 (IV/Infusion) or 2 (Parenteral)	4		1	

Case-Mix Adjustment and Variable Scores

28	Primary or Other Diagnosis = Skin 2 - Ulcers and other skin conditions	2	17	8	17
29	Primary or Other Diagnosis = Tracheostomy	4	19	4	11
30	Primary or Other Diagnosis = Urostomy/Cystostomy		19		14
31	M1030 (Therapy at home) = 1 (IV/Infusion) or 2 (Parenteral)		18	6	18
32	M1030 (Therapy at home) = 3 (Enteral)		15		7
33	M1200 (Vision)= 1 or more				
34	M1242 (Pain)= 3 or 4	2		1	
35	M1308 =Two or more pressure ulcers at stage 3 or 4	4	5	4	13
36	M1324 (Most problematic pressure ulcer stage)= 1 or 2	3	19	7	16
37	M1324 (Most problematic pressure ulcer stage)= 3 or 4	8	33	12	26
38	M1334 (Stasis ulcer status)= 2	4	13	8	22
39	M1334 (Stasis ulcer status)= 3	7	18	10	18
40	M1342 (Surgical wound status)= 2	1	7	6	14
41	M1342 (Surgical wound status)= 3		6	5	11
42	M1400 (Dyspnea)= 2, 3, or 4		2		3
43	M1620 (Bowel Incontinence)= 2 to 5		4		3
44	M1630 (Ostomy)= 1 or 2	4	11	3	11
45	M2030 (Injectable Drug Use)= 0, 1, 2, or 3				
FUNCTIONAL DIMENSION					
46	M1810 or M1820 (Dressing upper or lower body)=1, 2, or 3	2		2	
47	M1830 (Bathing)= 2 or more	6	3	5	
48	M1840 (Toilet transferring)= 2 or more	1	3		3
49	M1850 (Transferring)= 2 or more	3	4	2	1
50	M1860 (Ambulation) = 1, 2 or 3	7		3	
51	M1860 (Ambulation) = 4 or more	7	8	6	8

CY 2015 Clinical and Functional Thresholds		1 st and 2 nd Episodes		3 rd + episodes		All episodes
		0-13 therapy visits	14-19 therapy visits	01-13 therapy visits	14-19 therapy visits	20+ therapy visits
Grouping Step:		1	2	3	4	5
Equation(s) used to calculate points: (see table 8)		1	2	3	4	(2&4)
Dimension	Severity Level					
Clinical	C1	0 to 1	0 to 1	0	0 to 5	0 to 3
	C2	2 to 3	2 to 7	1	6 to 12	4 to 16
	C3	4+	8+	2+	13+	17+
Functional	F1	0 to 14	0 to 3	0 to 9	0	0 to 2
	F2	15	4 to 13	10	1 to 7	3 to 5
	F3	16+	14+	11+	8+	6+
Service	S1	0-5	14-15	0-5	14-15	20+
	S2	6	16-17	6	16-17	
	S3	7-9	18-19	7-9	18-19	
	S4	10		10		
	S5	11-13		11-13		