

REVIEW ARTICLE

Dan L. Longo, M.D., *Editor*

Adverse Health Effects of Marijuana Use

Nora D. Volkow, M.D., Ruben D. Baler, Ph.D., Wilson M. Compton, M.D.,
and Susan R.B. Weiss, Ph.D.

IN LIGHT OF THE RAPIDLY SHIFTING LANDSCAPE REGARDING THE LEGALIZATION of marijuana for medical and recreational purposes, patients may be more likely to ask physicians about its potential adverse and beneficial effects on health. The popular notion seems to be that marijuana is a harmless pleasure, access to which should not be regulated or considered illegal. Currently, marijuana is the most commonly used “illicit” drug in the United States, with about 12% of people 12 years of age or older reporting use in the past year and particularly high rates of use among young people.¹ The most common route of administration is inhalation. The greenish-gray shredded leaves and flowers of the *Cannabis sativa* plant are smoked (along with stems and seeds) in cigarettes, cigars, pipes, water pipes, or “blunts” (marijuana rolled in the tobacco-leaf wrapper from a cigar). Hashish is a related product created from the resin of marijuana flowers and is usually smoked (by itself or in a mixture with tobacco) but can be ingested orally. Marijuana can also be used to brew tea, and its oil-based extract can be mixed into food products.

The regular use of marijuana during adolescence is of particular concern, since use by this age group is associated with an increased likelihood of deleterious consequences² (Table 1). Although multiple studies have reported detrimental effects, others have not, and the question of whether marijuana is harmful remains the subject of heated debate. Here we review the current state of the science related to the adverse health effects of the recreational use of marijuana, focusing on those areas for which the evidence is strongest.

ADVERSE EFFECTS

RISK OF ADDICTION

Despite some contentious discussions regarding the addictiveness of marijuana, the evidence clearly indicates that long-term marijuana use can lead to addiction. Indeed, approximately 9% of those who experiment with marijuana will become addicted³ (according to the criteria for dependence in the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition [DSM-IV]). The number goes up to about 1 in 6 among those who start using marijuana as teenagers and to 25 to 50% among those who smoke marijuana daily.⁴ According to the 2012 National Survey on Drug Use and Health, an estimated 2.7 million people 12 years of age and older met the DSM-IV criteria for dependence on marijuana, and 5.1 million people met the criteria for dependence on any illicit drug¹ (8.6 million met the criteria for dependence on alcohol¹). There is also recognition of a bona fide cannabis withdrawal syndrome⁵ (with symptoms that include irritability, sleeping difficulties, dysphoria, craving, and anxiety), which makes cessation difficult and contributes to relapse. Marijuana use by adolescents is particularly troublesome. Adolescents' increased vulnerability to adverse long-term outcomes from marijuana use is probably related

From the National Institute on Drug Abuse, National Institutes of Health, Bethesda, MD. Address reprint requests to Dr. Volkow at the National Institute on Drug Abuse, 6001 Executive Blvd., Rm. 5274, Bethesda, MD 20892, or at nvolkow@nida.nih.gov.

N Engl J Med 2014;370:2219-27.

DOI: 10.1056/NEJMra1402309

Copyright © 2014 Massachusetts Medical Society.

Table 1. Adverse Effects of Short-Term Use and Long-Term or Heavy Use of Marijuana.**Effects of short-term use**

Impaired short-term memory, making it difficult to learn and to retain information

Impaired motor coordination, interfering with driving skills and increasing the risk of injuries

Altered judgment, increasing the risk of sexual behaviors that facilitate the transmission of sexually transmitted diseases

In high doses, paranoia and psychosis

Effects of long-term or heavy use

Addiction (in about 9% of users overall, 17% of those who begin use in adolescence, and 25 to 50% of those who are daily users)*

Altered brain development*

Poor educational outcome, with increased likelihood of dropping out of school*

Cognitive impairment, with lower IQ among those who were frequent users during adolescence*

Diminished life satisfaction and achievement (determined on the basis of subjective and objective measures as compared with such ratings in the general population)*

Symptoms of chronic bronchitis

Increased risk of chronic psychosis disorders (including schizophrenia) in persons with a predisposition to such disorders

* The effect is strongly associated with initial marijuana use early in adolescence.

to the fact that the brain, including the endocannabinoid system, undergoes active development during adolescence.⁶ Indeed, early and regular marijuana use predicts an increased risk of marijuana addiction, which in turn predicts an increased risk of the use of other illicit drugs.⁷ As compared with persons who begin to use marijuana in adulthood, those who begin in adolescence are approximately 2 to 4 times as likely to have symptoms of cannabis dependence within 2 years after first use.⁸

EFFECT ON BRAIN DEVELOPMENT

The brain remains in a state of active, experience-guided development from the prenatal period through childhood and adolescence until the age of approximately 21 years.⁹ During these developmental periods, it is intrinsically more vulnerable than a mature brain to the adverse long-term effects of environmental insults, such as exposure to tetrahydrocannabinol, or THC, the primary active ingredient in marijuana. This view has received considerable support from studies in animals, which have shown, for example, that prenatal or adolescent exposure to

THC can recalibrate the sensitivity of the reward system to other drugs¹⁰ and that prenatal exposure interferes with cytoskeletal dynamics, which are critical for the establishment of axonal connections between neurons.¹¹

As compared with unexposed controls, adults who smoked marijuana regularly during adolescence have impaired neural connectivity (fewer fibers) in specific brain regions. These include the precuneus, a key node that is involved in functions that require a high degree of integration (e.g., alertness and self-conscious awareness), and the fimbria, an area of the hippocampus that is important in learning and memory.¹² Reduced functional connectivity has also been reported in the prefrontal networks responsible for executive function (including inhibitory control) and the subcortical networks, which process habits and routines.¹³ In addition, imaging studies in persons who use cannabis have revealed decreased activity in prefrontal regions and reduced volumes in the hippocampus.¹⁴ Thus, certain brain regions may be more vulnerable than others to the long-term effects of marijuana. One study showed that selective down-regulation of cannabinoid-1 (CB1) receptors in several cortical brain regions in long-term marijuana smokers was correlated with years of cannabis smoking and was reversible after 4 weeks of abstinence.¹⁵ Changes in CB1 receptors were not seen in subcortical regions.

The negative effect of marijuana use on the functional connectivity of the brain is particularly prominent if use starts in adolescence or young adulthood,¹² which may help to explain the finding of an association between frequent use of marijuana from adolescence into adulthood and significant declines in IQ.¹⁶ The impairments in brain connectivity associated with exposure to marijuana in adolescence are consistent with preclinical findings indicating that the cannabinoid system plays a prominent role in synapse formation during brain development.¹⁷

POSSIBLE ROLE AS GATEWAY DRUG

Epidemiologic and preclinical data suggest that the use of marijuana in adolescence could influence multiple addictive behaviors in adulthood. In rodents exposed to cannabinoids during adolescence, there is decreased reactivity of the dopamine neurons that modulate the brain's reward regions.¹⁸ The exposure of rodents to

cannabis in utero alters the developmental regulation of the mesolimbic dopamine system of affected offspring.¹⁹ If reduced dopamine reactivity in the brain's reward regions does follow early exposure to marijuana, this effect could help to explain the increased susceptibility to drug abuse and addiction to several drugs later in life, which has been reported in most epidemiologic studies.²⁰ This theory is also consistent with animal models showing that THC can prime the brain for enhanced responses to other drugs.²¹ Although these findings support the idea that marijuana is a gateway drug, other drugs, such as alcohol and nicotine, can also be categorized as gateway drugs, since they also prime the brain for a heightened response to other drugs.²² However, an alternative explanation is that people who are more susceptible to drug-taking behavior are simply more likely to start with marijuana because of its accessibility and that their subsequent social interactions with other drug users would increase the probability that they would try other drugs.

RELATION TO MENTAL ILLNESS

Regular marijuana use is associated with an increased risk of anxiety and depression,²³ but causality has not been established. Marijuana is also linked with psychoses (including those associated with schizophrenia), especially among people with a preexisting genetic vulnerability,²⁴ and exacerbates the course of illness in patients with schizophrenia. Heavier marijuana use, greater drug potency, and exposure at a younger age can all negatively affect the disease trajectory (e.g., by advancing the time of a first psychotic episode by 2 to 6 years).²⁵

However, it is inherently difficult to establish causality in these types of studies because factors other than marijuana use may be directly associated with the risk of mental illness. In addition, other factors could predispose a person to both marijuana use and mental illness. This makes it difficult to confidently attribute the increased risk of mental illness to marijuana use.

EFFECT ON SCHOOL PERFORMANCE AND LIFETIME ACHIEVEMENT

In the 2013 Monitoring the Future survey of high-school students,²⁶ 6.5% of students in grade 12 reported daily or near-daily marijuana use, and this figure probably represents an underesti-

mate of use, since young people who have dropped out of school may have particularly high rates of frequent marijuana use.²⁷ Since marijuana use impairs critical cognitive functions, both during acute intoxication and for days after use,²⁸ many students could be functioning at a cognitive level that is below their natural capability for considerable periods of time. Although acute effects may subside after THC is cleared from the brain, it nonetheless poses serious risks to health that can be expected to accumulate with long-term or heavy use. The evidence suggests that such use results in measurable and long-lasting cognitive impairments,¹⁶ particularly among those who started to use marijuana in early adolescence. Moreover, failure to learn at school, even for short or sporadic periods (a secondary effect of acute intoxication), will interfere with the subsequent capacity to achieve increasingly challenging educational goals, a finding that may also explain the association between regular marijuana use and poor grades.²⁹

The relationship between cannabis use by young people and psychosocial harm is likely to be multifaceted, which may explain the inconsistencies among studies. For example, some studies suggest that long-term deficits may be reversible and remain subtle rather than disabling once a person abstains from use.³⁰ Other studies show that long-term, heavy use of marijuana results in impairments in memory and attention that persist and worsen with increasing years of regular use³¹ and with the initiation of use during adolescence.³² As noted above, early marijuana use is associated with impaired school performance and an increased risk of dropping out of school,^{27,29} although reports of shared environmental factors that influence the risks of using cannabis at a young age and dropping out of school³³ suggest that the relationship may be more complex. Heavy marijuana use has been linked to lower income, greater need for socioeconomic assistance, unemployment, criminal behavior, and lower satisfaction with life.^{2,34}

RISK OF MOTOR-VEHICLE ACCIDENTS

Both immediate exposure and long-term exposure to marijuana impair driving ability; marijuana is the illicit drug most frequently reported in connection with impaired driving and accidents, including fatal accidents.³⁵ There is a relationship between the blood THC concentration

and performance in controlled driving-simulation studies,³⁶ which are a good predictor of real-world driving ability. Recent marijuana smoking and blood THC levels of 2 to 5 ng per milliliter are associated with substantial driving impairment.³⁷ According to a meta-analysis, the overall risk of involvement in an accident increases by a factor of about 2 when a person drives soon after using marijuana.³⁷ In an accident culpability analysis, persons testing positive for THC (typical minimum level of detection, 1 ng per milliliter), and particularly those with higher blood levels, were 3 to 7 times as likely to be responsible for a motor-vehicle accident as persons who had not used drugs or alcohol before driving.³⁸ In comparison, the overall risk of a vehicular accident increases by a factor of almost 5 for drivers with a blood alcohol level above 0.08%, the legal limit in most countries, and increases by a factor of 27 for persons younger than 21 years of age.³⁹ Not surprisingly, the risk associated with the use of alcohol in combination with marijuana appears to be greater than that associated with the use of either drug alone.³⁷

RISK OF CANCER AND OTHER EFFECTS ON HEALTH

The effects of long-term marijuana smoking on the risk of lung cancer are unclear. For example, the use of marijuana for the equivalent of 30 or more joint-years (with 1 joint-year of marijuana use equal to 1 cigarette [joint] of marijuana smoked per day for 1 year) was associated with an increased incidence of lung cancer and several cancers of the upper aerodigestive tract; however, the association disappeared after adjustment for potential confounders such as cigarette smoking.⁴⁰ Although the possibility of a positive association between marijuana smoking and cancer cannot be ruled out,⁴¹ the evidence suggests that the risk is lower with marijuana than with tobacco.⁴⁰ However, the smoking of cigarettes that contain both marijuana and tobacco products is a potential confounding factor with a prevalence that varies dramatically among countries.


Marijuana smoking is also associated with inflammation of the large airways, increased airway resistance, and lung hyperinflation, associations that are consistent with the fact that regular marijuana smokers are more likely to report symptoms of chronic bronchitis than are nonsmokers⁴²; however, the long-term effect of low levels of marijuana exposure does not ap-

pear to be significant.⁴³ The immunologic competence of the respiratory system in marijuana smokers may also be compromised, as indicated by increased rates of respiratory infections and pneumonia.⁴⁴ Marijuana use has also been associated with vascular conditions that increase the risks of myocardial infarction, stroke, and transient ischemic attacks during marijuana intoxication.⁴⁵ The actual mechanisms underlying the effects of marijuana on the cardiovascular and cerebrovascular systems are complex and not fully understood. However, the direct effects of cannabinoids on various target receptors (i.e., CB1 receptors in arterial blood vessels) and the indirect effects on vasoactive compounds⁴⁶ may help explain the detrimental effects of marijuana on vascular resistance and coronary microcirculation.⁴⁷

LIMITATIONS OF THE EVIDENCE AND GAPS IN KNOWLEDGE

Most of the long-term effects of marijuana use that are summarized here have been observed among heavy or long-term users, but multiple (often hidden) confounding factors detract from our ability to establish causality (including the frequent use of marijuana in combination with other drugs). These factors also complicate our ability to assess the true effect of intrauterine exposure to marijuana. Indeed, despite the use of marijuana by pregnant women,⁴⁸ and animal models suggesting that cannabis exposure during pregnancy may alter the normal processes and trajectories of brain development,⁴⁹ our understanding of the long-term effects of prenatal exposure to marijuana in humans is very poor.

The THC content, or potency, of marijuana, as detected in confiscated samples, has been steadily increasing from about 3% in the 1980s to 12% in 2012⁵⁰ (Fig. 1A). This increase in THC content raises concerns that the consequences of marijuana use may be worse now than in the past and may account for the significant increases in emergency department visits by persons reporting marijuana use⁵¹ (Fig. 1B) and the increases in fatal motor-vehicle accidents.³⁵ This increase in THC potency over time also raises questions about the current relevance of the findings in older studies on the effects of marijuana use, especially studies that assessed long-term outcomes.



There is also a need to improve our understanding of how to harness the potential medical benefits of the marijuana plant without exposing people who are sick to its intrinsic risks. The authoritative report by the Institute of Medicine, *Marijuana and Medicine*,⁵² acknowledges the potential benefits of smoking marijuana in stimulating appetite, particularly in patients with the acquired immunodeficiency syndrome (AIDS) and the related wasting syndrome, and in combating chemotherapy-induced nausea and vomiting, severe pain, and some forms of spasticity. The report also indicates that there is some evidence for the benefit of using marijuana

to decrease intraocular pressure in the treatment of glaucoma. Nonetheless, the report stresses the importance of focusing research efforts on the therapeutic potential of synthetically pure cannabinoids.⁵² Some physicians continue to prescribe marijuana for medicinal purposes despite limited evidence of a benefit (see box). This practice raises particular concerns with regard to long-term use by vulnerable populations. For example, there is some evidence to suggest that in patients with symptoms of human immunodeficiency virus (HIV) infection or AIDS, marijuana use may actually exacerbate HIV-associated cognitive deficits.⁷⁵ Simi-

Clinical Conditions with Symptoms That May Be Relieved by Treatment with Marijuana or Other Cannabinoids.***Glaucoma**

Early evidence of the benefits of marijuana in patients with glaucoma (a disease associated with increased pressure in the eye) may be consistent with its ability to effect a transient decrease in intraocular pressure,^{53,54} but other, standard treatments are currently more effective. THC, cannabidiol, and nabilone (a synthetic cannabinoid similar to THC), but not cannabidiol, were shown to lower intraocular pressure in rabbits.^{55,56} More research is needed to establish whether molecules that modulate the endocannabinoid system may not only reduce intraocular pressure but also provide a neuroprotective benefit in patients with glaucoma.⁵⁷

Nausea

Treatment of the nausea and vomiting associated with chemotherapy was one of the first medical uses of THC and other cannabinoids.⁵⁸ THC is an effective antiemetic agent in patients undergoing chemotherapy,⁵⁹ but patients often state that marijuana is more effective in suppressing nausea. Other, unidentified compounds in marijuana may enhance the effect of THC (as appears to be the case with THC and cannabidiol, which operate through different antiemetic mechanisms).⁶⁰ Paradoxically, increased vomiting (hyperemesis) has been reported with repeated marijuana use.

AIDS-associated anorexia and wasting syndrome

Reports have indicated that smoked or ingested cannabis improves appetite and leads to weight gain and improved mood and quality of life among patients with AIDS.⁶¹ However, there is no long-term or rigorous evidence of a sustained effect of cannabis on AIDS-related morbidity and mortality, with an acceptable safety profile, that would justify its incorporation into current clinical practice for patients who are receiving effective antiretroviral therapy.⁶² Data from the few studies that have explored the potential therapeutic value of cannabinoids for this patient population are inconclusive.⁶²

Chronic pain

Marijuana has been used to relieve pain for centuries. Studies have shown that cannabinoids acting through central CB1 receptors, and possibly peripheral CB1 and CB2 receptors,⁶³ play important roles in modeling nociceptive responses in various models of pain. These findings are consistent with reports that marijuana may be effective in ameliorating neuropathic pain,^{64,65} even at very low levels of THC (1.29%).⁶⁶ Both marijuana and dronabinol, a pharmaceutical formulation of THC, decrease pain, but dronabinol may lead to longer-lasting reductions in pain sensitivity and lower ratings of rewarding effects.⁶⁷

Inflammation

Cannabinoids (e.g., THC and cannabidiol) have substantial antiinflammatory effects because of their ability to induce apoptosis, inhibit cell proliferation, and suppress cytokine production.⁶⁸ Cannabidiol has attracted particular interest as an antiinflammatory agent because of its lack of psychoactive effects.⁵⁸ Animal models have shown that cannabidiol is a promising candidate for the treatment of rheumatoid arthritis⁶⁸ and for inflammatory diseases of the gastrointestinal tract (e.g., ulcerative colitis and Crohn's disease).⁶⁹

Multiple sclerosis

Nabiximols (Sativex, GW Pharmaceuticals), an oromucosal spray that delivers a mix of THC and cannabidiol, appears to be an effective treatment for neuropathic pain, disturbed sleep, and spasticity in patients with multiple sclerosis. Sativex is available in the United Kingdom, Canada, and several other countries^{70,71} and is currently being reviewed in phase 3 trials in the United States in order to gain approval from the Food and Drug Administration.

Epilepsy

In a recent small survey of parents who use marijuana with a high cannabidiol content to treat epileptic seizures in their children,⁷² 11% (2 families out of the 19 that met the inclusion criteria) reported complete freedom from seizures, 42% (8 families) reported a reduction of more than 80% in seizure frequency, and 32% (6 families) reported a reduction of 25 to 60% in seizure frequency. Although such reports are promising, insufficient safety and efficacy data are available on the use of cannabis botanicals for the treatment of epilepsy.⁷³ However, there is increasing evidence of the role of cannabidiol as an antiepileptic agent in animal models.⁷⁴

* AIDS denotes acquired immunodeficiency syndrome, CB1 cannabinoid-1 receptor, and CB2 cannabinoid-2 receptor, HIV human immunodeficiency virus, and THC tetrahydrocannabinol.


larly, more research is needed to understand the potential effects of marijuana use on age-related cognitive decline in general and on memory impairment in particular.

Research is needed on the ways in which government policies on marijuana affect public health outcomes. Our understanding of the ef-

fects of policy on market forces is quite limited (e.g., the allure of new tax-revenue streams from the legal sale of marijuana, pricing wars, youth-targeted advertising, and the emergence of cannabis-based medicines approved by the Food and Drug Administration), as is our understanding of the interrelated variables of perceptions about

Figure 2. Use of Marijuana in Relation to Perceived Risk and Daily Use of Tobacco Cigarettes or Marijuana among U.S. Students in Grade 12, 1975–2013.

Panel A shows the inverse correlation between the perception of the risk associated with marijuana use and actual use. Perceived risk corresponds to the percentage of teenagers who reported that the use of marijuana is dangerous. Panel B shows the percentage of students who reported daily use of tobacco cigarettes or marijuana in the previous 30 days. Data for both graphs are from Johnston et al.²⁶



use, types of use, and outcomes. Historically, there has been an inverse correlation between marijuana use and the perception of its risks among adolescents (Fig. 2A). Assuming that this inverse relationship is causal, would greater permissiveness in culture and social policy lead to an increase in the number of young people who are exposed to cannabis on a regular basis? Among students in grade 12, the reported prevalence of regular marijuana smoking has been steadily increasing in recent years and may soon intersect the trend line for regular tobacco smoking (Fig. 2B). We also need information about the effects of second-hand exposure to cannabis smoke and cannabinoids. Second-hand exposure is an important public health issue in the context of tobacco smoking, but we do not have a clear understanding of the effects of second-hand exposure to marijuana smoking.⁷⁶ Studies in states (e.g., Colorado, California, and Washington) and countries (e.g., Uruguay, Portugal, and the Netherlands) where social and legal policies are shifting may provide important data for shaping future policies.

CONCLUSIONS

Marijuana use has been associated with substantial adverse effects, some of which have been determined with a high level of confidence (Table 2). Marijuana, like other drugs of abuse, can result in addiction. During intoxication, marijuana can interfere with cognitive function (e.g., memory and perception of time) and motor function (e.g., coordination), and these effects can have detrimental consequences (e.g., motor-vehicle accidents). Repeated marijuana use during adolescence may result in long-lasting changes in brain function that can jeopardize educational, professional, and social achievements. However, the ef-

Table 2. Level of Confidence in the Evidence for Adverse Effects of Marijuana on Health and Well-Being.

| Effect | Overall Level of Confidence* |
|---------------------------------------------|------------------------------|
| Addiction to marijuana and other substances | High |
| Abnormal brain development | Medium |
| Progression to use of other drugs | Medium |
| Schizophrenia | Medium |
| Depression or anxiety | Medium |
| Diminished lifetime achievement | High |
| Motor vehicle accidents | High |
| Symptoms of chronic bronchitis | High |
| Lung cancer | Low |

* The indicated overall level of confidence in the association between marijuana use and the listed effects represents an attempt to rank the strength of the current evidence, especially with regard to heavy or long-term use and use that starts in adolescence.

fects of a drug (legal or illegal) on individual health are determined not only by its pharmacologic properties but also by its availability and social acceptability. In this respect, legal drugs

(alcohol and tobacco) offer a sobering perspective, accounting for the greatest burden of disease associated with drugs⁷⁷ not because they are more dangerous than illegal drugs but because their legal status allows for more widespread exposure. As policy shifts toward legalization of marijuana, it is reasonable and probably prudent

to hypothesize that its use will increase and that, by extension, so will the number of persons for whom there will be negative health consequences.

No potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

REFERENCES

- Center for Behavioral Health Statistics and Quality. National survey on drug use and health. Rockville, MD: Substance Abuse & Mental Health Services Administration, 2013.
- Fergusson DM, Boden JM. Cannabis use and later life outcomes. *Addiction* 2008;103:969-76.
- Lopez-Quintero C, Pérez de los Cobos J, Hasin DS, et al. Probability and predictors of transition from first use to dependence on nicotine, alcohol, cannabis, and cocaine: results of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Drug Alcohol Depend* 2011;115:120-30.
- Hall W, Degenhardt L. Adverse health effects of non-medical cannabis use. *Lancet* 2009;374:1383-91.
- Gorelick DA, Levin KH, Copersino ML, et al. Diagnostic criteria for cannabis withdrawal syndrome. *Drug Alcohol Depend* 2012;123:141-7.
- Mechoulam R, Parker LA. The endocannabinoid system and the brain. *Annu Rev Psychol* 2013;64:21-47.
- Hall W, Degenhardt L. Prevalence and correlates of cannabis use in developed and developing countries. *Curr Opin Psychiatry* 2007;20:393-7.
- Chen CY, Storr CL, Anthony JC. Early-onset drug use and risk for drug dependence problems. *Addict Behav* 2009;34:319-22.
- Gogtay N, Giedd JN, Lusk L, et al. Dynamic mapping of human cortical development during childhood through early adulthood. *Proc Natl Acad Sci U S A* 2004;101:8174-9.
- Dinieri JA, Hurd YL. Rat models of prenatal and adolescent cannabis exposure. *Methods Mol Biol* 2012;829:231-42.
- Tortoriello G, Morris CV, Alpar A, et al. Miswiring the brain: Δ9-tetrahydrocannabinol disrupts cortical development by inducing an SCG10/stathmin-2 degradation pathway. *EMBO J* 2014;33:668-85.
- Zalesky A, Solowij N, Yücel M, et al. Effect of long-term cannabis use on axonal fibre connectivity. *Brain* 2012;135:2245-55.
- Filbey F, Yezhuvath U. Functional connectivity in inhibitory control networks and severity of cannabis use disorder. *Am J Drug Alcohol Abuse* 2013;39:382-91.
- Batalla A, Bhattacharyya S, Yücel M, et al. Structural and functional imaging studies in chronic cannabis users: a systematic review of adolescent and adult findings. *PLoS One* 2013;8(2):e55821.
- Hirvonen J, Goodwin RS, Li C-T, et al. Reversible and regionally selective down-regulation of brain cannabinoid CB1 receptors in chronic daily cannabis smokers. *Mol Psychiatry* 2012;17:642-9.
- Meier MH, Caspi A, Ambler A, et al. Persistent cannabis users show neuropsychological decline from childhood to mid-life. *Proc Natl Acad Sci U S A* 2012;109(40):E2657-E2564.
- Gaffuri AL, Ladarre D, Lenkei Z. Type-1 cannabinoid receptor signaling in neuronal development. *Pharmacology* 2012;90:19-39.
- Pistis M, Perra S, Pillolla G, Melis M, Muntoni AL, Gessa GL. Adolescent exposure to cannabinoids induces long-lasting changes in the response to drugs of abuse of rat midbrain dopamine neurons. *Biol Psychiatry* 2004;56:86-94.
- DiNieri JA, Wang X, Szutorisz H, et al. Maternal cannabis use alters ventral striatal dopamine D2 gene regulation in the offspring. *Biol Psychiatry* 2011;70:763-9.
- Agrawal A, Neale MC, Prescott CA, Kendler KS. A twin study of early cannabis use and subsequent use and abuse/dependence of other illicit drugs. *Psychol Med* 2004;34:1227-37.
- Panlilio LV, Zanettini C, Barnes C, Solinas M, Goldberg SR. Prior exposure to THC increases the addictive effects of nicotine in rats. *Neuropsychopharmacology* 2013;38:1198-208.
- Levine A, Huang Y, Drisaldi B, et al. Molecular mechanism for a gateway drug: epigenetic changes initiated by nicotine prime gene expression by cocaine. *Sci Transl Med* 2011;3:107ra109.
- Patton GC, Coffey C, Carlin JB, Degenhardt L, Lynskey M, Hall W. Cannabis use and mental health in young people: cohort study. *BMJ* 2002;325:1195-8.
- Caspi A, Moffitt TE, Cannon M, et al. Moderation of the effect of adolescent-onset cannabis use on adult psychosis by a functional polymorphism in the catechol-O-methyltransferase gene: longitudinal evidence of a gene X environment interaction. *Biol Psychiatry* 2005;57:1117-27.
- Di Forti M, Sallis H, Allegrì F, et al. Daily use, especially of high-potency cannabis, drives the earlier onset of psychosis in cannabis users. *Schizophr Bull* 2014 March 19 (Epub ahead of print).
- Johnston LD, O'Malley PM, Miech RA, et al. Monitoring the Future: national survey results on drug use, 1975-2013 — overview, key findings on adolescent drug use. Ann Arbor: Institute for Social Research, University of Michigan, 2014 (<http://monitoringthefuture.org/pubs/monographs/mtf-ove-view2013.pdf>).
- Bray JW, Zarkin GA, Ringwalt C, Qi J. The relationship between marijuana initiation and dropping out of high school. *Health Econ* 2000;9:9-18.
- Crean RD, Crane NA, Mason BJ. An evidence based review of acute and long-term effects of cannabis use on executive cognitive functions. *J Addict Med* 2011;5:1-8.
- Lynskey M, Hall W. The effects of adolescent cannabis use on educational attainment: a review. *Addiction* 2000;95:1621-30.
- Macleod J, Oakes R, Copello A, et al. Psychological and social sequelae of cannabis and other illicit drug use by young people: a systematic review of longitudinal, general population studies. *Lancet* 2004;363:1579-88.
- Solowij N, Stephens RS, Roffman RA, et al. Cognitive functioning of long-term heavy cannabis users seeking treatment. *JAMA* 2002;287:1123-31. [Erratum, *JAMA* 2002;287:1651.]
- Schweinsburg AD, Brown SA, Tapert SF. The influence of marijuana use on neurocognitive functioning in adolescents. *Curr Drug Abuse Rev* 2008;1:99-111.
- Verweij KJ, Huizink AC, Agrawal A, Martin NG, Lynskey MT. Is the relationship between early-onset cannabis use and educational attainment causal or due to common liability? *Drug Alcohol Depend* 2013;133:580-6.
- Brook JS, Lee JY, Finch SJ, Seltzer N, Brook DW. Adult work commitment, financial stability, and social environment as related to trajectories of marijuana use beginning in adolescence. *Subst Abuse* 2013;34:298-305.
- Brady JE, Li G. Trends in alcohol and other drugs detected in fatally injured drivers in the United States, 1999-2010. *Am J Epidemiol* 2014;179:692-9.
- Lenné MG, Dietze PM, Triggs TJ,

- Walmsley S, Murphy B, Redman JR. The effects of cannabis and alcohol on simulated arterial driving: influences of driving experience and task demand. *Accid Anal Prev* 2010;42:859-66.
37. Hartman RL, Huestis MA. Cannabis effects on driving skills. *Clin Chem* 2013; 59:478-92.
38. Ramaekers JG, Berghaus G, van Laar M, Drummer OH. Dose related risk of motor vehicle crashes after cannabis use. *Drug Alcohol Depend* 2004;73:109-19.
39. Peck RC, Gebers MA, Voas RB, Romano E. The relationship between blood alcohol concentration (BAC), age, and crash risk. *J Safety Res* 2008;39:311-9.
40. Hashibe M, Morgenstern H, Cui Y, et al. Marijuana use and the risk of lung and upper aerodigestive tract cancers: results of a population-based case-control study. *Cancer Epidemiol Biomarkers Prev* 2006; 15:1829-34.
41. Callaghan RC, Allebeck P, Sidorchuk A. Marijuana use and risk of lung cancer: a 40-year cohort study. *Cancer Causes Control* 2013;24:1811-20.
42. Tashkin DP. Effects of marijuana smoking on the lung. *Ann Am Thorac Soc* 2013;10:239-47.
43. Pletcher MJ, Vittinghoff E, Kalhan R, et al. Association between marijuana exposure and pulmonary function over 20 years. *JAMA* 2012;307:173-81.
44. Owen KP, Sutter ME, Albertson TE. Marijuana: respiratory tract effects. *Clin Rev Allergy Immunol* 2014;46:65-81.
45. Thomas G, Kloner RA, Rezkalla S. Adverse cardiovascular, cerebrovascular, and peripheral vascular effects of marijuana inhalation: what cardiologists need to know. *Am J Cardiol* 2014;113:187-90.
46. Stanley C, O'Sullivan SE. Vascular targets for cannabinoids: animal and human studies. *Br J Pharmacol* 2014;171:1361-78.
47. Rezkalla SH, Sharma P, Kloner RA. Coronary no-flow and ventricular tachycardia associated with habitual marijuana use. *Ann Emerg Med* 2003;42:365-9.
48. Brown HL, Graves CR. Smoking and marijuana use in pregnancy. *Clin Obstet Gynecol* 2013;56:107-13.
49. Juras-Aswad D, DiNieri JA, Harkany T, Hurd YL. Neurobiological consequences of maternal cannabis on human fetal development and its neuropsychiatric outcome. *Eur Arch Psychiatry Clin Neurosci* 2009;259:395-412.
50. ElSohly MA. Potency Monitoring Program quarterly report no.123 — reporting period: 09/16/2013-12/15/2013. Oxford: University of Mississippi, National Center for Natural Products Research, 2014.
51. Drug Abuse Warning Network, 2011: national estimates of drug-related emergency department visits. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2011 (<http://www.samhsa.gov/data/2k13/DAWN2k11ED/DAWN2k11ED.htm>).
52. Joy JE, Watson SJ Jr, Benson JA Jr, eds. Marijuana and medicine: assessing the science base. Washington, DC: National Academy Press, 1999.
53. Merritt JC, Crawford WJ, Alexander PC, Anduze AL, Gelbart SS. Effect of marijuana on intraocular and blood pressure in glaucoma. *Ophthalmology* 1980;87:222-8.
54. Hepler RS, Frank IR. Marijuana smoking and intraocular pressure. *JAMA* 1971;217:1392.
55. Chen J, Matias I, Dinh T, et al. Finding of endocannabinoids in human eye tissues: implications for glaucoma. *Biochem Biophys Res Commun* 2005;330:1062-7.
56. Song ZH, Slowey CA. Involvement of cannabinoid receptors in the intraocular pressure-lowering effects of WIN55212-2. *J Pharmacol Exp Ther* 2000;292:136-9.
57. Nucci C, Bari M, Spanò A, et al. Potential roles of (endo) cannabinoids in the treatment of glaucoma: from intraocular pressure control to neuroprotection. *Prog Brain Res* 2008;173:451-64.
58. Zuardi AW. Cannabidiol: from an inactive cannabinoid to a drug with wide spectrum of action. *Rev Bras Psiquiatr* 2008;30:271-80.
59. Sallan SE, Zinberg NE, Frei E III. Antiemetic effect of delta-9-tetrahydrocannabinol in patients receiving cancer chemotherapy. *N Engl J Med* 1975;293:795-7.
60. Parker LA, Kwiatkowska M, Burton P, Mechoulam R. Effect of cannabinoids on lithium-induced vomiting in the *Suncus murinus* (house musk shrew). *Psychopharmacology (Berl)* 2004;171:156-61.
61. D'Souza G, Matson PA, Grady CD, et al. Medicinal and recreational marijuana use among HIV-infected women in the Women's Interagency HIV Study (WIHS) cohort, 1994-2010. *J Acquir Immune Defic Syndr* 2012;61:618-26.
62. Lutge BE, Gray A, Siegfried N. The medical use of cannabis for reducing morbidity and mortality in patients with HIV/AIDS. *Cochrane Database Syst Rev* 2013;4:CD005175.
63. Chiou LC, Hu SS, Ho YC. Targeting the cannabinoid system for pain relief? *Acta Anaesthesiol Taiwan* 2013;51:161-70.
64. Wilsey B, Marcotte T, Tsodikov A, et al. A randomized, placebo-controlled, crossover trial of cannabis cigarettes in neuropathic pain. *J Pain* 2008;9:506-21.
65. Wallace M, Schulteis G, Atkinson JH, et al. Dose-dependent effects of smoked cannabis on capsaicin-induced pain and hyperalgesia in healthy volunteers. *Anesthesiology* 2007;107:785-96.
66. Wilsey B, Marcotte T, Deutsch R, Gouaux B, Sakai S, Donaghe H. Low-dose vaporized cannabis significantly improves neuropathic pain. *J Pain* 2013;14:136-48.
67. Cooper ZD, Comer SD, Haney M. Comparison of the analgesic effects of dronabinol and smoked marijuana in daily marijuana smokers. *Neuropsychopharmacology* 2013;38:1984-92.
68. Nagarkatti P, Pandey R, Rieder SA, Hegde VL, Nagarkatti M. Cannabinoids as novel anti-inflammatory drugs. *Future Med Chem* 2009;1:1333-49.
69. Esposito G, Filippis DD, Cirillo C, et al. Cannabidiol in inflammatory bowel diseases: a brief overview. *Phytother Res* 2013;5:633-6.
70. Collin C, Davies P, Mutiboko IK, Ratcliffe S. Randomized controlled trial of cannabis-based medicine in spasticity caused by multiple sclerosis. *Eur J* 2007; 14:290-6.
71. Centonze D, Mori F, Koch G, et al. Lack of effect of cannabis-based treatment on clinical and laboratory measures in multiple sclerosis. *Neuro Sci* 2009;30:531-4.
72. Porter BE, Jacobson C. Report of a parent survey of cannabidiol-enriched cannabis use in pediatric treatment-resistant epilepsy. *Epilepsy Behav* 2013;29:574-7.
73. Kogan NM, Mechoulam R. Cannabinoids in health and disease. *Dialogues Clin Neurosci* 2007;9:413-30.
74. Hill TD, Cascio MG, Romano B, et al. Cannabidiol-rich cannabis extracts are anticonvulsant in mouse and rat via a CB1 receptor-independent mechanism. *Br J Pharmacol* 2013;170:679-92.
75. Cristiani SA, Pukay-Martin ND, Bornstein RA. Marijuana use and cognitive function in HIV-infected people. *J Neuropsychiatry Clin Neurosci* 2004;16:330-5.
76. Niedbala S, Kardos K, Salamone S, Fritch D, Bronsgeest M, Cone EJ. Passive cannabis smoke exposure and oral fluid testing. *J Anal Toxicol* 2004;28:546-52.
77. Degenhardt L, Hall W. Extent of illicit drug use and dependence, and their contribution to the global burden of disease. *Lancet* 2012;379:55-70.

Copyright © 2014 Massachusetts Medical Society.

IMAGES IN CLINICAL MEDICINE

The *Journal* welcomes consideration of new submissions for Images in Clinical Medicine. Instructions for authors and procedures for submissions can be found on the *Journal's* website at NEJM.org. At the discretion of the editor, images that are accepted for publication may appear in the print version of the *Journal*, the electronic version, or both.

For Immediate Release: Wednesday, June 4, 2014, 5 p.m. EDT

NIDA review summarizes research on marijuana's negative health effects

Comprehensive review published in the New England Journal of Medicine also discusses why risks are greatest for teen users

The current state of science on the adverse health effects of marijuana use links the drug to several significant adverse effects including addiction, a review reports. The article, published today in the *New England Journal of Medicine*, is authored by scientists from the National Institute on Drug Abuse (NIDA), part of the National Institutes of Health.

The review describes the science establishing that marijuana can be addictive and that this risk for addiction increases for daily or young users. It also offers insights into research on the gateway theory indicating that marijuana use, similar to nicotine and alcohol use, may be associated with an increased vulnerability to other drugs.

The authors review literature showing that marijuana impairs driving, increasing the risk of being involved in a car accident and that these risks are further enhanced when combining marijuana with alcohol. The authors also discuss the implications of rising marijuana potencies and note that, because older studies are based on the effects of lower-potency (less THC) marijuana, stronger adverse health effects may occur with today's more potent marijuana. (THC is the psychoactive or mind-altering chemical delta-9-tetrahydrocannabinol found in marijuana.)

The reviewers consider areas in which little research has been conducted. This includes possible health consequences of secondhand marijuana smoke; the long-term impact of prenatal marijuana exposure; the therapeutic potential of the individual chemicals found in the marijuana plant; and effects of marijuana legalization policies on public health.

The scientists focus on marijuana's harmful effects on teens, an age group in which the brain rapidly develops, which is one factor that could help explain increased risks from marijuana use in this population. Research suggests that marijuana impairs critical thinking and memory functions during use and that these deficits persist for days after using. In addition, a long-term study showed that regular marijuana use in the early teen years lowers IQ into adulthood, even if users stopped smoking marijuana as adults.

The NIDA-supported 2013 Monitoring the Future Survey says that 6.5 percent of 12th graders report daily or near-daily marijuana use, with 60 percent perceiving that regular marijuana use can be harmful. "It is important to alert the public that using marijuana in the teen years brings health, social, and academic risk," said lead author and NIDA Director Dr. Nora D. Volkow. "Physicians in particular can play a role in conveying to families that early marijuana use can interfere with crucial social and developmental milestones and can impair cognitive development."

This review emphasizes that marijuana use is likely to increase as state and local policies move toward legalizing marijuana for medical or recreational purposes. As use increases, so might the number of people likely to suffer negative health consequences, the review says.

For more information on marijuana and its health consequences, go to: <http://www.drugabuse.gov/publications/drugfacts/marijuana>.

The National Institute on Drug Abuse is a component of the National Institutes of Health, U.S. Department of Health and Human Services. NIDA supports most of the world's research on the health aspects of drug abuse and addiction. The Institute carries out a large variety of programs to inform policy and improve practice. Fact sheets on the health effects of drugs of abuse and information on NIDA research and other activities can be found on the NIDA home page at <http://www.drugabuse.gov>, which is now compatible with your smartphone, iPad or tablet. To order publications in English or Spanish, call NIDA's DrugPubs research dissemination center at 1-877-NIDA-NIH or 240-645-0228 (TDD) or fax or email requests to 240-645-0227 or drugpubs@nida.nih.gov. Online ordering is available at <http://drugpubs.drugabuse.gov>. NIDA's media guide can be found at <http://drugabuse.gov/mediaguide>, and its new easy-to-read website can be found at <http://www.easyread.drugabuse.gov>.

About the National Institutes of Health (NIH): NIH, the nation's medical research agency, includes 27 Institutes and Centers and is a component of the U.S. Department of Health and Human Services. NIH is the primary federal agency conducting and supporting basic, clinical, and translational medical research, and is investigating the causes, treatments, and cures for both common and rare diseases. For more information about NIH and its programs, visit www.nih.gov.

NIH... Turning Discovery Into Health®

Reference

Adverse Health Effects of Marijuana Use, by Nora D. Volkow, M.D., Ruben D. Baler, Ph.D., Wilson M. Compton, M.D., and Susan R.B. Weiss, Ph.D., published online June 4, 2014 in *The New England Journal of Medicine*.

###

Institute/Center

National Institute on Drug Abuse (NIDA)

Contact

NIDA Press Office
301-443-6245

Subscribe

Receive NIH news releases by e-mail

"It is important to alert the public that using marijuana in the teen years brings health, social, and academic risk."

—Dr. Nora D. Volkow
Director, National Institute on Drug Abuse (NIDA)

WETA

PBS.org

Video


Programs


TV Schedules

Shop

Donate

WATCH | SCHEDULE | TOPICS | ABOUT FRONTLINE | SHOP | TEACHER CENTER

 busted: america's war on marijuana

 A Fact Sheet on the Effects of Marijuana This is from the Partnership for a Drug Free America which states all claims have been substantiated by the National Institute on Drug Abuse.

SUPPORT PROVIDED BY



MEMORY/PERCEPTION/BEHAVIOR

1. Attention, memory and learning are impaired among heavy marijuana users, even after users discontinued its use for at least 24 hours. Heavy marijuana use is associated with residual neuropsychological effects even after a day of supervised abstinence from the drug. Heavy users displayed significantly greater impairment than light users on attention/executive functions, as evidenced particularly by greater preservations on card sorting and reduced learning of word lists. These differences remained after controlling for potential confounding variables, such as estimated levels of premorbid cognitive functioning, and for use of alcohol and other substances in the two groups. However, the question remains open as to whether this impairment is due to a residue of drug in the brain, a withdrawal effect from the drug, or a frank neurotoxic effect of the drug. ("The Residual Cognitive Effects of Heavy Marijuana Use in College Students," Pope, HG Jr., Yurgelun-Todd, D., Biological Psychiatry Laboratory, McLean Hospital, Belmont, MA, JAMA February 21, 1996.)

2. Impaired memory for recent events, difficulty concentrating, dreamlike states, impaired motor coordination, impaired driving and other psychomotor skills, slowed reaction time, impaired goal-directed mental activity, and altered peripheral vision are common associated effects. (Adams and Martin 1996; Fehr and Kalant 1983; Hollister 1988a; Institute of Medicine 1982; Tart 1971)

3. A roadside study of reckless drivers who were not impaired by alcohol, showed that 45% of these drivers tested positive for marijuana. (Dr. Dan Brookoff, published in the New England Journal of Medicine)

4. Marijuana smoking affects the brain and leads to impaired short-term memory, perception, judgment and motor skills. (Marijuana Facts: Parents Need to Know, National Institute on Drug Abuse)

5. In a survey of 150 marijuana using students, 59% surveyed report they sometimes forget what a conversation is about before it has ended. 41% report if they read while stoned they remembered less of what they had read hours later. (Dr. Richard Schwartz, Vienna Pediatric Associates in Psychiatric Annals as reported in NIDA Capsules)

Neurobiological Effects

6. Marijuana activates the same pleasure centers in the brain that are targeted by heroin, cocaine and alcohol. (Dr. Gaetano Di Chiara, University of Caligari, Italy)

Cardiovascular Effects

NEXT ON FRONTLINE

Secrets of the Vatican
ENCORE PRESENTATION
July 8th

RECENT STORIES

July 3, 2014 / 10:42 am
What Comes Next
in Ukraine?

July 3, 2014 / 9:21 am
Coming in July on
FRONTLINE

July 2, 2014 / 4:31 pm
A Return to School
Segregation in
America?

JOIN OUR NEWSLETTER

e-mail address

Subscribe »

Like 370,107 people like this. Sign Up to see what your friends like.

7. Physiological effects of marijuana include an alteration of heart rate. Use of marijuana may result in intense anxiety, panic attacks or paranoia. (National Institute of Drug Abuse)

Chronic Effects

Respiratory System Effects

8. The daily use of 1 to 3 marijuana joints appears to produce approximately the same lung damage and potential cancer risk as smoking 5 times as many cigarettes. (UCLA) The study results suggest that the way smokers inhale marijuana, in addition to its chemical composition, increases the adverse physical effects. The same lung cancer risks associated with tobacco also apply to marijuana users, even though they smoke far less. (reported in NIDA Capsules)

9. Benzopyrene is the chemical in tobacco that causes lung cancer. An average marijuana cigarette contains nearly 50% more benzopyrene than a tobacco cigarette. An average marijuana cigarette contains 30 nanograms of this carcinogen compared to 21 nanograms in an average tobacco cigarette (Marijuana and Health, National Academy of Sciences, Institute of Medicine Report, 1982) Benzopyrene suppresses a gene that controls growth of cells. When this gene is damaged the body becomes more susceptible to cancer. This gene is related to half of all human cancers and as many as 70% of lung cancers.

10. Marijuana users may have many of the same respiratory problems that tobacco smokers have, such as chronic bronchitis and inflamed sinuses. ([Marijuana Facts: Parents Need to Know](#), National Institute on Drug Abuse)

11. Marijuana smokers, when compared to non marijuana smokers, have more respiratory illness. (Polen et al. 1993).

12. Marijuana smoke produces airway injury, acute and chronic bronchitis, lung inflammation, and decreased pulmonary defenses against infection. Smoking one marijuana cigarette leads to air deposition of four times as much cancer-causing tar as does tobacco smoke (Dr. D. Tashkin, Western Journal of Medicine)

13. Heavy marijuana use can affect hormones in both males and females. Heavy doses of the drugs may delay the onset of puberty in young men. Marijuana also can have adverse effects on sperm production. Among women, regular marijuana use can disrupt the normal monthly menstrual cycle and inhibit the discharge of eggs from the ovaries. ([Marijuana Facts: Parents Need to know](#), National Institute on Drug Abuse)

Adverse Mental Effects

14. An "amotivational syndrome" can develop in heavy, chronic marijuana users. It is characterized by decreased drive and ambition, shortened attention span, poor judgment, high distractibility, impaired communication skills, and diminished effectiveness in interpersonal situations. (National Institute of Drug Abuse)

15. Adults who smoked marijuana daily believed it helped them function better, improved self-awareness and improved relationships with others. However, researchers found that users were more willing to tolerate problems, suggesting that the drug served as a buffer for those who would rather avoid confronting problems than make changes that might increase their satisfaction with life. The study indicated that these

Sign Up Create an account or **Log In** to see what your friends are doing.

Few Answers on Border Patrol Agents' Use of Force – Criminal Justice - FRONTLINE
425 people recommend this.

Fourteen Years Later, Looking Back at a School Shooting – The Killer at Thurston High - FRONTLINE
500 people recommend this.

Holcer: Big Banks' Clout "Has an Inhibiting Impact" on Prosecutions – The Untouchables - FRONTLINE
1,083 people recommend this.

Never Forget to Lie – FRONTLINE
5,332 people recommend this.

Facebook social plugin

Follow [@frontlinepbs](#) 95.9K followers

FRONTLINE on

OWN FRONTLINE ON DVD TODAY

[SHOP NOW](#)

subjects used marijuana to avoid dealing with their difficulties and the avoidance inevitably made their problems worse. Although users believed the drug enhanced understanding of themselves, it actually served as a barrier against self-awareness. (case studies by research team from Center for Psychosocial Studies in New York.)

16. Marijuana and some of its compounds influence the immune system and affect the body's ability to resist viruses, bacteria, fungi and protozoa, and decreases the body's anti tumor activities. Marijuana has the potential to alter the backup safeguards of the immune system because it affects diverse types of cells in the body. This could compromise the immune system's ability to screen out cancer cells and eliminate infection. (Dr. Guy A. Cabral, Professor, Medical College of Virginia, speaking at NIDA's National Conference on Marijuana Use: Prevention, Treatment and Research.)

"Unfortunately, much of what is known about the human pharmacology of smoked marijuana comes from experiments with plant material containing about 2% THC or less, or occasionally up to 4% THC. In addition, human experiments typically are done in laboratory settings where only one or two smoked doses were administered to relatively young, medically screened, healthy male volunteers well experienced with the effects of marijuana. Females rarely participated in past marijuana research because of prohibitions (now removed) against their inclusion. Thus the clinical pharmacology of single or repeated smoked marijuana doses given to older people or to people with serious diseases has hardly been researched at all in a controlled laboratory or clinic setting. Some of the very few reports of experiments that have included older or sicker people, particularly patients less experienced in using marijuana suggest the profile of adverse effects may differ from healthy student volunteers smoking in a laboratory experiment (Hollister, 1986a, 1988a)

THC administered alone in its pure form is the most thoroughly research cannabinoid. Much of what has been written has been inferred from the results of experiments using only pure THC. Generally, in experiments actually using marijuana, the assumed dose of marijuana was based only on the concentration of THC in the plant material. The amounts of cannabidiol and other cannabinoids in the plant also vary so that pharmacological interactions modifying the effects THC may occur when marijuana is used instead of pure THC.

The result of this research strategy is that a good deal is known about the pharmacology of THC, but experimental confirmation that the pharmacology of a marijuana cigarette is indeed entirely or mainly determined by the amount of THC it contains remains to be completed. The scientific literature contains occasional hints that the pharmacology of pure THC, although similar, is not always the same as the clinical pharmacology of smoked marijuana containing the same amount of THC (Graham 1976, Harvey 1985, Institute of Medicine 1982)" (Report to the Director, National Institutes of Health, by the Ad-Hoc Group of Experts, "Workshop on the Medical Utility of Marijuana.")