

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Stinger®

EPA Reg. No. 62719-73

This supplemental label expires on March 21, 2021 and must not be used or distributed after this date.

Post-Harvest Broadleaf Weed Control in Strawberries

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Stinger according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

Directions for Use

Perennial Strawberries (Not for use in California)

Use Stinger for control of various annual and perennial broadleaf weeds infesting perennial strawberries post-harvest.

Application Rate

Apply one application of Stinger after harvest at 1/3 to 2/3 pint per acre. Apply uniformly with ground equipment in a minimum of 10 gallons of water per acre. For control of Canada thistle from after harvest to early fall apply Stinger after the majority of basal leaves have emerged but prior bud stage.

Specific Use Precautions and Restrictions

- Do not exceed 2/3 pint per acre per year.
- Do not tank mix with other herbicides registered for use on strawberries.
- Do not use surfactants when applying Stinger to strawberry.
- Some leaf cupping may occur. Do not use if unwilling to accept minor leaf injury.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R043-099

EPA accepted: 03/11/19

Initial Printing