


Emergency Care

Fourteenth Edition



Chapter 20

Cardiac Emergencies

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Topics

- [Cardiac Anatomy and Physiology](#)
- [Acute Coronary Syndrome](#)
- [Cardiovascular Disorders](#)

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Cardiac Anatomy and Physiology

[Back to Topics](#)

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Cardiac Anatomy and Physiology (1 of 2)

- Review of the cardiovascular system:
 - Composition of the blood
 - Flow of blood through the chambers of the heart
 - Flow of blood through arteries, veins, arterioles, venules, and capillaries
 - Circulation of blood between heart and lungs, and between heart and the rest of body

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved


Cardiac Anatomy and Physiology (2 of 2)

- Only job of the heart is to pump blood.
 - Generation and distribution of electrical charge
 - Mechanical response to create rhythmic and unceasing pumping action
 - Requires constant supply of oxygen and nutrients

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Coronary Arteries



The coronary arteries.

Pearson

Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Think About It

- How does the normal function of the heart and blood vessels relate to blood pressure and distal pulses?
- How is shock related to the function of the heart and blood vessels?



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Coronary Syndrome

[Back to Topics](#)



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Coronary Syndrome (1 of 4)

- Sometimes called cardiac compromise
- Refers to any time the blood supply to the cells of the heart is blocked or disrupted
- Heart muscle cells go without oxygen, causing cell death.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Coronary Syndrome (2 of 4)

- Hypoxic condition of the heart is called ischemia
- Cells die quickly without adequate blood supply



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Coronary Syndrome (3 of 4)

- Chest discomfort is best-known symptom.
 - Can be described as pressure, squeezing, or aching
- Radiates to the jaw, neck, either arm, or upper abdomen
- Dyspnea is another common finding in older patients and women



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Coronary Syndrome (4 of 4)

- Other symptoms
 - Nausea and/or vomiting
 - Syncope
 - Sudden onset of sweating
 - Abnormal pulse (tachycardia/bradycardia)
 - Abnormal blood pressure
 - Anxiety, feeling of impending doom



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (1 of 8)

- Acute coronary syndrome
 - Perform primary assessment.
 - Explore chief complaint.
 - Use OPQRST to get history of present illness.
 - Obtain past medical history.
 - Take baseline vital signs.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (2 of 8)

- Signs and symptoms of ACS
 - Pain, pressure, or discomfort in the chest, jaw, neck, arms, or upper abdomen
 - Difficulty breathing
 - Palpitations
 - Sudden onset of sweating and nausea or vomiting



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (3 of 8)

- Signs and symptoms of ACS
 - Syncope
 - Anxiety
 - Unusual generalized weakness
 - Abnormal pulse
 - Abnormal blood pressure



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (4 of 8)

- Physical examination
 - Grabbing or clutching the center of the chest
 - Sweating
 - Pale or gray skin
 - Anxiousness or restlessness



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (5 of 8)

- Physical examination
 - Acute pulmonary edema
 - Swollen ankles and feet
 - Medic alert jewelry indicating cardiac problems



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (6 of 8)

- 12-lead ECG
 - Speed up process of recognizing electrocardial findings
 - Interpretation is outside scope of EMT
 - EMT can place leads and acquire print tracing
 - EMT can transmit ECG to destination hospital



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Patient Assessment (7 of 8)

- Preparing a patient for an ECG
 - Place electrodes on chest
 - Electrodes need secure contact with the skin
 - Remove clothing and jewelry
 - Remove sweat and dead skin cells
 - Remove hair



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved


Patient Assessment (8 of 8)

- 12-lead ECG placement
 - Step 1—Place limb leads
 - Step 2—Place V1 and V2 leads
 - Step 3—Place V4 and V3 leads
 - Step 4—Place V5 and V6 leads



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

12-Lead ECG (1 of 2)




Placement of arm and leg leads.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

12-Lead ECG (2 of 2)



Placement of V5 and V6



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (1 of 8)

- Fundamental Principles of Care
 - Place patient in position of comfort.
 - Determine if oxygen should be administered.
 - Respiratory failure, agonal breaths, or apneic should receive high-concentration oxygen via ventilations



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (2 of 8)

- Fundamental Principles of Care
 - Determine if oxygen should be administered.
 - Low oxygen saturations receive high-concentration oxygen via mask or nasal cannula
 - No significant distress and oxygen saturation of at least 94 percent should not receive oxygen



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (3 of 8)

- Fundamental Principles of Care
 - If trained, equipped, and authorized to do so, obtain a 12-lead electrocardiogram (ECG).
 - Follow local protocol as to whether to transmit it to hospital for interpretation.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (4 of 8)

- Fundamental Principles of Care
 - Administer 325 mg of aspirin by mouth if allowed
 - Indications for NOT administering aspirin
 - Risk for aspiration
 - Already taken a full dose
 - Aspirin allergy
 - Recent GI bleeding
 - Taking blood thinner



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Treatment (1 of 2)



Aspirin



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (5 of 8)

- Fundamental Principles of Care
 - Indications for administering nitroglycerin
 - Chest pain
 - History of cardiac problems
 - Physician has prescribed nitroglycerin to patient
 - Patient has nitroglycerin at hand



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Treatment (2 of 2)



Nitroglycerin



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (6 of 8)

- Fundamental Principles of Care
 - Indications for administering nitroglycerin
 - Systolic blood pressure meets protocol criteria
 - Patient has not had Viagra or similar drug for erectile dysfunction within 48 to 72 hours
 - Medical direction authorizes administration of the medication



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (7 of 8)

- Fundamental Principles of Care
 - After giving one dose of nitroglycerin, repeat dose in 5 minutes if:
 - Patient experiences no relief or only partial relief.
 - Systolic blood pressure remains greater than 90 to 100 systolic.
 - Medical direction authorizes another dose of medication.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Care of the Patient with Acute Coronary Syndrome (8 of 8)

- Fundamental Principles of Care
 - Administer maximum of three doses of nitroglycerin
 - Reassess vital signs and chest pain after each dose
 - Treat for shock if blood pressure falls below 90 to 100 systolic
 - Transport promptly



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Cardiovascular Disorders

[Back to Topics](#)



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Cardiovascular Disorders

- Heart problems caused by a number of disorders affecting condition and function of blood vessels and heart



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Coronary Artery Disease (1 of 2)

- Conditions that narrow or block arteries of heart
- Often result from fatty deposit buildup on inner walls of arteries
- Buildup narrows inner vessel diameter, restricting flow of blood.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Coronary Artery Disease (2 of 2)

- Risk factors
 - Heredity
 - Age
 - Hypertension
 - Obesity
 - Lack of exercise
 - Elevated blood cholesterol and triglycerides
 - Cigarette smoking



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Angina Pectoris (1 of 2)

- Chest pain caused by insufficient blood flow to the myocardium
- Typically due to narrowed arteries secondary to coronary artery disease
- Pain usually during times of increased myocardial oxygen demand, such as exertion or stress



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Angina Pectoris (2 of 2)

- Possession of nitroglycerin is good indication that patient has history of angina
- Nitroglycerin dilates blood vessels so heart has less blood to pump



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessing and Treating Angina Pectoris

- If symptoms subside with rest, it is angina pectoris.
- Assume patient is having myocardial infarction until proven otherwise.
- Consider oxygen, administer nitroglycerin and aspirin, obtain 12-lead ECG, transport.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Myocardial Infarction (1 of 6)

- Hardened plaque can cause rupture (aneurysm) of a blood vessel.
- Narrowing of a vessel causes blood flow to be blocked by an occlusion.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved


Acute Myocardial Infarction (2 of 6)

- Thrombus
 - Occlusion of blood flow caused by formation of a clot on rough inner surface of diseased artery
 - Can break loose and form an embolism
- Emboli can move to occlude flow of blood downstream in a smaller artery.
- Blocking of coronary artery by thrombus or embolism is an acute myocardial infarction (AMI).



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Myocardial Infarction (3 of 6)



Cross-section of a myocardial infarction.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Myocardial Infarction (4 of 6)

- Acute myocardial infarction can cause:
 - Ischemia
 - Leads to injury of cells and cell death
 - Disturbs electrical function of the heart
 - Dysrhythmias
 - Harmful changes to rate, rhythm, and pumping ability



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Myocardial Infarction (5 of 6)

- Acute myocardial infarction can cause:
 - Mechanical problems within the heart
 - Cardiogenic shock
 - Sudden death
 - Cardiac arrest within 2 hours of the onset of symptoms



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Acute Myocardial Infarction (6 of 6)

- Treatment of AMI at the hospital
 - Fibrinolytics
 - Balloon angioplasty
- Treatment upon discharge
 - Aspirin every day
 - Beta blocker



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of a Myocardial Infarction (1 of 2)

- Primary and secondary assessment
- Pattern of symptoms similar to ACS
 - Chest discomfort
 - Dyspnea
 - Nausea/vomiting
 - Syncope
 - Sweating



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of a Myocardial Infarction (2 of 2)

- Most important treatment is transport
- 12-lead ECG
- Contact ALS
- Be prepared for cardiac arrest
- Aspirin and nitroglycerin



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Heart Failure and Acute Pulmonary Edema (1 of 3)

- Heart failure
 - Congestive heart failure (CHF)
 - Heart is unable to pump blood efficiently
 - Ability to perfuse body tissues is impaired
 - Exercise-related tasks are difficult
 - Limited ability to compensate for challenge
 - Causes fluid buildup



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Heart Failure and Acute Pulmonary Edema (2 of 3)

- Right-sided failure
 - Pressure builds up in right atrium and superior and inferior vena cava
 - Jugular venous distention
 - Swelling and fluid buildup in abdomen
 - Pedal edema



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Heart Failure and Acute Pulmonary Edema (3 of 3)

- Left-sided failure
 - Pressure builds up in left atrium and pulmonary vein
 - Fluid leaks into alveoli
 - Gas exchange is impaired
 - Pulmonary edema
 - Can be life-threatening



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of Heart Failure (1 of 4)

- Assessment:
 - History of heart failure
 - Medications designed to control fluid levels
 - Pedal edema
 - Swelling in the abdomen or buttocks
 - JVD



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of Heart Failure (2 of 4)

- Assessment:
 - Pulmonary edema
 - Difficulty breathing
 - Crackles
 - Coughing up pink, frothy sputum



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of Heart Failure (3 of 4)

- Examination:
 - Primary assessment
 - May require positive pressure ventilation
 - Use OPQRST and SAMPLE
 - Detailed assessment of cardiovascular system
 - Vital signs



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Assessment and Treatment of Heart Failure (4 of 4)

- Treatment:
 - Treat signs and symptoms
 - Contact ALS for pulmonary edema
 - Consider application of CPAP
 - Administer nitroglycerin



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved


Aneurysm (1 of 2)

- Weakened sections of arterial walls begin to dilate (balloon).
- Bursting can cause rapid, life-threatening internal bleeding.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Aneurysm (2 of 2)



A weakened area in the wall of an artery will tend to balloon out, forming a sac-like aneurysm, which may eventually burst.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (1 of 6)

- Patients with acute coronary syndrome (ACS) can have many different presentations. Some complain of pressure or pain in the chest with difficulty breathing and a history of heart problems. Others may have just mild discomfort that they ignore for several hours or that goes away and returns. Some patients having heart attacks have no chest discomfort at all.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (2 of 6)

- Use a thorough secondary assessment to identify the signs and symptoms associated with the pattern of acute coronary syndrome.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (3 of 6)

- Because of the many possible presentations and the potentially severe complications of acute coronary syndrome, it is important to have a high index of suspicion and to treat patients with these symptoms aggressively. The treatment will not hurt them and may help them.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (4 of 6)

- Patients with suspected ACS who are hypoxic or short of breath need oxygen and prompt, safe transportation to definitive care. You may be able to assist patients who have their own nitroglycerin in taking it, thereby relieving pain and anxiety.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (5 of 6)

- Definitive treatment for an acute myocardial infarction occurs at the hospital. Treat suspected AMI as a time-sensitive disorder. Consider appropriate transportation destinations, activate systems of care, and obtain early 12-lead ECG, if possible.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Chapter Review (6 of 6)

- Aspirin and nitroglycerin are first-line medications associated with the treatment of acute coronary syndrome.
- Heart failure is a chronic condition that can have life-threatening acute presentations. Recognize the signs and symptoms of acute pulmonary edema and treat aggressively.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Remember (1 of 4)

- The heart is a simple pump that moves deoxygenated blood to the lungs and oxygenated blood to the body. Pressure within the cardiovascular system is critical to the moving of blood.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Remember (2 of 4)

- Acute coronary syndrome (ACS) is a blanket term that refers to a number of situations in which perfusion of the heart is inadequate.
- Although there are common symptoms of ACS, EMTs must recognize atypical findings and err on the side of caution.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Remember (3 of 4)

- Oxygen, nitroglycerine, and aspirin are key medications indicated to treat ACS. However, the definitive treatment is transportation of the patient to a facility that can open the blocked artery.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Remember (4 of 4)

- Most cardiac conditions are caused by arterial problems. Angina pectoris and acute myocardial infarction are caused by inadequate perfusion of the heart.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Questions to Consider

- What position is best for a patient with:
 - Difficulty breathing and a blood pressure of 100/70?
 - Chest pain and a blood pressure of 180/90?



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Critical Thinking

- A 62-year-old female has been complaining of severe shortness of breath and chest pain for 15 minutes prior to your arrival. When you arrive, you find the patient conscious but with difficulty breathing and oxygen saturation of 88 percent. What steps should you take in caring for this patient?



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved

Copyright



This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.



Copyright © 2021, 2016, 2012 Pearson Education, Inc. All Rights Reserved