TREBUTE® KALM PERFORMER

A high-fat, textured sweet feed formulated for mature horses at maintenance or in training.

PRODUCT CODE: T910TEN

FEATURES & BENEFITS:

- **No Corn Formula!**...to keep sugar and starch levels low.
- Higher fat, highly digestible fiber and lower sugar and starch for calm performances without loss of energy or condition.
- **Steam-crimped grains** for improved digestion.
- **Essential** amino acids for growth and muscle maintenance.
- **Essential** organic minerals replenish body tissue stores depleted during daily activities and exercise.
- **Essential** fatty acids, **Omega 3 and 6**, for healthy skin and haircoat, top performances and quick recovery.
- Optimum levels of antioxidants Vitamin E, Vitamin C and Selenium reduce damage from aging and exercise. Also boosts immunity.
- Excellent for hard-keepers and horses requiring less sugar and starch in their diets, like those who are hyperactive or have metabolic conditions such as insulin resistance, laminitis, ulcers, obesity or tying-up issues.
- With microencapsulated active dry yeast; acts as both a pre-and probiotic, increases total diet digestibility and improves overall gut health.

FEEDING DIRECTIONS:

I ELDING DIKECTIONS.		
Training Level	T910TEN (Lbs. per day)	Lbs. Hay per day
Mature Weight: 660-880 l	bs	
Idle	3-5	6-10
Light Training	5-7	8-12
Moderate Training	6-8	8-12
Intense Training	8-10	9-14
Mature Weight: 880-1,100 lbs		
Idle	4-8	9-14
Light Training	6-10	10-16
Moderate Training	8-12	10-16
Intense Training	10-14	12-18
Mature Weight: 1,100-1,3	20 lbs	
Idle	5-9	12-18
Light Training	7-11	14-20
Moderate Training	10-14	14-20
Intense Training	12-16	16 - 22

GUARANTEED ANALYSIS:

Crude Protein (Min.)
Lysine (Min.)
Methionine (Min.)
Methionine & Cystine (Min.)
Threonine (Min.)
Crude Fat (Min.)
Crude Fiber (Max.)
Calcium (Ca) (Min.)
Calcium (Ca) (Max.)
Phosphorus (P) (Min.)
Copper (Cu) (Min.)
Zinc (Zn) (Min.)
Selenium (Se) (Min.)
Potassium (K) (Min.)
Magnesium (Mg) (Min.)
Manganese (Mn) (Min.)
Cobalt (Co) (Min.)
Iron (Fe) (Min.)
Iodine (I) (Min.)
Salt (NaCl) (Min.)
Salt (NaCl) (Max.)
Vitamin A (Min.)
Vitamin D (Min.)
Vitamin E (Min.)
Vitamin C (Min.)
Biotin (Min.)
Thiamine (Min.)
Riboflavin (Min.)
Omega 6 Fatty Acids (Min.)
Omega 3 Fatty Acids (Min.)
Saccharomyces cervisiae Yeast Culture (Min.)350 million CFU/lb
Direct-Fed Microorganisms (Min.)
(Saccharomyces cervisiae, Lactobacillus acidophilus, Bacillus subtilus, Bacillus
lichenformis, Bacillus coagulans, Enterococcus faecium, Bifidobacterium
thermophilum, and Bifidobacterium longum)
Cellulase (Aspergillus Oryzae) (Min.)
Protease (Aspergillus Oryzae) (Min.)
Lipase (Aspergillus Oryzae) (Min.)
Hemicullulase (Aspergillus Niger) (Min.)10.8 Enzyme Units
Phytase (Trichoderma reesei) (Min.)

*An Enzyme Unit is defined as milligrams of substrate liberated/minute/lb. of feed

*A Phytase Unit (FTU) is defined as the quantity of enzyme which liberates one micromole of inorganic phosphate per minute from sodium phytate at 37°C, 5.5 pH

This feed contains a dry source of cellulase that breaks down cellulose, a dry source of protease that hydrolyzes proteins and increases the digestibility of protein in soybean meal based diets, a dry source of lipase that hydrolyzes triglycerides, a dry source of hemicellulase that breaks down hemicellulose, and a dry source of phytase which hydrolyzes phytate and increases the digestibility of phytin-bound phosphorous.