

## RDO

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Age: \_\_\_\_\_ Score: \_\_\_\_\_

When your back hurts, you may find it difficult to do some of the things you normally do.

*Mark only the sentences that describe you lately....*

1.  I stay at home most of the time because of my back.
2.  I walk more slowly than usual because of my back.
3.  Because of my back, I am not doing any jobs that I usually do around the house.
4.  Because of my back, I use a handrail to get upstairs.
5.  Because of my back, I lie down to rest more often.
6.  Because of my back, I have to hold onto something to get out of an easy chair.
7.  Because of my back, I try to get other people to do things for me.
8.  I get dressed more slowly than usual because of my back.
9.  I stand up only for short periods of time because of my back.
10.  Because of my back, I try not to bend or kneel down.
11.  I find it difficult to get out of a chair because of my back.
12.  My back or leg is painful almost all of the time.
13.  I find it difficult to turn over in bed because of my back.
14.  I have trouble putting on my socks (or stockings) because of pain in my back.
15.  I sleep less well because of my back.
16.  I avoid heavy jobs around the house because of my back.
17.  Because of back pain, I am more irritable and bad tempered with people than usual.
18.  Because of my back, I go upstairs more slowly than usual.

## Roland Morris Disability Questionnaire

### Scoring: **Instructions for Roland-Morris :**

- ✘ The patient is instructed to put a mark next to each appropriate statement.
- ✘ The total number of marked statements are added by the clinician. Unlike the authors of the Oswestry Disability Questionnaire, Roland and Morris did not provide descriptions of the varying degrees of disability (e.g. 40%-60% is severe disability).
- ✘ Clinical improvements over time can be graded based on the analysis of serial questionnaire scores. If, for example, at the beginning of treatment, a patient's score was 12 and, at the conclusion of treatment, her score was 2 (10 points of improvement), we would calculate an 83%  $(10/12 \times 100)$  improvement.

#### References

1. Deyo RA, Battie M, Beurskens AJ, Bombardier C, Croft P, Koes B, et al. Outcome measures for low back pain research. *Spine* 1998;23:2003-2013.
2. Roland M, Morris R. A study of the natural history of back pain: part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine* 1983;8:141-144.
3. Deyo RA. Comparative validity of the sickness impact profile and shorter scales for functional assessment in low back pain. *Spine* 1986;11:951-954.
4. Jensen MP, Strom SE, Turner JA, Romano JM. Validity of the Sickness Impact Profile Roland scale as a measure of dysfunction in chronic pain patients. *Pain* 1992;50:157-162.
5. Patrick DL, Deyo RA, Atlas SJ, Singer DE, Chapin A, Keller RB. Assessing health related quality of life in patients with sciatica. *Spine* 1995;20:1899-909.
6. Roberts A. The conservative treatment of low back pain. MD thesis, University of Nottingham, 1991.
7. Waddell G. *The Back Pain Revolution*. Edinburgh: Churchill Livingstone, 1998.
8. Baker CD, Pynsent PB, Fairbank JCT. The Oswestry Disability Index revisited: its reliability, repeatability and validity, and a comparison with the St. Thomas's Disability Index. In: Roland MO, Jenner JR, eds. *Back Pain: New Approaches to Education and Rehabilitation*. Manchester University Press, 1989:174-86.
9. Stratford PW, Binkley JM. Measurement properties of the RM 18: a modified version of the Roland-Morris disability scale. *Spine* 1997;22:2416-2421.
10. CareTrak outcomes software. Grand Rapids, MN; (800) 393-7255, [www.caretrak-outcomes.com](http://www.caretrak-outcomes.com).