

Solomon Press

Core Mathematics C3

Paper J

(Mark Scheme)

All exam papers are issued free to students for education purpose only.
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com

GCE Examinations
Advanced Subsidiary

Core Mathematics C3

Paper J

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

C3 Paper J – Marking Guide

1. (a) $\cos^2 x = (\sqrt{3} - 1)^2 = 3 - 2\sqrt{3} + 1 = 4 - 2\sqrt{3}$ M1
 $\cos 2x = 2 \cos^2 x - 1 = 2(4 - 2\sqrt{3}) - 1 = 7 - 4\sqrt{3}$ M1 A1
- (b) $2(\cos y \cos 30 - \sin y \sin 30) = \sqrt{3} (\sin y \cos 30 - \cos y \sin 30)$ M1 A1
 $\sqrt{3} \cos y - \sin y = \frac{3}{2} \sin y - \frac{1}{2}\sqrt{3} \cos y$ B1
 $\frac{3}{2}\sqrt{3} \cos y = \frac{5}{2} \sin y$
 $\tan y = \frac{3}{2}\sqrt{3} \div \frac{5}{2} = \frac{3}{5}\sqrt{3}$ M1 A1 (8)
-

2. (a) $f(x) = (x - \frac{3}{2})^2 - \frac{9}{4} + 7 = (x - \frac{3}{2})^2 + \frac{19}{4}$ M1 A1
 $\therefore f(x) \geq \frac{19}{4}$ A1
- (b) $= g(11) = 21$ M1 A1
- (c) $fg(x) = f(2x - 1) = (2x - 1)^2 - 3(2x - 1) + 7$ M1
 $\therefore 4x^2 - 4x + 1 - 6x + 3 + 7 = 17$
 $2x^2 - 5x - 3 = 0$ A1
 $(2x + 1)(x - 3) = 0$ M1
 $x = -\frac{1}{2}, 3$ A1 (9)
-

3. (a)
$$\begin{array}{r} x^2 + 4x - 4 \\ x^2 - 3x + 3 \overline{) x^4 + x^3 - 13x^2 + 26x - 17} \\ x^4 - 3x^3 + 3x^2 \\ \hline 4x^3 - 16x^2 + 26x \\ 4x^3 - 12x^2 + 12x \\ \hline - 4x^2 + 14x - 17 \\ - 4x^2 + 12x - 12 \\ \hline 2x - 5 \end{array}$$
 M1
- $\therefore f(x) = x^2 + 4x - 4 + \frac{2x-5}{x^2-3x+3}, A = 4, B = -4, C = 2, D = -5$ A3
- (b) $f'(x) = 2x + 4 + \frac{2 \times (x^2 - 3x + 3) - (2x - 5) \times (2x - 3)}{(x^2 - 3x + 3)^2}$ M1 A2
 $x = 1 \Rightarrow y = -2, \text{ grad} = 5$
 $\therefore \text{grad of normal} = -\frac{1}{5}$ M1
 $\therefore y + 2 = -\frac{1}{5}(x - 1)$ M1
 $5y + 10 = -x + 1$
 $x + 5y + 9 = 0$ A1 (10)
-

4. (a) $\frac{dx}{dy} = \frac{1}{2} \sec \frac{y}{2} \tan \frac{y}{2}$ M1
 $0 \leq y < \pi \therefore \tan \frac{y}{2} \geq 0 \therefore \frac{dx}{dy} = \frac{1}{2} \sec \frac{y}{2} \sqrt{\sec^2 \frac{y}{2} - 1} = \frac{1}{2} x \sqrt{x^2 - 1}$ M1 A1
 $\frac{dy}{dx} = 1 \div \frac{dx}{dy} = \frac{2}{x \sqrt{x^2 - 1}}$ M1 A1
- (b) $\frac{dy}{dx} = \frac{1}{2}(3 + 2 \cos x)^{-\frac{1}{2}} \times (-2 \sin x) = -\frac{\sin x}{\sqrt{3 + 2 \cos x}}$ M1 A1
 $x = \frac{\pi}{3}, y = 2, \text{ grad} = -\frac{1}{4}\sqrt{3}$ M1 A1
 $\therefore y - 2 = -\frac{1}{4}\sqrt{3}(x - \frac{\pi}{3}) \quad [3\sqrt{3}x + 12y - 24 - \pi\sqrt{3} = 0]$ M1 A1 (11)
-

5.	(a) $f(x) > 5$	B1
	(b) $y = 5 + e^{2x-3}$	
	$2x - 3 = \ln(y - 5)$	M1
	$x = \frac{1}{2}[3 + \ln(y - 5)]$	M1
	$\therefore f^{-1}(x) = \frac{1}{2}[3 + \ln(x - 5)], x \in \mathbb{R}, x > 5$	A2
	(c) $x = f^{-1}(7) = \frac{1}{2}(3 + \ln 2)$	M1 A1
	(d) $f'(x) = 2e^{2x-3}$	M1
	grad = 4	A1
	$\therefore y - 7 = 4[x - \frac{1}{2}(3 + \ln 2)]$	M1 A1
	[$y = 4x + 1 - 2\ln 2$]	(11)

6.	(a) $LHS \equiv \frac{2\cos 2x}{\sin 2x} + \frac{\sin x}{\cos x}$	M1
	$\equiv \frac{\cos 2x}{\sin x \cos x} + \frac{\sin x}{\cos x}$	M1
	$\equiv \frac{\cos 2x + \sin^2 x}{\sin x \cos x}$	A1
	$\equiv \frac{(\cos^2 x - \sin^2 x) + \sin^2 x}{\sin x \cos x}$	M1
	$\equiv \frac{\cos^2 x}{\sin x \cos x}$	
	$\equiv \frac{\cos x}{\sin x}$	
	$\equiv \cot x \equiv RHS$	A1
	(b) $\cot x = \operatorname{cosec}^2 x - 7$	
	$\cot x = 1 + \cot^2 x - 7$	M1
	$\cot^2 x - \cot x - 6 = 0$	
	$(\cot x + 2)(\cot x - 3) = 0$	M1
	$\cot x = -2 \text{ or } 3$	A1
	$\tan x = -\frac{1}{2} \text{ or } \frac{1}{3}$	M1
	$x = \pi - 0.4636 \text{ or } 0.32$	
	$x = 0.32, 2.68 \text{ (2dp)}$	A2
		(11)

7.	(a) $f(x) \geq 0$	B1
	(b) $= f(0) = 5$	M1 A1
	(c) $fg(x) = f[\ln(x + 3)] = 2 \ln(x + 3) - 5 $	M1
	$\therefore 2 \ln(x + 3) - 5 = 3$	
	$2 \ln(x + 3) = 2, 8$	M1
	$\ln(x + 3) = 1, 4$	A1
	$x = e - 3, e^4 - 3$	M1 A1
	(d) let $h(x) = f(x) - g(x)$	
	$h(3) = -0.79, f(4) = 1.1$	M1
	sign change, $h(x)$ continuous \therefore root	A1
	(e) $x_1 = 3.396, x_2 = 3.428, x_3 = 3.430, x_4 = 3.431$	M1 A2
	(f) $h(3.4305) = -0.000052, f(3.4315) = 0.0018$	M1
	sign change, $h(x)$ continuous \therefore root $\therefore \alpha = x_4$ to 4sf	A1
		(15)

Total (75)