



Cancer as a Metabolic Disease: Implications for Novel Therapeutics

Journal:	<i>Carcinogenesis</i>
Manuscript ID:	CARCIN-2013-00896.R2
Manuscript Type:	Review
Date Submitted by the Author:	19-Nov-2013
Complete List of Authors:	Seyfried, Thomas; Boston College, Biology Flores, Roberto; Boston College, Biology Poff, Angela; University of South Florida, Molecular Pharmacology and Physiology D'Agostino, Dominic; University of South Florida, Molecular Pharmacology and Physiology
Keywords:	somatic mutations, Warburg effect, Crabtree effect, metabolism, mitochondria

SCHOLARONE™
Manuscripts

Cancer as a Metabolic Disease: Implications for Novel Therapeutics

Thomas N. Seyfried¹, Roberto Flores¹, Angela M. Poff², and Dominic P. D’Agostino²

¹Biology Department, Boston College, Chestnut Hill, MA 02467

²Department of Molecular Pharmacology and Physiology, University of South Florida, Tampa, Florida,

Corresponding Author: Thomas N. Seyfried

Phone: +1-617-552-3563

Fax: +1- 617-552-2011

Email: thomas.seyfried@bc.edu

Abstract

Emerging evidence indicates that cancer is primarily a metabolic disease involving disturbances in energy production through respiration and fermentation. The genomic instability observed in tumor cells and all other recognized hallmarks of cancer are considered downstream epiphenomena of the initial disturbance of cellular energy metabolism. The disturbances in tumor cell energy metabolism can be linked to abnormalities in the structure and function of the mitochondria. When viewed as a mitochondrial metabolic disease, the evolutionary theory of Lamarck can better explain cancer progression than can the evolutionary theory of Darwin. Cancer growth and progression can be managed following a whole-body transition from fermentable metabolites, primarily glucose and glutamine, to respiratory metabolites, primarily ketone bodies. As each individual is a unique metabolic entity, personalization of metabolic therapy as a broad-based cancer treatment strategy will require fine-tuning to match the therapy to an individual's unique physiology.

Abbreviations:

ROS (reactive oxygen species)
OxPhos (oxidative phosphorylation)
CR (calorie restriction)
KD (ketogenic diet)
HBO₂T (hyperbaric oxygen therapy)
SLP (substrate level phosphorylation)
RTG (retrograde response)
TCA (tricarboxylic acid)
GBM (glioblastoma multiforme)

Introduction

Cancer is a disease involving multiple time- and space-dependent changes in the health status of cells and tissues that ultimately lead to malignant tumors. Neoplasia (abnormal cell growth) is the biological endpoint of the disease. Tumor cell invasion into surrounding tissues and their spread (metastasis) to distant organs is the primary cause of morbidity and mortality of most cancer patients (1-5). A major impediment in the effort to control cancer has been due in large part to the confusion surrounding the origin of the disease. Contradictions and paradoxes continue to plague the field (6-10). Much of the confusion surrounding cancer origin arises from the absence of a unifying theory that can integrate the many diverse observations on the nature of the disease. Without a clear understanding of how cancer arises, it becomes difficult to formulate a successful strategy for effective long-term management and prevention. The failure to clearly define the origin of cancer is responsible in large part for the failure to significantly reduce the death rate from the disease (2). Although cancer metabolism is receiving increased attention, cancer is generally considered a genetic disease (10,11). This general view is now under serious reevaluation (2,12). The information in this review comes in part from our previous articles and treatise on the subject (2,13-17).

Provocative Question: Does cancer arise from somatic mutations?

Most of those who conduct academic research on cancer would consider it a type of somatic genetic disease where damage to a cell’s nuclear DNA underlies the transformation of a normal cell into a potentially lethal cancer cell (7,10,11,18). Abnormalities in dominantly expressed oncogenes and in recessively expressed tumor suppressor genes have been the dogma driving the field for several decades (7,10). The discovery of millions of gene changes in different cancers has led to the perception that cancer is not a single disease, but is a collection of many different diseases (6,11,19,20). Consideration of cancer as a “disease complex” rather than as a single disease has contributed to the notion that management of the various forms of the disease will require individual or “personalized” drug therapies (2,21-23). Tailored therapies, unique to the genomic defects within individual tumors, are viewed as the

future of cancer therapeutics (2,24). This therapeutic strategy would certainly be logical if the nuclear somatic mutations detected in tumors were the drivers of the disease. How certain are we that tumors arise from somatic mutations and that some of these mutations drive the disease? It would therefore be important to revisit the origin of the gene theory of cancer.

The gene theory of cancer originated with Theodor Boveri's suggestion in 1914 that cancer could arise from defects in the segregation of chromosomes during cell division (18,25-29). As chromosomal instability in the form of aneuploidy (extra chromosomes, missing chromosomes, or broken chromosomes) is present in many tumor tissues (21,30-32), it was logical to extend these observations to somatic mutations within individual genes including oncogenes and tumor suppressor genes (18,33-36). Boveri's hypothesis on the role of chromosomes in the origin of malignancy was based primarily on his observations of chromosome behavior in nematodes (*Ascaris*) and sea urchins (*Paracentrotus*) and from his consideration of von Hansemann's earlier observations of abnormal chromosome behavior in tumors (18,25,29). In contrast to Boveri's view of aneuploidy as the origin of cancer, von Hansemann considered the abnormal chromosome behavior in tumors as an effect rather than as a cause of the disease (25). Although Boveri's hypothesis emerged as the foundation for the somatic mutation theory of cancer, it appears that he never directly experimented on the disease (18,25,29). The reason for the near universal acceptance of Boveri's hypothesis for the origin of cancer is not clear, but might have been linked to his monumental achievement in showing that Gregor Mendel's abstract heredity factors resided on chromosomes (29). Boveri's cancer theory was also consistent with the gradual accumulation of evidence showing that DNA abnormalities are abundant in cancer cells.

In his 2002 review, Knudson stated that, "considerable evidence has been amassed in support of Boveri's early hypothesis that cancer is a somatic genetic disease" (37). The seeds of the somatic mutation theory of cancer might have been sowed even before the work of von Hansemann and Boveri. Virchow considered that cancer cells arose from other cancer cells (38). Robert Wagner provided a good overview of those early studies

leading to the idea that somatic mutations give rise to cancer (38). It gradually became clear that almost every kind of genomic defect could be found in tumor cells whether or not the mutations were connected to carcinogenesis (10,11,18,26,31). The current somatic mutation theory involves a genomic landscape of incomprehensible complexity that also includes mysterious genomic “Dark Matter” (2,10,11,19). While massive evidence exists showing that genomic instability is present to some degree in all tumor cells, it is unclear how this phenotype relates to the origin of the disease. It appears that almost every neoplastic cell within a naturally arising human tumor is heterogeneous in containing a unique genetic architecture (31).

Inconsistencies with a nuclear gene origin of cancer

The distinguished British geneticist, C. D. Darlington, was one of the first to raise concerns regarding the nuclear genetic origin of cancer (39). Based on several inconsistencies in the association of mutagens with cancer, Darlington argued persuasively that nuclear genomic defects could not be the origin of cancer. Rather, he was convinced that cancer cells arose from defects in cytoplasmic elements, which he referred to as “plasmagenes”. While Darlington did not specifically characterize the nature of the plasmagene, several characteristics of the plasmagenes suggested that they were mitochondria. It was unclear, however, if the radiation damage to the plasmagenes acted alone in causing cancer or also acted in conjunction with mutations in nuclear genes.

Inconsistencies regarding the somatic nuclear gene theory of cancer also come from nuclear/cytoplasmic transfer experiments between tumorigenic and non-tumorigenic cells. Several investigators showed that tumorigenicity is suppressed when cytoplasm from non-tumorigenic cells, containing normal mitochondria, is combined with nuclei from tumor cells (40-44). Moreover, the *in vivo* tumorigenicity of multiple human and animal tumor types is suppressed when the nucleus from the tumor cell is introduced into the cytoplasm of a non-tumorigenic cell (45-48). Tumors generally did not form despite the continued presence of the tumor-associated mutations. The nuclear gene mutations documented in mouse brain tumors and melanomas were also detected in

the normal embryonic tissues of the mice derived from the tumor nuclei (47,48). Some embryos derived from tumor nuclei, which contained major chromosomal imbalances, proceeded through early development forming normal appearing tissues before dying. Despite the presence of tumor-associated aneuploidy and somatic mutations, tumors did not develop from these tumor-derived nuclei (49). Boveri also found that sea urchin embryos with chromosomal imbalances developed normally to gastrulation, but then aborted (25,29). Hochedlinger and colleagues showed that nuclei derived from melanoma cells were unable to direct complete mouse development due presumably to the chromosomal imbalances and irreversible tumor-associated mutations in the melanoma nucleus (48). Tumors did not arise in the embryos derived from the melanoma nuclei. These findings suggest that the nuclear genomic defects in these tumor cells have more to do with directing development than with causing tumors.

More recent mitochondrial transfer experiments support the general findings of the nuclear transfer experiments. (50,51). The tumorigenic phenotype is suppressed when normal mitochondria are transferred to the tumor cell cytoplasm. On the other hand, the tumorigenic phenotype is enhanced when tumor mitochondria are transferred to a normal cell cytoplasm. These findings further suggest that tumorigenesis is dependent more on mitochondrial function than on the types of mutations in the nucleus.

In contrast to the suppressive effects of normal mitochondria on tumorigenicity, tumorigenicity is enhanced when nuclei of non-tumorigenic cells are combined with cytoplasm from tumor cells (52,53). These observations are consistent with the original view of Darlington that tumor cells arise from defects in the cytoplasm rather than from defects in the nucleus (39). Wallace and co-workers also showed that introduction of mitochondrial DNA mutations into non-tumorigenic cybrids could reverse the anti-tumorigenic effect of normal mitochondria leading to the conclusion that cancer can be best defined as a type of mitochondrial disease (53). The nuclear transfer studies are summarized in **Fig. 1** and highlight the role of the mitochondria in suppressing tumorigenesis. These studies also raise questions regarding the role of somatic mutations as drivers of tumorigenesis. Further studies will be needed to determine

whether tumors arise from defects in the nuclear genome alone, in the mitochondria alone, or require defects in both the mitochondria and the nuclear genome. Such studies will provide evidence for or against the nuclear gene driver hypothesis of cancer initiation.

Respiratory insufficiency as the origin of cancer and the “Warburg Effect”

Otto Warburg first proposed that all cancers originate from dysfunctional cellular respiration (54,55). Warburg stated, *“Just as there are many remote causes of plague, heat, insects, rats, but only one common cause, the plague bacillus, there are a great many remote causes of cancer-tar, rays, arsenic, pressure, urethane- but there is only one common cause into which all other causes of cancer merge, the irreversible injuring of respiration”*. The key points of Warburg’s theory are; 1) insufficient respiration initiates tumorigenesis and ultimately cancer, 2) energy through glycolysis gradually compensates for insufficient energy through respiration, 3) cancer cells continue to ferment lactate in the presence of oxygen, and 4) respiratory insufficiency eventually becomes irreversible (54-58). Efraim Racker was the first to describe the increased aerobic glycolysis seen cancer cells as the “Warburg effect” (59). Warburg, however, referred to the phenomenon in cancer cells as “aerobic fermentation” to highlight the abnormal production of lactate in the presence of oxygen (54-58). As lactate production is widely recognized as an indicator of respiratory insufficiency in biological systems (60), Warburg also viewed the aerobic production of lactate in tumor cells as an indicator of respiratory insufficiency.

A deficiency in OxPhos energy is responsible for lactate production in most cases (61,62). For example, muscle cells significantly increase their metabolic rate when oxygen becomes limiting during intense exercise. The oxygen deficiency causes a lack of energy through OxPhos prompting lactate production in an effort to provide compensatory energy from fermentation (glycolytic energy) (60). A competing argument would be that OxPhos is not insufficient during intense exercise and that aerobic fermentation is needed to provide more energy and growth metabolites in response to the increased work demand. This would be similar to the suggestion of

Weinhouse and others for the increased aerobic glycolysis in tumor cells (63,64). Indeed, Dang and co-workers suggest that the increased lactate production in tumor cells arises from damage to the regulation of glycolysis and not to insufficient respiration (64). However, the competing argument is inconsistent with the observation that the lactate made by muscle cells during intense exercise falls significantly after oxygen is restored to the muscle tissue. This would indicate that the lactate was made primarily because O_2 was unavailable for robust OxPhos (65). In addition, oxygen deprivation or hypoxia causes all known cultured mammalian cells to increase lactate production (66-68). An increase in lactate is also seen in adequately oxygenated cells when respiration is inhibited either by respiratory poisons or null mutations in key respiratory enzymes (69-71). It is therefore clear from established bio-energetic principles that the excess lactate made by most mammalian cells is needed to sustain fermentation energy in order to compensate for insufficient energy from respiration. It is our view that tumor cells are not an exception to this general principle and that their lactate production results in part from insufficient respiratory activity. It is expected that an up-regulation of glycolytic genes would be needed to facilitate compensatory energy production through glycolysis when cellular respiration is deficient for protracted periods (56). The reduction of pyruvate to lactate is needed to enhance the glycolytic pathway when respiration becomes insufficient.

It is important to recognize that pyruvate is produced through aerobic glycolysis in most normal cells of the body that use glucose for energy. The reduction of pyruvate to lactate distinguishes the tumor cells from most normal cells, which fully oxidize pyruvate to CO_2 and water for ATP production through the TCA cycle and the electron transport chain (56). Aerobic glycolysis with lactate production can occur in normal retina though more ATP is produced through respiration than through glycolysis, as is the case in most respiring tissues (72). On the other hand, enhanced aerobic glycolysis without significant lactate production or energy through fermentation can occur in normal cardiac and brain tissues under conditions of increased activity (73-75). The slight transient increase in lactate production under these conditions is not associated with a significant increase in total energy production. As enhanced aerobic glycolysis does not

produce significant lactate in normal cells under well-oxygenated conditions, a phenotype of enhanced aerobic glycolysis is therefore not synonymous with a Warburg effect.

Lactate will be produced in normal tissues under low oxygen conditions. Tumor cells also produce lactate under hypoxia through anaerobic glycolysis. Although many investigators of tumor cell energy metabolism use the term “aerobic glycolysis” in referring to the Warburg effect, we consider the term “aerobic fermentation” as a more accurate description of the Warburg effect since aerobic glycolysis occurs in most normal cells of the body. A key issue is whether the lactate produced in tumor cells under aerobic conditions results from insufficient respiration as Warburg proposed, or is due to some other phenomenon. The origin of the Warburg effect is an issue of controversy that persists today despite Warburg’s data showing that it arose from insufficient respiration.

According to Warburg and Burk respiratory insufficiency together with lactate production are the key features of tumor cell energy metabolism (55,57,76,77). Respiratory insufficiency as the origin of tumorigenesis has remained controversial, however, due to observations of high oxygen consumption rates in many tumor cells (63,64,78-82). It is generally assumed that oxygen consumption rate is a good indicator of cellular respiration and OxPhos. While low oxygen consumption rate seen together with high lactate production can be indicative of insufficient respiration, high oxygen consumption might not be indicative of sufficient respiration especially if lactate is also produced. It is now recognized from numerous studies that oxygen consumption rates are not always linked to a normally coupled oxidative phosphorylation (83-86). It can be difficult to determine the degree to which mitochondrial ATP production arises from coupled respiration or from TCA cycle substrate level phosphorylation (87-90). The origin of mitochondrial ATP production in tumor cells requires further clarification in light of these issues.

Mitochondrial structure is intimately connected to mitochondrial function. This fact

cannot be overemphasized. We have reviewed substantial evidence of morphological, proteomic, and lipidomic abnormalities in mitochondria of numerous types of cancer cells (17,85,91). Tumor cells can have abnormalities in both the content and composition of their mitochondria. The work of Arismendi-Morillo and Oudard et al showed that the ultrastructure of tumor tissue mitochondria differs markedly from the ultrastructure of normal tissue mitochondria (17,92-94). In contrast to normal mitochondria, which contain numerous cristae, mitochondria from tumor tissue samples showed swelling with partial or total cristolysis (**Fig. 2**). Cristae contain the proteins of the respiratory complexes and play an essential structural role in facilitating energy production through OxPhos (95). The structural defects in human glioma mitochondria are also consistent with lipid biochemical defects in murine gliomas (96,97).

More recent electron micrographic studies from Elliott and co-workers showed that mitochondria ultrastructure was abnormal to some degree in 778 patients with breast cancer (51). Remarkably, mitochondria were severely reduced in number or were undetectable in the tumor tissue from over 80% of the patients. These findings together with the evidence from the Pedersen review would support Warburg's central hypothesis that respiration is insufficient in tumor cells (67). *It is obvious that mitochondrial function or OxPhos sufficiency cannot be normal in tumor cells that contain few if any mitochondria.* Glycolysis and lactate fermentation would need to be upregulated in these tumor cells in order to compensate for the absence of OxPhos. Furthermore, the degree of malignancy in these breast tumors was correlated directly with the degree of mitochondrial structural abnormality (51). The high glycolytic activity and lactate production seen in the most malignant tumors were also linked to the mitochondrial structural abnormalities seen in the tumors (91,98-102). In contrast to inherited mitochondriopathies, where glycolysis might not compensate completely for mitochondrial energy failure, fermentation energy appears capable of compensating completely for the respiratory insufficiency in tumor cells (18,103). Further studies will be needed to distinguish the differences in glycolytic and respiratory energy metabolism in tumor cells and in cells with mitochondriopathies (18).

Pedersen presented massive evidence showing that mitochondria in tumor cells are abnormal when compared with mitochondria from normal cells (67). His review provides a comprehensive discussion of mitochondrial bioenergetics and dysfunction in cancer cells. It was clearly shown that the mitochondria of cancer cells contain numerous qualitative and quantitative abnormalities when compared with mitochondria from tissue specific control cells. Summarized here are just a few of the conclusions from the Pedersen review. 1) Tumor mitochondria are abnormal in morphology and ultrastructure and respond differently to changes in growth media than mitochondria from normal cells. 2) The protein and lipid composition of tumor mitochondria are markedly different from that of normal mitochondria. 3) Proton leak and uncoupling is greater in tumor mitochondria than in normal mitochondria. 4) Calcium regulation is impaired in tumor mitochondria. 5) Anion membrane transport systems are abnormal or dysregulated in mitochondria from many tumors. 6) Defective shuttle systems are not responsible for elevated glucose fermentation in tumor cells. 7) Pyruvate is not effectively oxidized in tumor mitochondria. 8) Tumor mitochondria contain a surface-bound, fetal-like hexokinase. 9) A deficiency in some aspect of respiration can account for excessive lactic acid production in tumor cells. Clearly, substantial evidence exists showing that mitochondrial structure, function, and respiratory capacity is defective to some degree in all types of tumor cells. This information should be addressed in discussions of tumor cell energy metabolism.

Besides a generalized defect at the level of the mitochondrial electron transport chain in most tumor cells, numerous other mitochondrial abnormalities do exist that would diminish respiratory function (104,105). Interestingly, Warburg never stated that a generalized defect in electron transport was responsible for the origin of cancer despite suggestions from others (106,107). Rather, Warburg stated that insufficient respiration was responsible for aerobic fermentation and the origin of cancer (54,55,57,58,82). We know from the work of numerous investigators that electron transport may not be coupled to ATP synthesis in cancer cells (91,104). Any mitochondrial defect that would uncouple electron transport from OxPhos could reduce respiratory sufficiency and thus contribute to lactate formation or a Warburg effect.

Influence of unnatural growth environment on cellular energy metabolism

Much of the evidence arguing against Warburg's central theory that respiratory insufficiency is the origin of the aerobic fermentation seen in cancer cells (Warburg effect) was derived from investigations of tumor cells grown *in vitro* (64,78,79,108-110). In contrast to the structural defects, reduced numbers, or the absence of mitochondria observed in human cancerous tissues, such mitochondrial abnormalities are not generally seen in many human and animal tumor cells when they are grown in the *in vitro* environment. It is interesting that oxygen consumption rate can be similar or even greater in cultured tumor cells than in non-tumorigenic cells (83,86,111). The presence of mitochondria and robust oxygen consumption rates in tumor cells grown *in vitro* suggested to some that mitochondria are normal in tumor cells and that Warburg's central theory was incorrect (64,81,109). As mentioned above, however, oxygen consumption rate is not always an indicator of coupled respiration. Some tumor cells consume oxygen while importing and hydrolyzing glycolytically derived ATP through the mitochondrial adenine nucleotide transporter 2 in order to maintain the proton motive gradient (112). We also showed that the growth of tumorigenic and non-tumorigenic cells in typical cell culture media changes the content and fatty acid composition of lipids especially cardiolipin, the signature phospholipid of the inner mitochondrial membrane that regulates OxPhos (96). No tumor cells have yet been described with a normal content and composition of cardiolipin (97,113,114). Cells cannot respire effectively if the content or composition of their cardiolipin is abnormal (97,115,116). This point cannot be overemphasized.

It is not clear why mitochondria might appear functionally normal in many types of cultured tumor cells, but appear structurally abnormal when evaluated in the tumor cells of many primary malignant cancers. Cultured cell lines are usually derived from only a single cell or a few cells of a heterogeneous tumor. *Is it possible that only those tumor cells with some level of mitochondrial function are capable of growing in vitro?* Also the *in vitro* environment forces many cells into a state of aerobic fermentation whether or not they are tumorigenic. We showed that the typical culture environment produces

immature cardiolipin in non-tumorigenic glial cells, which reduces the activity of mitochondrial respiratory chain complexes (96). Further studies are needed on the structure and function of mitochondria in tumor tissue and their derived cell lines.

Lactate production should be minimal in adequately oxygenated cells that have the capacity to respire normally. However, significant lactate production is often observed in proliferating non-tumorigenic cells grown in well-oxygenated cultures (96,103,117). It is not likely that the high aerobic fermentation seen in normal cells grown in culture is due to deregulated glycolysis, as suggested for tumor cells (64). Enhanced glycolysis in tumor cells cannot be considered only as deregulated, but can also be considered as necessary to compensate for respiratory insufficiency.

Some investigators consider lactate production as necessary for normal cell proliferation (118,119). It is important to consider the differences in the metabolic requirements of tumorigenic and non-tumorigenic cells when grown in the *in vivo* and *in vitro* environments (117,120). In contrast to what is seen in cultured cells, no lactate production is seen in the rapidly growing embryonic chorion under aerobic conditions (57). Moreover, lactate production is minimal in rapidly growing hepatocytes during liver regeneration (121,122). Instead, regenerating liver cells use fatty acids rather than glucose to fuel proliferation. Fatty acid metabolism produces mostly water and CO₂, but not lactate. In contrast to hepatomas, which have abnormal cardiolipin composition, the content and composition of cardiolipin is similar in resting liver cells and in proliferating liver cells during regeneration (123,124). These findings suggest that respiration can occur normally in rapidly proliferating liver cells during liver regeneration. Viewed together, these findings indicate that lactate production is not required for rapid cell proliferation *in vivo*. Tumor cells are an exception in this regard, as lactate production in these cells arises as a consequence of abnormal respiration, which can be linked to either the structural defects seen in tumor tissue mitochondria or to reduced number of mitochondria. If lactate production is not required for rapid cell growth, why are significant amounts of lactate produced in many types of rapidly growing tumorigenic and non-tumorigenic cells when grown in culture?

The *Crabtree effect* can confound the interpretation of energy metabolism in cultured cells. The Crabtree effect involves a glucose-induced suppression of respiration leading to lactate production whether or not mitochondria are damaged (96,120,125,126). The Crabtree effect differs from the Warburg effect, which involves lactate production arising from insufficient respiration. In other words, the aerobic lactate produced under the Crabtree effect arises from a suppressed respiration rather than from insufficient respiration as occurs in the Warburg effect. It can be difficult to determine with certainty, however, whether the aerobic fermentation (aerobic glycolysis) observed in cultured cells arises from a Crabtree effect, a Warburg effect, or from some combination of these effects (126-128). We consider the Crabtree effect as an artifact of the *in vitro* environment that causes some non-tumorigenic mammalian cells to ferment lactate even in the presence of oxygen. It would therefore be important for investigators to exclude the influence of a Crabtree effect on the assessment of energy measurements in cultured cells. Although a Crabtree effect might suppress OxPhos, the TCA cycle should remain functional and produce ATP through substrate level phosphorylation (87-90). Under certain conditions (hypoxia), the tumor TCA cycle can work in both forward and reverse (reductive) directions (129,130). Although some tumor cells can have a functional TCA cycle linked to insufficient respiration, sufficient respiration is unlikely to occur without a functional TCA cycle. Support for this comes from findings that some rare cancers can arise from inherited mutations in TCA enzymes, e.g., fumarate hydratase and succinate dehydrogenase, which impede the TCA cycle (131,132). Based on the data presented over many years by numerous investigators, we consider that OxPhos is universally insufficient to some degree in all tumor cells. However, the Crabtree effect and the unnatural conditions of the *in vitro* environment can obscure this insufficiency. While respiratory insufficiency might be more profound in some tumor cells than in others, most if not all tumor cells will express some degree of OxPhos insufficiency when compared with appropriate controls matched for species, age, and tissue type.

Besides the confounding influence of the *in vitro* environment on energy metabolism,

1
2
3
4 abnormalities and misinformation can be obtained when human tumor cells are grown in
5 non-syngeneic hosts (133). This is especially relevant with respect to the mouse
6 xenograft models including the *patient derived xenografts*. We found that human
7 U87MG brain cancer cells express mouse carbohydrates on their surface when grown
8 as a xenograft in immune deficient mice (134). Over 65% of the sialic acid composition
9 on the U87MG tumor cells consisted of the nine-carbon sugar, N-glycolylneuraminic
10 acid. Humans, however, are unable to synthesize N-glycolylneuraminic acid due to a
11 mutation in the gene that encodes a common mammalian hydroxylase enzyme
12 (134,135). The hydroxylase mutation occurred in the human genome sometime after
13 our evolutionary split with the great apes (135). The acquisition of murine
14 carbohydrates and lipids will likely occur in any human tumor cell grown in the body of a
15 mouse or rat. N-glycolylneuraminic acid alters the characteristics of human embryonic
16 stem cells when grown on non-human feeder cells (136). The influence of the murine
17 host on gene expression in human tumor cells is a confounding variable that can create
18 difficulties for data interpretation in tumor cells. Few investigators address these issues.

19
20
21
22
23
24
25
26
27
28
29
30
31
32 Expression of mouse carbohydrates and lipids on human tumor cells when grown as
33 xenografts can alter gene expression patterns and growth behavior of the tumor cells
34 thus altering their response to changes in the microenvironment. It might be reasonable
35 to view the human xenograft tumor models as a type of human-mouse centaur (133). In
36 addition, the basal metabolic rate of the mouse is 7-8-fold greater than that of humans
37 (137,138). The basal metabolic rate is the energy needed for the maintenance of all
38 physiological processes under rest. Little attention is given to differences in metabolic
39 rate when comparing metabolism among human and animal tumors (117). The
40 difference in metabolic rate could cause the human tumor cells to grow slow or not at all
41 in xenografts due to competition for energy metabolites with mouse host stromal cells
42 that have a higher metabolic rate than the human tumor cells. This could account in
43 part for the low incidence of systemic metastasis seen in xenograft models implanted
44 with tumor cells taken from human metastases. Solid tumors that do not metastasize or
45 are not invasive are generally considered benign (4). Further studies will be needed to
46 determine if the human tumor cells that are selected to grow in the mouse have a

metabolic rate more similar to that of the mouse than to that of the human.

Many human tumor cells or tissues are grown in mice that are **Non-Obese Diabetic** and have **Severely Compromised Immuno-Deficiency (NOD-SCID)** (139). These mice not only have a compromised innate and/or adaptive immune system, but they also express characteristics of both type-1 diabetes and type-2 diabetes (140). This is not a usual situation for most cancer patients. Despite some limited success, it is naive to assume that the growth behavior and response to therapies of human tumors grown as xenografts would be similar to the situation in the natural host. The evaluation of cancer drugs against tumor cells grown in unnatural environments together with the misunderstanding on the origin of cancer is responsible in large part for widespread failure in developing new cancer therapies (133). The use of syngeneic mouse tumor models will be more representative of the natural physiological state in humans than will the xenograft models.

Connecting the links from respiratory insufficiency to cancer origin

The path from normal cell physiology to malignant behavior, where all major cancer hallmarks are expressed is depicted in **Fig. 3**. Any unspecific condition that damages a cell's respiratory capacity but is not severe enough to kill the cell can potentially initiate the path to a malignant cancer. Reduced respiratory capacity could arise from damage to any mitochondrial protein, lipid, or mtDNA. Some of the many unspecific conditions that can diminish a cell's respiratory capacity thus initiating carcinogenesis include inflammation, carcinogens, radiation (ionizing or ultraviolet), intermittent hypoxia, rare germline mutations, viral infections, and age. The evidence supporting this statement also addresses Szent Giorgy's "oncogenic paradox", as was described in a recent treatise on the subject (141). The paradox addresses the difficulty in knowing how a plethora of disparate carcinogenic agents might produce cancer through a common mechanism. Some of the rare germline mutations that increase risk for cancer through an effect on cellular respiration include *p53*, *BRACA1*, *RB*, and xeroderma pigmentosum (18). Cancer-causing viruses can be linked to mitochondrial dysfunction (18). If respiratory damage is acute, the cell will die. On the other hand, if damage is

mild and protracted, the cell will elevate lactate or amino acid fermentation in order to compensate for insufficient OxPhos. Recent evidence also shows that mitochondrial dysfunction is the initial event in the path to tumorigenesis induced by the mutated *Ras* oncogene and is closely linked to the action of the *BRAF* oncogene (83,142,143). Cells will enter their default state of proliferation following loss of respiratory control (9,141). Several cancer hallmarks can be linked to the transition from quiescence to proliferation (**Fig. 3**). Unbridled proliferation is linked to fermentation, which was the dominant form of energy metabolism during the oxygen deficient α period of earth's history (144). OxPhos insufficiency in fusion hybrids between immune cells (mostly macrophages) and cancer stem cells can underlie the ability of tumor cells to intravasate the circulation locally and to extravasate the circulation at distant sites (145,146). As macrophages are already mesenchymal and naturally capable of systemic tissue dispersion, it is not necessary to explain the phenomenon of metastasis in terms of complicated gene-linked epithelial to mesenchymal (EMT), and mesenchymal to epithelial (MET) transitions. Metastasis in our view would arise from the dysregulation of normal macrophage functions in fusion hybrids including intravasation and extravasation (146-148). All major hallmarks of cancer including genomic instability can be linked directly or indirectly to the respiratory dysfunction and the compensatory fermentation of the tumor cell.

Are mutations in the *P53* and the *Ras* genes primary or secondary causes of cancer?

Although germline or somatic mutations in the *P53* tumor suppressor gene and somatic mutations the *Ras* oncogene occur frequently in many tumor cells and cancers (149,150), it is not clear if these genes or their products are primary or secondary causes of cancer. Hwang and co-workers showed that p53 regulates mitochondrial respiration through its transcriptional target gene *Synthesis of Cytochrome c Oxidase 2* (*SCO2*) (151-153). In these studies the Warburg effect was linked directly to impaired respiration. Singh and co-workers showed that mitochondrial energy metabolism is impaired in human cancer cells containing defects in p53 (154). Huang and co-workers recently showed that the common K-Ras^{G12V} mutation causes a metabolic switch from

OxPhos to glycolysis (Warburg effect) due to mitochondrial dysfunction (**Fig. 4**). Lee and co-workers showed that transfection of human diploid cells with V^{12} Ras significantly increased damaging oxygen species in mitochondria (155), whereas Weinberg and colleagues showed that mitochondrial ROS generation and damage to complex III was essential for K-Ras-induced cell proliferation and tumorigenesis (156). Moreover, Yang and colleagues showed that H-Ras transformation of mouse fibroblasts damaged respiration thus forcing the cells into a glycolytic metabolism (157). This is notable since activated RAS has been proposed to induce MYC activity and to enhance non-hypoxic levels of HIF-1 α (64). As MYC and HIF-1 drive glycolysis, their up-regulation would be necessary to prevent senescence following respiratory impairment (18,158). As constitutive *Ras* activation is incompatible with prenatal development, a disruption of mitochondrial energy metabolism could underlie tumor formation in mice cloned from melanoma nuclei following the inadvertent expression of the *Ras* oncogene (48). Viewed collectively, these and other observations are consistent with Warburg's theory and suggest that mutations in the *P53* and *Ras* genes initiate cancer through their adverse effects on respiratory function. It will be up to each investigator to determine whether they consider these mutations as primary or secondary causes of cancer according to the Warburg's central theory (55,57,58). It is our view that all roads to the origin of cancer pass through the mitochondria (**Fig. 3**).

Can tumor somatic mutations arise as a downstream epiphenomenon of abnormal energy metabolism?

How might protracted respiratory insufficiency cause somatic mutations and the nuclear genomic instability seen in tumor cells? The integrity of the nuclear genome is dependent to a large extent on the efficiency of mitochondrial respiratory function (159). Evidence indicates that a persistent retrograde response (RTG) or mitochondrial stress response leads to abnormalities in DNA repair mechanisms and to the up-regulation of fermentation pathways (50,160-164). Oncogene up-regulation becomes essential for increased glucose and glutamine metabolism following respiratory impairment (83,165). The metabolic waste products of fermentation can destabilize the morphogenetic field of the tumor microenvironment thus contributing to inflammation, angiogenesis and

progression (166-168). Normal mitochondrial function is necessary for maintaining intracellular calcium homeostasis, which is required for chromosomal integrity and the fidelity of cell division. Aneuploidy can arise during cell division from abnormalities in calcium homeostasis (159). In this general picture, the abnormal genomic landscape seen in tumor cells is considered a downstream epiphenomenon of dysfunctional respiration and protracted oncogene-driven fermentation. In other words the somatic mutations arise as effects rather than as causes of tumorigenesis. The nuclear transfer experiments support this view (**Fig. 1**). In light of this perspective, it would be important for those working in cancer genomics field to justify the logic of their experimental approach to the cancer problem (169,170).

Cancer progression is more consistent with Lamarckian than Darwinian evolution

When viewed as a mitochondrial metabolic disease cancer progression is more in line with the evolutionary theory of Lamarck than with the theory of Darwin (20). Many investigators in the cancer field have attempted to link the Darwinian theory of evolution to the phenomenon of tumor progression (171-174). The attempt to link cancer progression to Darwinian evolution is based largely on the view that nuclear somatic mutations are drives of the disease. According to Lamarck's theory, it is the environment that produces changes in biological structures (175). Through adaptation and differential use, these changes lead to modifications in the structures. The modifications of structures would then be passed on to successive generations as acquired traits. Lamarck's evolutionary synthesis was based on his belief that the degree of use or disuse of biological structures shaped evolution along with the inheritance of acquired adaptability. Lamarck's ideas could also accommodate a dominant role for epigenetics and horizontal gene transfer as factors that could facilitate tumor progression (176,177). In addition to nuclear epigenetic events involving acetylation and phosphorylations, mitochondria are also recognized as a powerful extra nuclear epigenetic system (159,178-180). Other epigenetic phenomena such as cytomegalovirus infection, cell fusion and horizontal gene transfer can also contribute to cancer progression and metastasis (147,159,181-184).

Considering the dynamic behavior of mitochondria involving regular fusions and fissions, abnormalities in mitochondrial structure can be rapidly disseminated throughout the cellular mitochondrial network and passed along to daughter cells somatically, through cytoplasmic inheritance (17,185). The capacity for mitochondrial respiratory function becomes progressively less with each cell division as adaptability to substrate level phosphorylation increases (**Fig. 3**). The somatic progression of cancer would therefore embody the concept of the somatic inheritance of an acquired trait. The acquired trait in this case is alteration to mitochondrial structure. The most malignant cancer cells would sustain the near-complete replacement of their respiration with fermentation. This is obvious in those tumor cells with quantitative and qualitative abnormalities in their mitochondria (**Fig. 2**). The somatic inheritance of mitochondrial dysfunction in tumor cells could contribute in part to the appearance of a clonal origin, but not directly involving the nuclear genome. However, the degree of nuclear genomic instability can be linked to mitochondrial dysfunction and both defects together can contribute to tumor progression. A Lamarckian view can account for the non-uniform accumulation of mutations and drug resistance seen during cancer progression. Drug resistance is linked to enhanced lactate fermentation, which is acquired during tumor progression (61,186). It is our opinion that the evolutionary concepts of Lamarck can better explain the phenomena of tumor progression than can the evolutionary concepts of Darwin. We encourage further research on this perspective of tumor progression.

Exploiting mitochondrial dysfunction for the metabolic management of cancer

If cancer is primarily a disease of energy metabolism, then rational strategies for cancer management should be found in those therapies that specifically target tumor cell energy metabolism. These therapeutic strategies should be applicable to the majority of cancers regardless of tissue origin, as nearly all cancers suffer from a common malady, i.e., insufficient respiration with compensatory fermentation (2,54,55,57). As glucose is the major fuel for tumor energy metabolism through lactate fermentation, the restriction of glucose becomes a prime target for management. However, most normal cells of the body also need glycolytic pathway products, such as pyruvate, for energy production through OxPhos. It therefore becomes important to protect normal cells from drugs or

therapies that disrupt glycolytic pathways or cause systemic reduction of glucose. It is well known that ketones can replace glucose as an energy metabolite and can protect the brain from severe hypoglycemia (187-189). Hence, the shift in energy metabolism associated with a low carbohydrate, high fat ketogenic diet administered in restricted amounts (KD-R) can protect normal cells from glycolytic inhibition and the brain from hypoglycemia.

When systemic glucose availability becomes limiting, most normal cells of the body will transition their energy metabolism to fats and ketone bodies. Ketone bodies are generated almost exclusively in liver hepatocytes largely from fatty acids of triglyceride origin during periods of fasting (187,190). There are no metabolic pathways described that can produce ketone bodies from carbohydrates despite suggestions to the contrary (191). A restriction of total caloric intake will facilitate a reduction in blood glucose and insulin levels and an elevation in ketone bodies (β -hydroxybutyrate and acetoacetate). Most tumor cells are unable to use ketone bodies for energy due to abnormalities in mitochondria structure or function (13,192). Ketone bodies can also be toxic to some cancer cells (193,194). Nutritional ketosis induces metabolic stress on tumor tissue that is selectively vulnerable to glucose deprivation (13). Hence, metabolic stress will be greater in tumor cells than in normal cells when the whole body is transitioned away from glucose and to ketone bodies for energy.

The metabolic shift from glucose metabolism to ketone body metabolism creates an anti-angiogenic, anti-inflammatory, and pro-apoptotic environment within the tumor mass (192,195-199). The general concept of a survival advantage of tumor cells over normal cells occurs when fermentable fuels are abundant, but not when they are limited (20). **Fig. 5** illustrates the changes in whole body levels of blood glucose and ketone bodies (β -hydroxybutyrate) that will metabolically stress tumor cells while enhancing the metabolic efficiency of normal cells. This therapeutic strategy was illustrated previously in cancer patients and in preclinical models (200-205).

Implications for novel therapeutics

Once the whole body enters the metabolic zone described in **Fig. 5**, relatively low doses of a variety of drugs can be used to further target energy metabolism in any surviving tumor cells (192). It is interesting that the therapeutic success of imatinib (Gleevec) and trastuzumab (Herceptin) in managing BCR-ABL leukemia cells and ErbB2-positive breast cancers, respectively, is dependent on their ability to target signaling pathways linked to glucose metabolism (206,207). In contrast to these drugs, which target energy metabolism primarily in those individuals with mutations in specific receptors linked to the IGF-1/PI3K/Akt pathway, calorie restricted ketogenic diets will target similar pathways in any cancer cell regardless of the mutations involved (197,208). Dietary energy reduction will simultaneously target multiple metabolic signaling pathways without causing adverse effects or toxicity (208). Non-toxic metabolic therapies might also be a preferable alternative to toxic immunotherapies for cancer management especially if both therapies target the same pathways. It must be emphasized that the therapeutic efficacy of the KD is strongly dependent on restricted intake, as consumption of the KD in unrestricted amounts can cause insulin insensitivity and glucose elevation despite the complete absence of carbohydrates in the diet (205). Elevated consumption of the KD is not often seen, however, as humans usually restrict intake due to the high fat content of the diet.

Poff et al also recently showed a synergistic interaction between the KD and hyperbaric oxygen therapy (HBO₂T) (**Fig. 6**). The KD reduces glucose for glycolytic energy, while also reducing NADPH levels for anti-oxidant potential through the pentose-phosphate-pathway. HBO₂T will increase ROS in the tumor cells while the ketones will protect normal cells against ROS damage and from the potential for central nervous system oxygen toxicity (189,209). Glucose deprivation will enhance oxidative stress in tumor cells, while increased oxygen can reduce tumor cell proliferation (210,211). A dependency on glucose and an inability to use ketones for energy makes tumor cells selectively vulnerable to this therapy. Although this metabolic therapy is effective against those tumor cells that contain mitochondria, it remains to be determined if this therapy would be equally effective against those tumor cells containing few or no mitochondria (51). In contrast to radiation therapy, which also kills tumor cells through

ROS production (212), the KD + HBO₂T will kill tumor cells without causing toxic collateral damage to normal cells. Cancer patients and their oncologists should know about this. Some ketogenic diets might also enhance the therapeutic action of radiation therapy against brain and lung tumors (213,214). It will be important to compare and contrast the therapeutic efficacy of conventional radiation therapy with HBO₂T when used with the KD-R. While radiation is widely used as a cancer therapy, it should be recognized that radiation damages respiration in normal cells and can itself cause cancer (55). Radiation therapy for malignant brain cancer creates a necrotic microenvironment that can facilitate recurrence and progression through enhanced glucose and glutamine metabolism (13,215).

Besides drugs that target glucose, drugs that target glutamine can also be effective in killing systemic metastatic cancer cells (192,216,217). Many metastatic cancers express multiple characteristics of macrophages (146,218). Glutamine is a major fuel of macrophages and other cells of the immune system (146,219). As glutamine is the most abundant amino acid in the body and is used in multiple metabolic reactions, targeting glutamine without toxicity might be more difficult than targeting glucose (220,221). While glutamine interacts synergistically with glucose to drive energy metabolism in cultured tumor cells, there are reports suggesting that glutamine can have chemo-preventive effects (222). Further studies are needed to evaluate the role of glutamine as a facilitator of tumor energy metabolism *in vivo*.

The novelty of the metabolic approach to cancer management involves the implementation of a synergistic combination of nutritional ketosis, cancer metabolic drugs, and HBO₂T. This therapeutic approach would be similar to the “Press-Pulse” scenario for the mass extinction of organisms in ecological communities (223,224). The KD-R would act as a sustained “Press”, while HBO₂T and metabolic drugs would act as a “Pulse” for the mass elimination of tumor cells in the body. Some of the cancer metabolic drugs could include 2-deoxyglucose, 3-bromopyruvate, and dichloroacetate (56,120,225-227). This therapeutic strategy produces a shift in metabolic physiology that will not only kill tumor cells, but will also enhance the general health and metabolic

1
2
3 efficiency of normal cells, and consequently the whole body (189,209). We view this
4 therapeutic approach as a type of “*mitochondrial enhancement therapy*” (192). As we
5 consider OxPhos insufficiency with compensatory fermentation as the origin of cancer,
6 enhanced OxPhos efficiency would be anti-carcinogenic.
7
8
9

10
11
12 Many cancers are infected with human cytomegalovirus (HCMV), which acts as an
13 oncomodulator of tumor progression (228). Products of the virus can damage
14 mitochondria in the infected tumor cells thus contributing to a further dependence on
15 glucose and glutamine for energy metabolism (18,229-231). The virus often infects
16 cells of monocyte/macrophage origin, which are considered the origin of many
17 metastatic cancers (145,146,232,233). We predict that the KD-R used together with
18 anti-viral therapy will also be an effective Press-Pulse strategy for reducing progression
19 of those cancers infected with HCMV (234).
20
21
22
23
24
25
26
27

28 Advanced metastatic cancers can become manageable when their access to
29 fermentable fuels becomes restricted. The metabolic shift associated with the KD-R
30 involves “keto-adaptation”. However, the adaptation to this new metabolic state can be
31 challenging for some people. The administration of ketone esters could conceivably
32 enable patients to circumvent the dietary restriction generally required for sustained
33 nutritional ketosis. Ketone ester-induced ketosis would make sustained hypoglycemia
34 more tolerable, and thus assist in metabolic management of cancer (235,236). As each
35 person is a unique metabolic entity, personalization of metabolic therapy as a broad
36 based cancer treatment strategy will require fine-tuning based on an understanding of
37 individual human physiology. Also, personalized molecular therapies developed
38 through the genome projects could be useful in targeting and killing those tumor cells
39 that might survive the non-toxic whole-body metabolic therapy. The number of
40 molecular targets should be less in a few survivor cells of a small tumor than in a
41 heterogeneous cell population of a large tumor. We would therefore consider
42 personalized molecular therapy as a final strategy rather than as an initial strategy for
43 cancer management. Non-toxic metabolic therapy should become the future of cancer
44 treatment if the goal is to manage the disease without harming the patient. While it will
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

be important for researchers to elucidate the mechanistic minutia responsible for the therapeutic benefits, this should not impede an immediate application of this therapeutic strategy for cancer management or prevention.

Acknowledgements: This work was supported, in part, by the National Institutes of Health (HD-39722, NS-55195 and CA-102135) a grant from the American Institute of Cancer Research, and the Boston College Expense Fund (TNS). The work was also funded in part by Scivation and Office of Naval Research (ONR) (DPA). We also that Miriam Kalamian for helpful comments.

Legends to Figures

Figure 1. Role of the nucleus and mitochondria in the origin of tumors. This image summarizes the experimental evidence supporting a dominant role of the mitochondria in the origin of tumorigenesis as previously described (49). Normal cells are depicted in green with mitochondrial and nuclear morphology indicative of normal respiration and nuclear gene expression, respectively. Tumor cells are depicted in red with abnormal mitochondrial and nuclear morphology indicative of abnormal respiration and genomic instability. 1) Normal cells beget normal cells. 2) Tumor cells beget tumor cells. 3) Delivery of a tumor cell nucleus into a normal cell cytoplasm begets normal cells despite the persistence of tumor-associated genomic abnormalities. 4) Delivery of a normal cell nucleus into a tumor cell cytoplasm begets tumor cells or dead cells, but not normal cells. The results suggest that tumors do not arise from nuclear genomic defects alone, and that normal mitochondria can suppress tumorigenesis. Original diagram from Jeffrey Ling and Thomas N. Seyfried, with permission.

Figure 2. Typical ultrastructure of a normal mitochondrion and a mitochondrion from a human glioblastoma. Normal mitochondria contain elaborate cristae, which are extensions of the inner membrane and contain the protein complexes of the electron transport chain necessary for producing ATP through OxPhos. The mitochondrion from the glioblastoma (m) is enlarged and shows a near total breakdown of cristae (cristolysis) and an electron-lucent matrix. The absence of cristae in glioblastoma mitochondria indicates that OxPhos would be deficient. The arrow indicates an inner membrane fold. Bar: 0.33 μm . Method of staining: uranyl acetate/lead citrate. The GBM mitochondrion was reprinted with permission from *Journal of Electron Microscopy* (94). The normal mitochondrion diagram was from <http://academic.brooklyn.cuny.edu/biology/bio4fv/page/mito.htm>.

Figure 3. Mitochondrial respiratory dysfunction as the origin of cancer. Cancer can arise from any number of non-specific events that damage the respiratory capacity of cells over time. The path to carcinogenesis will occur only in those cells capable of enhancing energy production through fermentation (substrate level phosphorylation, SLP). Despite the shift from respiration to SLP the $\Delta G'$ of ATP hydrolysis remains fairly constant at approximately -56 kJ indicating that the energy from SLP compensates for the reduced energy from OxPhos. The mitochondrial stress response or retrograde signaling (RTG) will initiate oncogene up-regulation and tumor suppressor gene inactivation that are necessary to maintain viability of incipient cancer cells when respiration becomes unable to maintain energy homeostasis. Genomic instability will arise as a secondary consequence of protracted mitochondrial stress from disturbances in the intracellular and extracellular microenvironment. Metastasis arises from respiratory damage in cells of myeloid/macrophage origin (146). The degree of malignancy is linked directly to the energy transition from OxPhos to substrate level phosphorylation. This scenario links all major cancer hallmarks to an extrachromosomal respiratory dysfunction (141). The **T** signifies an arbitrary threshold when the shift from OxPhos to SLP might become irreversible. Reprinted with modifications from (17).

Figure 4. Time line of events following expression of *K-Ras*. The time axis depicts the various events after stimulation of the *K-RAS* gene expression. The findings of Huang and co-workers indicate that mitochondria-linked changes are observed around the time of the increase in the K-Ras protein (142). This is then followed by other changes, such as alteration in cell metabolism. Gene mutations would form as a downstream epiphenomenon of altered metabolism. Malignant transformation, documented by the colony-forming activity of the cells and their propensity to form tumors ensue much later. According to these findings the Warburg effect (aerobic fermentation), arises as a consequence of *K-RAS*-induced respiratory injury. This timeline is in general agreement with Warburg's central theory and with other similar findings that respiratory disturbance is an initial event in *K-RAS*-induced tumorigenesis (237-239). The timeline will be greatly protracted *in vivo* as shown from the Roskelley et al., experiments (240).

Reprinted from Neuzil et al with permission (143).

Figure 5. Relationship of circulating levels of glucose and ketones (β -hydroxybutyrate, β -OHB) to tumor management. The glucose and ketone values are within normal physiological ranges under fasting conditions in humans and will produce anti-angiogenic, anti-inflammatory, and pro-apoptotic effects. We refer to this state as the zone of metabolic management. Metabolic stress will be greater in tumor cells than in normal cells when the whole body enters the metabolic zone. The values for blood glucose in mg/dl can be estimated by multiplying the mM values by 18. The glucose and ketone levels predicted for tumor management in human cancer patients are 3.1-3.8 mM (55-65 mg/dl) and 2.5-7.0 mM, respectively. These ketone levels are well below the levels associated with ketoacidosis (blood ketone values greater than 15 mmol). Elevated ketones will protect the brain from hypoglycemia. Modified from a previous version (241).

Figure 6.

The ketogenic diet (KD) and hyperbaric oxygen therapy (HBO₂T) are synergistic in reducing systemic metastatic cancer in the syngeneic VM mouse model. VM-M3/Fluc tumor cells were implanted subcutaneously and systemic organ metastasis was evaluated *ex vivo* using bioluminescent imaging as previously described (242,243). Tumor growth was slower in mice fed the KD than in mice fed a standard high carbohydrate diet (SD). (A) Representative animals from each treatment group demonstrating tumor bioluminescence at day 21 after tumor cell inoculation. Treated animals showed less bioluminescence than controls with KD + HBO₂T mice exhibiting a profound decrease in tumor bioluminescence compared to all groups. (B) Total body bioluminescence was measured weekly as a measure of tumor size; error bars represent standard error of the mean. KD+HBO₂T mice exhibited significantly less tumor bioluminescence than control animals at week 3 ($p < 0.01$; two-tailed student's t-test) and an overall trend of notably slower tumor growth than controls and other treated animals throughout the study. (C & D) Day 21 organ bioluminescence of SD and KD +

HBO₂T animals (N = 8) demonstrated a trend of reduced metastatic tumor burden in animals receiving the combined therapy. Spleen bioluminescence was significantly decreased in KD + HBO₂T mice (*p < 0.02; two-tailed student's t-test). Reprinted with permission from (243).

References

1. Sporn, M.B. (1996) The war on cancer. *Lancet*, **347**, 1377-81.
2. Seyfried, T.N. (2012) *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken.
3. Fidler, I.J. (2003) The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat Rev Cancer*, **3**, 453-8.
4. Lazebnik, Y. (2010) What are the hallmarks of cancer? *Nat Rev Cancer*, **10**, 232-3.
5. Tarin, D. (2011) Cell and tissue interactions in carcinogenesis and metastasis and their clinical significance. *Semin Cancer Biol*, **21**, 72-82.
6. Seyfried, T.N. (2012) Confusion surrounds the origin of cancer. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 15-29.
7. Hanahan, D., et al. (2011) Hallmarks of cancer: the next generation. *Cell*, **144**, 646-74.
8. Baker, S.G., et al. (2007) Paradoxes in carcinogenesis: new opportunities for research directions. *BMC Cancer*, **7**, 151.
9. Soto, A.M., et al. (2004) The somatic mutation theory of cancer: growing problems with the paradigm? *Bioessays*, **26**, 1097-107.
10. Vogelstein, B., et al. (2013) Cancer genome landscapes. *Science*, **339**, 1546-58.
11. Alexandrov, L.B., et al. (2013) Signatures of mutational processes in human cancer. *Nature*, **500**, 415-21.
12. Soto, A.M., et al. (2012) Is systems biology a promising approach to resolve controversies in cancer research? *Cancer cell international*, **12**, 12.
13. Seyfried, T.N., et al. (2012) Is the restricted ketogenic diet a viable alternative to the standard of care for managing malignant brain cancer? *Epilepsy research*, **100**, 310-26.
14. Seyfried, T.N. (2013) Cancer as a metabolic disease: Implications for novel therapeutics. In *Education Book*. Amer. Assoc. Cancer Res.
15. Seyfried, T.N., et al. (2010) Metabolic management of brain cancer. *Biochim Biophys Acta*, **1807**, 577-94.
16. Seyfried, T.N., et al. (2005) Targeting energy metabolism in brain cancer: review and hypothesis. *Nutr Metab (Lond)*, **2**, 30.
17. Seyfried, T.N., et al. (2010) Cancer as a metabolic disease. *Nutr Metab (Lond)*, **7**, 7.
18. Seyfried, T.N. (2012) Genes, respiration, viruses, and cancer. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 145-176.

19. Stratton, M.R. (2011) Exploring the genomes of cancer cells: progress and promise. *Science*, **331**, 1553-8.
20. Seyfried, T.N. (2012) Nothing in cancer biology makes sense except in the light of evolution. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 261-275.
21. Nowell, P.C. (1976) The clonal evolution of tumor cell populations. *Science*, **194**, 23-8.
22. Fojo, T., *et al.* (2010) Biologically targeted cancer therapy and marginal benefits: are we making too much of too little or are we achieving too little by giving too much? *Clin Cancer Res*, **16**, 5972-80.
23. Rosell, R., *et al.* (2009) Customized treatment in non-small-cell lung cancer based on EGFR mutations and BRCA1 mRNA expression. *PLoS One*, **4**, e5133.
24. McLeod, H.L. (2013) Cancer pharmacogenomics: early promise, but concerted effort needed. *Science*, **339**, 1563-6.
25. Hardy, P.A., *et al.* (2005) Reappraisal of the Hanseemann-Boveri hypothesis on the origin of tumors. *Cell biology international*, **29**, 983-92.
26. Gibbs, W.W. (2003) Untangling the roots of cancer. *Sci Am*, **289**, 56-65.
27. Hameroff, S.R. (2004) A new theory of the origin of cancer: quantum coherent entanglement, centrioles, mitosis, and differentiation. *Biosystems*, **77**, 119-36.
28. Manchester, K. (1997) The quest by three giants of science for an understanding of cancer. *Endeavour*, **21**, 72-6.
29. Wolf, U. (1974) Theodor Boveri and his book, *On the Problem of the Origin of Malignant Tumors*. In Sons, J.W. (ed.), *Chromosomes and Cancer*, pp. 1-20.
30. Duesberg, P., *et al.* (2000) Aneuploidy, the somatic mutation that makes cancer a species of its own. *Cell Motil Cytoskeleton*, **47**, 81-107.
31. Salk, J.J., *et al.* (2010) Mutational heterogeneity in human cancers: origin and consequences. *Annu Rev Pathol*, **5**, 51-75.
32. Cairns, J. (1981) The origin of human cancers. *Nature*, **289**, 353-7.
33. Loeb, L.A. (2001) A mutator phenotype in cancer. *Cancer Res*, **61**, 3230-9.
34. Whitman, R.C. (1919) Somatic mutations as a factor in the production of cancer; a critical review of von Hanseemann's theory of anaplasia in light of modern knowledge of genetics. *J. Cancer Res.*, **4**, 181-202.
35. Nigro, J.M., *et al.* (1989) Mutations in the p53 gene occur in diverse human tumour types. *Nature*, **342**, 705-8.
36. Fearon, E.R., *et al.* (1990) A genetic model for colorectal tumorigenesis. *Cell*, **61**, 759-67.
37. Knudson, A.G. (2002) Cancer genetics. *Am J Med Genet*, **111**, 96-102.
38. Wagner, R.P. (1999) Anecdotal, historical and critical commentaries on genetics. Rudolph Virchow and the genetic basis of somatic ecology. *Genetics*, **151**, 917-20.
39. Darlington, C.D. (1948) The plasmagene theory of the origin of cancer. *British journal of cancer*, **2**, 118-26.
40. Koura, M., *et al.* (1982) Suppression of tumorigenicity in interspecific reconstituted cells and cybrids. *Gann*, **73**, 574-80.
41. Israel, B.A., *et al.* (1987) Cytoplasmic suppression of malignancy. *In Vitro Cell Dev Biol*, **23**, 627-32.
42. Shay, J.W., *et al.* (1988) Cytoplasmic suppression of tumorigenicity in reconstructed mouse cells. *Cancer Res*, **48**, 830-3.

43. Howell, A.N., *et al.* (1978) Tumorigenicity and its suppression in cybrids of mouse and Chinese hamster cell lines. *Proc Natl Acad Sci U S A*, **75**, 2358-62.

44. Jonasson, J., *et al.* (1977) The analysis of malignancy by cell fusion. VIII. Evidence for the intervention of an extra-chromosomal element. *J Cell Sci*, **24**, 255-63.

45. McKinnell, R.G., *et al.* (1969) Transplantation of pluripotential nuclei from triploid frog tumors. *Science*, **165**, 394-6.

46. Mintz, B., *et al.* (1975) Normal genetically mosaic mice produced from malignant teratocarcinoma cells. *Proc Natl Acad Sci U S A*, **72**, 3585-9.

47. Li, L., *et al.* (2003) Mouse embryos cloned from brain tumors. *Cancer Res*, **63**, 2733-6.

48. Hochedlinger, K., *et al.* (2004) Reprogramming of a melanoma genome by nuclear transplantation. *Genes Dev*, **18**, 1875-85.

49. Seyfried, T.N. (2012) Mitochondria: The ultimate tumor suppressor. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 195-205.

50. Kaiparettu, B.A., *et al.* (2013) Crosstalk from non-cancerous mitochondria can inhibit tumor properties of metastatic cells by suppressing oncogenic pathways. *PloS one*, **8**, e61747.

51. Elliott, R.L., *et al.* (2012) Mitochondria organelle transplantation: introduction of normal epithelial mitochondria into human cancer cells inhibits proliferation and increases drug sensitivity. *Breast cancer research and treatment*, **136**, 347-54.

52. Israel, B.A., *et al.* (1988) Cytoplasmic mediation of malignancy. *In Vitro Cell Dev Biol*, **24**, 487-90.

53. Petros, J.A., *et al.* (2005) mtDNA mutations increase tumorigenicity in prostate cancer. *Proc Natl Acad Sci U S A*, **102**, 719-24.

54. Warburg, O. (1931) *The Metabolism of Tumours*. Richard R. Smith, New York.

55. Warburg, O. (1956) On the origin of cancer cells. *Science*, **123**, 309-14.

56. Pedersen, P.L. (2007) Warburg, me and Hexokinase 2: Multiple discoveries of key molecular events underlying one of cancers' most common phenotypes, the "Warburg Effect", i.e., elevated glycolysis in the presence of oxygen. *J Bioenerg Biomembr*, **39**, 211-22.

57. Warburg, O. (1956) On the respiratory impairment in cancer cells. *Science*, **124**, 269-270.

58. Warburg, O. (1969) Revidsed Lindau Lectures: The prime cause of cancer and prevention - Parts 1 & 2. In Burk, D. (ed.), *Meeting of the Nobel-Laureates* K. Triltsch, Lindau, Lake Constance, Germany, pp. <http://www.hopeforcancer.com/OxyPlus.htm>.

59. Racker, E. (1972) Bioenergetics and the problem of tumor growth. *American scientist*, **60**, 56-63.

60. Nelson, D.L., *et al.* (2008) *Lehninger Principles of Biochemistry* W. H. Freeman, New York.

61. Cuezva, J.M., *et al.* (2002) The bioenergetic signature of cancer: a marker of tumor progression. *Cancer Res*, **62**, 6674-81.

62. Acebo, P., *et al.* (2009) Cancer abolishes the tissue type-specific differences in the phenotype of energetic metabolism. *Transl Oncol*, **2**, 138-45.

63. Weinhouse, S. (1956) On respiratory impairment in cancer cells. *Science*, **124**, 267-9.

64. Koppenol, W.H., *et al.* (2011) Otto Warburg's contributions to current concepts of cancer metabolism. *Nat Rev Cancer*, **11**, 325-37.

65. Stellingwerff, T., *et al.* (2006) Hyperoxia decreases muscle glycogenolysis, lactate production, and lactate efflux during steady-state exercise. *American journal of physiology. Endocrinology and metabolism*, **290**, E1180-90.
66. Gatenby, R.A., *et al.* (2004) Why do cancers have high aerobic glycolysis? *Nat Rev Cancer*, **4**, 891-9.
67. Pedersen, P.L. (1978) Tumor mitochondria and the bioenergetics of cancer cells. *Prog Exp Tumor Res*, **22**, 190-274.
68. Chevrollier, A., *et al.* (2005) ANT2 expression under hypoxic conditions produces opposite cell-cycle behavior in 143B and HepG2 cancer cells. *Molecular carcinogenesis*, **42**, 1-8.
69. Wijburg, F.A., *et al.* (1989) Studies on the formation of lactate and pyruvate from glucose in cultured skin fibroblasts: implications for detection of respiratory chain defects. *Biochemistry international*, **19**, 563-70.
70. Tiefenthaler, M., *et al.* (2001) Increased lactate production follows loss of mitochondrial membrane potential during apoptosis of human leukaemia cells. *British journal of haematology*, **114**, 574-80.
71. Donnelly, M., *et al.* (1976) Energy metabolism in respiration-deficient and wild type Chinese hamster fibroblasts in culture. *J Cell Physiol*, **89**, 39-51.
72. Fiske, B.P., *et al.* (2012) Seeing the Warburg effect in the developing retina. *Nature cell biology*, **14**, 790-1.
73. Prichard, J., *et al.* (1991) Lactate rise detected by ¹H NMR in human visual cortex during physiologic stimulation. *Proceedings of the National Academy of Sciences of the United States of America*, **88**, 5829-31.
74. Fox, P.T., *et al.* (1988) Nonoxidative glucose consumption during focal physiologic neural activity. *Science*, **241**, 462-4.
75. Krasnow, N., *et al.* (1962) Myocardial lactate and pyruvate metabolism. *The Journal of clinical investigation*, **41**, 2075-85.
76. Burk, D., *et al.* (1956) On respiratory impairment in cancer cells. *Science*, **124**, 270-2.
77. Burk, D., *et al.* (1967) On the significance of glucolysis for cancer growth, with special reference to Morris rat hepatomas. *J Natl Cancer Inst*, **38**, 839-63.
78. Vaupel, P., *et al.* (2012) Availability, not respiratory capacity governs oxygen consumption of solid tumors. *The international journal of biochemistry & cell biology*, **44**, 1477-81.
79. Moreno-Sanchez, R., *et al.* (2007) Energy metabolism in tumor cells. *Febs J*, **274**, 1393-418.
80. Haq, R., *et al.* (2013) Oncogenic BRAF regulates oxidative metabolism via PGC1alpha and MITF. *Cancer cell*, **23**, 302-15.
81. Weinhouse, S. (1976) The Warburg hypothesis fifty years later. *Z Krebsforsch Klin Onkol Cancer Res Clin Oncol*, **87**, 115-26.
82. Ferreira, L.M. (2010) Cancer metabolism: the Warburg effect today. *Exp Mol Pathol*, **89**, 372-80.
83. Hall, A., *et al.* (2013) Dysfunctional oxidative phosphorylation makes malignant melanoma cells addicted to glycolysis driven by the V600EBRAF oncogene. *Oncotarget*, **4**, 584-99.
84. Hochachka, P.W., *et al.* (2002) *Biochemical Adaptation: Mechanism and Process in Physiological Evolution*. Oxford Press, New York.

85. Seyfried, T.N. (2012) Is respiration normal in cancer cells? In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 119-132.

86. Ramanathan, A., *et al.* (2005) Perturbational profiling of a cell-line model of tumorigenesis by using metabolic measurements. *Proc Natl Acad Sci U S A*, **102**, 5992-7.

87. Seyfried, T.N. (2012) Is mitochondrial glutamine fermentation a missing link in the metabolic theory of cancer? In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 133-144.

88. Chinopoulos, C., *et al.* (2010) Forward operation of adenine nucleotide translocase during F0F1-ATPase reversal: critical role of matrix substrate-level phosphorylation. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **24**, 2405-16.

89. Phillips, D., *et al.* (2009) Succinyl-CoA synthetase is a phosphate target for the activation of mitochondrial metabolism. *Biochemistry*, **48**, 7140-9.

90. Schwimmer, C., *et al.* (2005) Increasing mitochondrial substrate-level phosphorylation can rescue respiratory growth of an ATP synthase-deficient yeast. *J Biol Chem*, **280**, 30751-9.

91. Seyfried, T.N. (2012) Respiratory dysfunction in cancer cells. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 73-105.

92. Arismendi-Morillo, G. (2011) Electron microscopy morphology of the mitochondrial network in gliomas and their vascular microenvironment. *Biochimica et biophysica acta*, **1807**, 602-8.

93. Arismendi-Morillo, G. (2009) Electron microscopy morphology of the mitochondrial network in human cancer. *Int J Biochem Cell Biol*, **41**, 2062-8.

94. Arismendi-Morillo, G.J., *et al.* (2008) Ultrastructural mitochondrial pathology in human astrocytic tumors: potentials implications pro-therapeutics strategies. *J Electron Microsc (Tokyo)*, **57**, 33-9.

95. Galluzzi, L., *et al.* (2010) Mitochondrial gateways to cancer. *Molecular aspects of medicine*, **31**, 1-20.

96. Kiebish, M.A., *et al.* (2009) In vitro growth environment produces lipidomic and electron transport chain abnormalities in mitochondria from non-tumorigenic astrocytes and brain tumours. *ASN Neuro*, **1**.

97. Kiebish, M.A., *et al.* (2008) Cardiolipin and electron transport chain abnormalities in mouse brain tumor mitochondria: lipidomic evidence supporting the Warburg theory of cancer. *J Lipid Res*, **49**, 2545-56.

98. Gonzalez, M.J., *et al.* (2012) The bio-energetic theory of carcinogenesis. *Medical hypotheses*, **79**, 433-9.

99. Benard, G., *et al.* (2008) Ultrastructure of the mitochondrion and its bearing on function and bioenergetics. *Antioxidants & redox signaling*, **10**, 1313-42.

100. Shapovalov, Y., *et al.* (2011) Mitochondrial dysfunction in cancer cells due to aberrant mitochondrial replication. *J Biol Chem*.

101. Alirol, E., *et al.* (2006) Mitochondria and cancer: is there a morphological connection? *Oncogene*, **25**, 4706-16.

102. Oudard, S., *et al.* (1997) Gliomas are driven by glycolysis: putative roles of hexokinase, oxidative phosphorylation and mitochondrial ultrastructure. *Anticancer Res*, **17**, 1903-11.

103. Wu, S.B., *et al.* (2012) AMPK-mediated increase of glycolysis as an adaptive response to oxidative stress in human cells: implication of the cell survival in mitochondrial diseases. *Biochimica et biophysica acta*, **1822**, 233-47.
104. Fulda, S., *et al.* (2010) Targeting mitochondria for cancer therapy. *Nature reviews. Drug discovery*, **9**, 447-64.
105. Shapovalov, Y., *et al.* (2011) Mitochondrial dysfunction in cancer cells due to aberrant mitochondrial replication. *The Journal of biological chemistry*, **286**, 22331-8.
106. Chance, B., *et al.* (1959) Spectroscopic evidence of metabolic control. *Science*, **129**, 700-8.
107. Colowick, S.P. (1961) The status of Warburg's theory of glycolysis and respiration in tumors. *Quart. Rev. Biol.*, vol. 36, pp. 273-276.
108. Cairns, R.A., *et al.* (2011) Regulation of cancer cell metabolism. *Nature reviews. Cancer*, **11**, 85-95.
109. Ward, P.S., *et al.* (2012) Metabolic reprogramming: a cancer hallmark even warburg did not anticipate. *Cancer cell*, **21**, 297-308.
110. Vander Heiden, M.G., *et al.* (2009) Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*, **324**, 1029-33.
111. Rossignol, R., *et al.* (2004) Energy substrate modulates mitochondrial structure and oxidative capacity in cancer cells. *Cancer Res*, **64**, 985-93.
112. Chevrollier, A., *et al.* (2011) Adenine nucleotide translocase 2 is a key mitochondrial protein in cancer metabolism. *Biochimica et biophysica acta*, **1807**, 562-7.
113. Jahnke, V.E., *et al.* (2010) Evidence for mitochondrial respiratory deficiency in rat rhabdomyosarcoma cells. *PLoS One*, **5**, e8637.
114. Schild, L., *et al.* (2012) Composition of molecular cardiolipin species correlates with proliferation of lymphocytes. *Experimental biology and medicine*, **237**, 372-9.
115. Kocherginsky, N. (2009) Acidic lipids, H(+)-ATPases, and mechanism of oxidative phosphorylation. Physico-chemical ideas 30 years after P. Mitchell's Nobel Prize award. *Prog Biophys Mol Biol*, **99**, 20-41.
116. Claypool, S.M., *et al.* (2012) The complexity of cardiolipin in health and disease. *Trends in biochemical sciences*, **37**, 32-41.
117. Zu, X.L., *et al.* (2004) Cancer metabolism: facts, fantasy, and fiction. *Biochem Biophys Res Commun*, **313**, 459-65.
118. Wang, T., *et al.* (1976) Aerobic glycolysis during lymphocyte proliferation. *Nature*, **261**, 702-5.
119. Lunt, S.Y., *et al.* (2011) Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annual review of cell and developmental biology*, **27**, 441-64.
120. Papandreou, I., *et al.* (2011) Anticancer drugs that target metabolism: Is dichloroacetate the new paradigm? *International journal of cancer. Journal international du cancer*, **128**, 1001-8.
121. Burk, D., *et al.* (1941) Metabolism of butter yellow rat liver cancers. *Canc. Res.*, **1**, 733-734.
122. Simek, J., *et al.* (1965) Effect of glucose administered in vivo or in vitro on the respiratory quotient of rat liver tissue after partial hepatectomy. *Nature*, **207**, 761-2.
123. Diatlovitskaia, E.V., *et al.* (1972) [Cardiolipins of mitochondria and microsomes of Jensen's sarcoma]. *Dokl Akad Nauk SSSR*, **206**, 737-9.

1
2
3
4 124. Diatlovitskaia, E.V., *et al.* (1976) [Positional distribution of fatty acids in the
5 phospholipids of regenerating rat liver]. *Biokhimiia*, **41**, 538-42.
6 125. Crabtree, H.G. (1929) Observations on the carbohydrate metabolism of tumors. *Biochem.*
7 *J.*, **23**, 536-545
8 126. Redman, E.K., *et al.* (2013) Role of p90(RSK) in regulating the Crabtree effect:
9 implications for cancer. *Biochemical Society transactions*, **41**, 124-6.
10 127. Diaz-Ruiz, R., *et al.* (2008) Mitochondrial oxidative phosphorylation is regulated by
11 fructose 1,6-bisphosphate. A possible role in Crabtree effect induction? *The Journal of*
12 *biological chemistry*, **283**, 26948-55.
13 128. Diaz-Ruiz, R., *et al.* (2011) The Warburg and Crabtree effects: On the origin of cancer
14 cell energy metabolism and of yeast glucose repression. *Biochim Biophys Acta*.
15 129. Holleran, A.L., *et al.* (1995) Glutamine metabolism in AS-30D hepatoma cells. Evidence
16 for its conversion into lipids via reductive carboxylation. *Molecular and cellular*
17 *biochemistry*, **152**, 95-101.
18 130. Fendt, S.M., *et al.* (2013) Reductive glutamine metabolism is a function of the alpha-
19 ketoglutarate to citrate ratio in cells. *Nature communications*, **4**, 2236.
20 131. Pollard, P.J., *et al.* (2003) The TCA cycle and tumorigenesis: the examples of fumarate
21 hydratase and succinate dehydrogenase. *Ann Med*, **35**, 632-9.
22 132. Gottlieb, E., *et al.* (2005) Mitochondrial tumour suppressors: a genetic and biochemical
23 update. *Nat Rev Cancer*, **5**, 857-66.
24 133. Seyfried, T.N. (2012) Cancer Models. In *Cancer as a Metabolic Disease: On the Origin,*
25 *Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 31-46.
26 134. Ecsedy, J.A., *et al.* (1999) Expression of mouse sialic acid on gangliosides of a human
27 glioma grown as a xenograft in SCID mice. *J. Neurochem*, **73**, 254-259.
28 135. Chou, H.H., *et al.* (1998) A mutation in human CMP-sialic acid hydroxylase occurred
29 after the Homo-Pan divergence. *Proc. Natl. Acad. Sci. USA*, **95**, 11751-11756.
30 136. Martin, M.J., *et al.* (2005) Human embryonic stem cells express an immunogenic
31 nonhuman sialic acid. *Nat Med*, **11**, 228-32.
32 137. Davies, B., *et al.* (1993) Physiological parameters in laboratory animals and humans.
33 *Pharmaceutical research*, **10**, 1093-5.
34 138. Mahoney, L.B., *et al.* (2006) Caloric restriction in C57BL/6J mice mimics therapeutic
35 fasting in humans. *Lipids Health Dis*, **5**, 13.
36 139. Tentler, J.J., *et al.* (2012) Patient-derived tumour xenografts as models for oncology drug
37 development. *Nature reviews. Clinical oncology*, **9**, 338-50.
38 140. Chaparro, R.J., *et al.* (2006) Nonobese diabetic mice express aspects of both type 1 and
39 type 2 diabetes. *Proc Natl Acad Sci U S A*, **103**, 12475-80.
40 141. Seyfried, T.N. (2012) Mitochondrial respiratory dysfunction and the extrachromosomal
41 origin of cancer. In *Cancer as a Metabolic Disease: On the Origin, Management, and*
42 *Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 253-259.
43 142. Hu, Y., *et al.* (2012) K-ras(G12V) transformation leads to mitochondrial dysfunction and
44 a metabolic switch from oxidative phosphorylation to glycolysis. *Cell research*, **22**, 399-
45 412.
46 143. Neuzil, J., *et al.* (2012) K-Ras and mitochondria: Dangerous liaisons. *Cell research*, **22**,
47 285-7.
48 144. Szent-Gyorgyi, A. (1977) The living state and cancer. *Proc Natl Acad Sci U S A*, **74**,
49 2844-7.
50
51
52
53
54
55
56
57
58
59
60

145. Pawelek, J.M., *et al.* (2008) The cancer cell--leukocyte fusion theory of metastasis. *Adv Cancer Res*, **101**, 397-444.
146. Seyfried, T.N., *et al.* (2013) On the origin of cancer metastasis. *Critical reviews in oncogenesis*, **18**, 43-73.
147. Powell, A.E., *et al.* (2011) Fusion between Intestinal epithelial cells and macrophages in a cancer context results in nuclear reprogramming. *Cancer research*, **71**, 1497-505.
148. Pawelek, J.M. (2005) Tumour-cell fusion as a source of myeloid traits in cancer. *Lancet Oncol*, **6**, 988-93.
149. Malkin, D., *et al.* (1990) Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science*, **250**, 1233-8.
150. Funes, J.M., *et al.* (2007) Transformation of human mesenchymal stem cells increases their dependency on oxidative phosphorylation for energy production. *Proc Natl Acad Sci U S A*, **104**, 6223-8.
151. Sung, H.J., *et al.* (2011) Mitochondrial respiration protects against oxygen-associated DNA damage. *Nat Commun*, **1**, 1-8.
152. Lago, C.U., *et al.* (2010) p53, Aerobic Metabolism and Cancer. *Antioxid Redox Signal*.
153. Matoba, S., *et al.* (2006) p53 regulates mitochondrial respiration. *Science*, **312**, 1650-3.
154. Zhou, S., *et al.* (2003) Mitochondrial impairment in p53-deficient human cancer cells. *Mutagenesis*, **18**, 287-92.
155. Lee, A.C., *et al.* (1999) Ras proteins induce senescence by altering the intracellular levels of reactive oxygen species. *J Biol Chem*, **274**, 7936-40.
156. Weinberg, F., *et al.* (2010) Mitochondrial metabolism and ROS generation are essential for Kras-mediated tumorigenicity. *Proc Natl Acad Sci U S A*, **107**, 8788-93.
157. Yang, D., *et al.* (2010) Impairment of mitochondrial respiration in mouse fibroblasts by oncogenic H-RAS(Q61L). *Cancer Biol Ther*, **9**, 122-33.
158. Moiseeva, O., *et al.* (2009) Mitochondrial dysfunction contributes to oncogene-induced senescence. *Mol Cell Biol*, **29**, 4495-507.
159. Seyfried, T.N. (2012) Respiratory insufficiency, the retrograde response, and the origin of cancer. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 177-194.
160. Guha, M., *et al.* (2013) Mitochondrial Retrograde Signaling at the crossroads of tumor bioenergetics, genetics and epigenetics. *Mitochondrion*.
161. Singh, K.K., *et al.* (2005) Inter-genomic cross talk between mitochondria and the nucleus plays an important role in tumorigenesis. *Gene*, **354**, 140-6.
162. Jazwinski, S.M. (2005) The retrograde response links metabolism with stress responses, chromatin-dependent gene activation, and genome stability in yeast aging. *Gene*, **354**, 22-7.
163. Nargund, A.M., *et al.* (2012) Mitochondrial import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science*, **337**, 587-90.
164. Al Mamun, A.A., *et al.* (2012) Identity and function of a large gene network underlying mutagenic repair of DNA breaks. *Science*, **338**, 1344-8.
165. Dang, C.V. (2010) Glutaminolysis: supplying carbon or nitrogen or both for cancer cells? *Cell Cycle*, **9**, 3884-6.
166. Bissell, M.J., *et al.* (2011) Why don't we get more cancer? A proposed role of the microenvironment in restraining cancer progression. *Nat Med*, **17**, 320-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

167. Gatenby, R.A., *et al.* (2007) Glycolysis in cancer: a potential target for therapy. *Int J Biochem Cell Biol*, **39**, 1358-66.

168. Husain, Z., *et al.* (2013) Tumor-derived lactate modifies antitumor immune response: effect on myeloid-derived suppressor cells and NK cells. *Journal of immunology*, **191**, 1486-95.

169. Liu, E.T. (2013) Grappling with cancer. *Science*, **339**, 1493.

170. Pennisi, E. (2013) Steering cancer genomics into the fast lane. *Science*, **339**, 1540-2.

171. Stratton, M.R., *et al.* (2009) The cancer genome. *Nature*, **458**, 719-24.

172. Crespi, B., *et al.* (2005) Evolutionary biology of cancer. *Trends Ecol Evol*, **20**, 545-52.

173. Merlo, L.M., *et al.* (2006) Cancer as an evolutionary and ecological process. *Nat Rev Cancer*, **6**, 924-35.

174. Davies, P.C., *et al.* (2011) Cancer tumors as Metazoa 1.0: tapping genes of ancient ancestors. *Phys Biol*, **8**, 015001.

175. Mayer, E. (1982) Evolution before Darwin. In *The Growth of Biological Thought: Diversity, Evolution, and Inheritance*. Belknap Harvard, Cambridge, MA, pp. 343-362.

176. Handel, A.E., *et al.* (2010) Is Lamarckian evolution relevant to medicine? *BMC Med Genet*, **11**, 73.

177. Koonin, E.V., *et al.* (2009) Is evolution Darwinian or/and Lamarckian? *Biol Direct*, **4**, 42.

178. Smiraglia, D.J., *et al.* (2008) A novel role for mitochondria in regulating epigenetic modification in the nucleus. *Cancer Biol Ther*, **7**, 1182-90.

179. Minocherhomji, S., *et al.* (2012) Mitochondrial regulation of epigenetics and its role in human diseases. *Epigenetics : official journal of the DNA Methylation Society*, **7**, 326-34.

180. Feinberg, A.P., *et al.* (2004) The history of cancer epigenetics. *Nat Rev Cancer*, **4**, 143-53.

181. Pawelek, J.M. (2000) Tumour cell hybridization and metastasis revisited. *Melanoma Res*, **10**, 507-14.

182. Huysentruyt, L.C., *et al.* (2010) Perspectives on the mesenchymal origin of metastatic cancer. *Cancer Metastasis Rev*, **29**, 695-707.

183. Holmgren, L., *et al.* (1999) Horizontal transfer of DNA by the uptake of apoptotic bodies. *Blood*, **93**, 3956-63.

184. Dziurzynski, K., *et al.* (2012) Consensus on the role of human cytomegalovirus in glioblastoma. *Neuro-oncology*, **14**, 246-55.

185. Detmer, S.A., *et al.* (2007) Functions and dysfunctions of mitochondrial dynamics. *Nat Rev Mol Cell Biol*, **8**, 870-9.

186. Xu, R.H., *et al.* (2005) Inhibition of glycolysis in cancer cells: a novel strategy to overcome drug resistance associated with mitochondrial respiratory defect and hypoxia. *Cancer research*, **65**, 613-621.

187. VanItallie, T.B., *et al.* (2003) Ketones: metabolism's ugly duckling. *Nutr Rev*, **61**, 327-41.

188. Drenick, E.J., *et al.* (1972) Resistance to symptomatic insulin reactions after fasting. *The Journal of clinical investigation*, **51**, 2757-62.

189. Veech, R.L. (2004) The therapeutic implications of ketone bodies: the effects of ketone bodies in pathological conditions: ketosis, ketogenic diet, redox states, insulin resistance, and mitochondrial metabolism. *Prostaglandins Leukot Essent Fatty Acids*, **70**, 309-19.

190. Krebs, H.A., *et al.* (1971) The role of ketone bodies in caloric homeostasis. *Adv. Enzyme Reg.*, **9**, 387-409.

191. Bonuccelli, G., *et al.* (2010) Ketones and lactate "fuel" tumor growth and metastasis: Evidence that epithelial cancer cells use oxidative mitochondrial metabolism. *Cell cycle*, **9**, 3506-14.
192. Seyfried, T.N. (2012) Metabolic management of cancer. In *Cancer as a Metabolic Disease: On the Origin, Management, and Prevention of Cancer*. John Wiley & Sons, Hoboken, NJ, pp. 291-354.
193. Maurer, G.D., *et al.* (2011) Differential utilization of ketone bodies by neurons and glioma cell lines: a rationale for ketogenic diet as experimental glioma therapy. *BMC cancer*, **11**, 315.
194. Skinner, R., *et al.* (2009) Ketone bodies inhibit the viability of human neuroblastoma cells. *J Pediatr Surg*, **44**, 212-6; discussion 216.
195. Seyfried, T.N., *et al.* (2003) Role of glucose and ketone bodies in the metabolic control of experimental brain cancer. *Br J Cancer*, **89**, 1375-82.
196. Jiang, Y.S., *et al.* (2013) Caloric restriction reduces edema and prolongs survival in a mouse glioma model. *Journal of neuro-oncology*, **114**, 25-32.
197. Mukherjee, P., *et al.* (2004) Antiangiogenic and proapoptotic effects of dietary restriction on experimental mouse and human brain tumors. *Clin Cancer Res*, **10**, 5622-9.
198. Mukherjee, P., *et al.* (2002) Dietary restriction reduces angiogenesis and growth in an orthotopic mouse brain tumour model. *Br J Cancer*, **86**, 1615-21.
199. Mulrooney, T.J., *et al.* (2011) Influence of Caloric Restriction on Constitutive Expression of NF-kappaB in an Experimental Mouse Astrocytoma. *PLoS One*, **6**, e18085.
200. Simone, B.A., *et al.* (2013) Selectively starving cancer cells through dietary manipulation: methods and clinical implications. *Future oncology*, **9**, 959-76.
201. Stafford, P., *et al.* (2010) The ketogenic diet reverses gene expression patterns and reduces reactive oxygen species levels when used as an adjuvant therapy for glioma. *Nutr Metab (Lond)*, **7**, 74.
202. Fine, E.J., *et al.* (2012) Targeting insulin inhibition as a metabolic therapy in advanced cancer: a pilot safety and feasibility dietary trial in 10 patients. *Nutrition*, **28**, 1028-35.
203. Zuccoli, G., *et al.* (2010) Metabolic management of glioblastoma multiforme using standard therapy together with a restricted ketogenic diet: Case Report. *Nutr Metab (Lond)*, **7**, 33.
204. Nebeling, L.C., *et al.* (1995) Effects of a ketogenic diet on tumor metabolism and nutritional status in pediatric oncology patients: two case reports. *J. Am. Coll. Nutr.*, **14**, 202-8.
205. Zhou, W., *et al.* (2007) The calorically restricted ketogenic diet, an effective alternative therapy for malignant brain cancer. *Nutr Metab (Lond)*, **4**, 5.
206. Gottschalk, S., *et al.* (2004) Imatinib (STI571)-mediated changes in glucose metabolism in human leukemia BCR-ABL-positive cells. *Clinical cancer research : an official journal of the American Association for Cancer Research*, **10**, 6661-8.
207. Zhao, Y., *et al.* (2011) Overcoming trastuzumab resistance in breast cancer by targeting dysregulated glucose metabolism. *Cancer research*, **71**, 4585-97.
208. Marsh, J., *et al.* (2008) Akt-dependent proapoptotic effects of dietary restriction on late-stage management of a phosphatase and tensin homologue/tuberous sclerosis complex 2-deficient mouse astrocytoma. *Clin Cancer Res*, **14**, 7751-62.
209. Veech, R.L., *et al.* (2001) Ketone bodies, potential therapeutic uses. *IUBMB Life*, **51**, 241-7.

210. Chen, Y., *et al.* (2009) Oxygen consumption can regulate the growth of tumors, a new perspective on the warburg effect. *PLoS One*, **4**, e7033.

211. Spitz, D.R., *et al.* (2000) Glucose deprivation-induced oxidative stress in human tumor cells. A fundamental defect in metabolism? *Ann N Y Acad Sci*, **899**, 349-62.

212. Harrison, L., *et al.* (2004) Hypoxia and anemia: factors in decreased sensitivity to radiation therapy and chemotherapy? *The oncologist*, **9 Suppl 5**, 31-40.

213. Allen, B.G., *et al.* (2013) Ketogenic diets enhance oxidative stress and radio-chemotherapy responses in lung cancer xenografts. *Clinical cancer research : an official journal of the American Association for Cancer Research*, **19**, 3905-13.

214. Abdelwahab, M.G., *et al.* (2012) The ketogenic diet is an effective adjuvant to radiation therapy for the treatment of malignant glioma. *PloS one*, **7**, e36197.

215. Seyfried, T.N., *et al.* (2010) Does the existing standard of care increase glioblastoma energy metabolism? *Lancet Oncol*, **11**, 811-3.

216. Yuneva, M. (2008) Finding an "Achilles' heel" of cancer: the role of glucose and glutamine metabolism in the survival of transformed cells. *Cell Cycle*, **7**, 2083-9.

217. DeBerardinis, R.J., *et al.* (2010) Q's next: the diverse functions of glutamine in metabolism, cell biology and cancer. *Oncogene*, **29**, 313-24.

218. Lazova, R., *et al.* (2013) A Melanoma Brain Metastasis with a Donor-Patient Hybrid Genome following Bone Marrow Transplantation: First Evidence for Fusion in Human Cancer. *PloS one*, **8**, e66731.

219. Newsholme, P. (2001) Why is L-glutamine metabolism important to cells of the immune system in health, postinjury, surgery or infection? *J Nutr*, **131**, 2515S-22S; discussion 2523S-4S.

220. Mates, J.M., *et al.* (2012) Glutaminase isoenzymes as key regulators in metabolic and oxidative stress against cancer. *Current Mol. Med.*, **12**, 1-21.

221. Shelton, L.M., *et al.* (2010) Glutamine targeting inhibits systemic metastasis in the VM-M3 murine tumor model. *Inter. J. Cancer.*, **127**, 2478-85.

222. Lim, V., *et al.* (2009) Glutamine prevents DMBA-induced squamous cell cancer. *Oral oncology*, **45**, 148-55.

223. Vincent, M.D. (2011) Cancer: beyond speciation. *Advances in cancer research*, **112**, 283-350.

224. Arens, N.C., *et al.* (2008) Press-pulse: a general theory of mass extinction? *Paleobiology*, **34**, 456-471.

225. Ko, Y.H., *et al.* (2004) Advanced cancers: eradication in all cases using 3-bromopyruvate therapy to deplete ATP. *Biochem Biophys Res Commun*, **324**, 269-75.

226. Bonnet, S., *et al.* (2007) A mitochondria-K⁺ channel axis is suppressed in cancer and its normalization promotes apoptosis and inhibits cancer growth. *Cancer Cell*, **11**, 37-51.

227. Marsh, J., *et al.* (2008) Drug/diet synergy for managing malignant astrocytoma in mice: 2-deoxy-D-glucose and the restricted ketogenic diet. *Nutr Metab (Lond)*, **5**, 33.

228. Michaelis, M., *et al.* (2009) The story of human cytomegalovirus and cancer: increasing evidence and open questions. *Neoplasia*, **11**, 1-9.

229. Yu, Y., *et al.* (2011) Viral effects on metabolism: changes in glucose and glutamine utilization during human cytomegalovirus infection. *Trends in microbiology*.

230. Bozidis, P., *et al.* (2010) Trafficking of UL37 proteins into mitochondrion-associated membranes during permissive human cytomegalovirus infection. *Journal of virology*, **84**, 7898-903.

231. Williamson, C.D., *et al.* (2009) Access of viral proteins to mitochondria via mitochondria-associated membranes. *Reviews in medical virology*, **19**, 147-64.
232. Dziurzynski, K., *et al.* (2011) Glioma-associated cytomegalovirus mediates subversion of the monocyte lineage to a tumor propagating phenotype. *Clinical cancer research : an official journal of the American Association for Cancer Research*, **17**, 4642-9.
233. Munzarova, M., *et al.* (1991) Do some malignant melanoma cells share antigens with the myeloid monocyte lineage? *Neoplasma*, **38**, 401-5.
234. Soderberg-Naucler, C., *et al.* (2013) Survival in patients with glioblastoma receiving valganciclovir. *The New England journal of medicine*, **369**, 985-6.
235. Clarke, K., *et al.* (2012) Kinetics, safety and tolerability of (R)-3-hydroxybutyl (R)-3-hydroxybutyrate in healthy adult subjects. *Regulatory toxicology and pharmacology : RTP*, **63**, 401-8.
236. D'Agostino, D.P., *et al.* (2013) Therapeutic ketosis with ketone ester delays central nervous system oxygen toxicity seizures in rats. *American journal of physiology. Regulatory, integrative and comparative physiology*, **304**, R829-36.
237. Lu, W., *et al.* (2012) Novel role of NOX in supporting aerobic glycolysis in cancer cells with mitochondrial dysfunction and as a potential target for cancer therapy. *PLoS biology*, **10**, e1001326.
238. de Groof, A.J., *et al.* (2009) Increased OXPHOS activity precedes rise in glycolytic rate in H-RasV12/E1A transformed fibroblasts that develop a Warburg phenotype. *Molecular cancer*, **8**, 54.
239. Baracca, A., *et al.* (2010) Mitochondrial Complex I decrease is responsible for bioenergetic dysfunction in K-ras transformed cells. *Biochimica et biophysica acta*, **1797**, 314-23.
240. Roskelley, R.C., *et al.* (1943) Studies in Cancer. Vii. Enzyme Deficiency in Human and Experimental Cancer. *J Clin Invest*, **22**, 743-51.
241. Seyfried, T.N., *et al.* (2008) Targeting energy metabolism in brain cancer with calorically restricted ketogenic diets. *Epilepsia*, **49 Suppl 8**, 114-6.
242. Huysentruyt, L.C., *et al.* (2008) Metastatic cancer cells with macrophage properties: evidence from a new murine tumor model. *Int J Cancer*, **123**, 73-84.
243. Poff, A.M., *et al.* (2013) The ketogenic diet and hyperbaric oxygen therapy prolong survival in mice with systemic metastatic cancer. *PloS one*, **8**, e65522.

Figure-1




Figure-2




Figure-3




Figure-4




Figure-5




Figure-6

