

Please join us for a discussion on



On behalf of Daiichi Sankyo, Inc. and AstraZeneca, we cordially invite you to join us for an informative, expert-led discussion.

This program will cover:

- An FDA-approved, HER2-directed antibody-drug conjugate for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received a prior anti-HER2-based regimen either in the metastatic setting, or in the neoadjuvant or adjuvant setting, and have developed disease recurrence during or within six months of completing therapy
- Results from a head-to-head Phase 3 clinical trial, DESTINY-Breast03
- Key safety considerations for ENHERTU
- Dosage and administration of ENHERTU

Program Details:

Saturday, September 10, 2022

6:00 PM Pacific
The Capital Grille
700 West 9th Street
Suite B
Los Angeles, CA 90015

Faculty:

Linda Buck, MSN, AOCNP

City of Hope
Orange, CA

RSVP: Noemi Sklar

Representative Program Host

nskлар@dsi.com

(917) 526-2202

By: 09/03/2022

This program is being sponsored by Daiichi Sankyo, Inc. and AstraZeneca. The speaker is being compensated for the presentation. The program is not CME accredited and may not be used for CME accreditation. In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs. Please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you may receive may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity that limits or prohibits meals from pharmaceutical companies, please identify yourself so that compliance with such requirements can be ensured.

Important Safety Information

Indication

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received a prior anti-HER2-based regimen either:

- In the metastatic setting, or
- In the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy

WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY

- **Interstitial lung disease (ILD) and pneumonitis, including fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**

Contraindications

None.

Warnings and Precautions

Interstitial Lung Disease / Pneumonitis

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU. In clinical studies, of the 491 patients with unresectable or metastatic HER2-positive breast cancer treated with ENHERTU 5.4 mg/kg, ILD occurred in 13% of patients. Fatal outcomes due to ILD and/or pneumonitis occurred in 1.4% of patients treated with ENHERTU. Median time to first onset was 5.5 months (range: 1.1 to 20.8).

Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in ≤ 28 days from date of onset, maintain dose. If resolved in > 28 days from date of onset, reduce dose one level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥ 0.5 mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥ 1 mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

Neutropenia

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU. In clinical studies, of the 491 patients with unresectable or metastatic HER2-positive breast cancer who received ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 68% of patients. Eighteen percent had Grade 3 or 4 decrease in neutrophil count. Median time to first onset of decreased neutrophil count was 22 days (range: 6 to 664). Febrile neutropenia was reported in 1.2% of patients.

Please see Important Safety Information and accompanying full Prescribing Information, including Boxed WARNINGS, and Medication Guide.



Important Safety Information (continued)

Neutropenia (continued)

Monitor complete blood counts prior to initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC] <1.0 to $0.5 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC $<0.5 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 2 or less. Reduce dose by one level. For febrile neutropenia (ANC $<1.0 \times 10^9/L$ and temperature $>38.3^\circ C$ or a sustained temperature of $\geq 38^\circ C$ for more than 1 hour), interrupt ENHERTU until resolved. Reduce dose by one level.

Left Ventricular Dysfunction

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular ejection fraction (LVEF) decrease has been observed with anti-HER2 therapies, including ENHERTU. In the 491 patients with unresectable or metastatic HER2-positive breast cancer who received ENHERTU 5.4 mg/kg, 13 cases (2.6%) of asymptomatic LVEF decrease were reported.

Assess LVEF prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. Manage LVEF decrease through treatment interruption. When LVEF is $>45\%$ and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is $<10\%$, continue treatment with ENHERTU and repeat LVEF assessment within 3 weeks. When LVEF is 40-45% and absolute decrease from baseline is 10-20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF has not recovered to within 10% from baseline, permanently discontinue ENHERTU. If LVEF recovers to within 10% from baseline, resume treatment with ENHERTU at the same dose. When LVEF is $<40\%$ or absolute decrease from baseline is $>20\%$, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of $<40\%$ or absolute decrease from baseline of $>20\%$ is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF $<50\%$ prior to initiation of treatment.

Embryo-Fetal Toxicity

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for at least 7 months following the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 4 months after the last dose of ENHERTU.

Additional Dose Modifications Thrombocytopenia

For Grade 3 thrombocytopenia (platelets <50 to $25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets $<25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less. Reduce dose by one level.

Adverse Reactions

The pooled safety population for patients with metastatic breast cancer reflects exposure to ENHERTU at 5.4 mg/kg given as an intravenous infusion once every 3 weeks (21-day cycle) in 491 patients in DESTINY-Breast03, DESTINY-Breast01, and Study DS8201-A-J101. The median duration of treatment was 13 months (range: 0.7 to 37). In this pooled safety population, the most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (78%), decreased white blood cell count (74%), decreased hemoglobin (68%), decreased neutrophil count (68%), increased aspartate aminotransferase (58%), fatigue (57%), decreased lymphocyte count (56%), vomiting (50%), decreased platelet count (49%), increased alanine aminotransferase (48%), increased blood alkaline phosphatase (45%), alopecia (41%), constipation (35%), hypokalemia (33%), decreased appetite (32%), diarrhea (31%), musculoskeletal pain (28%), increased transaminases (27%), respiratory infection (24%), headache (21%), and abdominal pain (21%).

DESTINY-Breast03

The safety of ENHERTU was evaluated in 257 patients with unresectable or metastatic HER2-positive breast cancer who received at least one dose of ENHERTU 5.4 mg/kg in DESTINY-Breast03. ENHERTU was administered by intravenous infusion once every three weeks. The median duration of treatment was 14 months (range: 0.7 to 30).

Serious adverse reactions occurred in 19% of patients receiving ENHERTU. Serious adverse reactions in $>1\%$ of patients who received ENHERTU were vomiting, interstitial lung disease, pneumonia, pyrexia, and urinary tract infection. Fatalities due to adverse reactions occurred in 0.8% of patients including COVID-19 and sudden death (one patient each).

ENHERTU was permanently discontinued in 14% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 44% of patients treated with ENHERTU. The most frequent adverse reactions ($>2\%$) associated with dose interruption were neutropenia, leukopenia, anemia, thrombocytopenia, pneumonia, nausea, fatigue, and ILD/pneumonitis. Dose reductions occurred in 21% of patients treated with ENHERTU. The most frequent adverse reactions ($>2\%$) associated with dose reduction were nausea, neutropenia, and fatigue.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (74%), decreased neutrophil count (70%), increased aspartate aminotransferase (67%), decreased hemoglobin (64%), decreased lymphocyte count (55%), increased alanine aminotransferase (53%), decreased platelet count (52%), fatigue (49%), vomiting (49%), increased blood alkaline phosphatase (49%), alopecia (37%), hypokalemia (35%), constipation (34%), musculoskeletal pain (31%), diarrhea (29%), decreased appetite (29%), respiratory infection (22%), headache (22%), abdominal pain (21%), increased blood bilirubin (20%), and stomatitis (20%).

Use in Specific Populations

- Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months following the last dose of ENHERTU.
- Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- Females and Males of Reproductive Potential:** **Pregnancy testing:** Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. **Contraception: Females:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 7 months following the last dose. **Males:** Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for at least 4 months following the last dose. **Infertility:** ENHERTU may impair male reproductive function and fertility.
- Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- Geriatric Use:** Of the 491 patients with HER2-positive breast cancer treated with ENHERTU 5.4 mg/kg, 22% were ≥ 65 years and 4% were ≥ 75 years. No overall differences in efficacy within clinical studies were observed between patients ≥ 65 years of age compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged ≥ 65 years (60%) as compared to younger patients (49%).
- Renal Impairment:** A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Monitor patients with moderate or severe renal impairment.
- Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, closely monitor for increased toxicities related to the topoisomerase inhibitor.

To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or fda.gov/medwatch.

Please see accompanying full Prescribing Information, including Boxed WARNINGS, and Medication Guide.



ENHERTU[®] is a registered trademark of Daiichi Sankyo Company, Limited.
© 2022 Daiichi Sankyo, Inc. and AstraZeneca. PP-US-ENB-0882 05/22

