



Core Mathematics C34(GCE)

Practice Question 4

Standard A[★]

Mr. S. V. Swarnaraja

(Team Leader, Marking Examiner & Author)

www. *sw* *nash*. com

**CRITICAL THINKING IS THE KEY TO SOLVE REAL WORLD PROBLEMS.
CHILDREN MUST BE TAUGHT HOW TO THINK, NOT WHAT TO THINK.
A GREAT TEACHER WILL BE CREATING STUDENTS TO DO NEW THINGS
THROUGH CRITICAL THINKING, NOT SIMPLY REPEATING WHAT OTHER
GENERATIONS HAVE DONE BEFORE. WE DO NOT NEED ANOTHER
ALBERT EINSTEIN OR ISAAC NEWTON.... WE NEED A PERSON BETTER
THAN THEM.**

MR.S.V. SWARNARAJA

Trigonometry

Question:

Find the *exact* value of

$$\sin \left[2 \cos^{-1} \left(\frac{2}{3} \right) \right]$$

(4 marks)



Golden Rules

- $\sin(A + B) = \sin A \cos B + \cos A \sin B \Rightarrow \sin(A + A) = \sin A \cos A + \cos A \sin A \Rightarrow \sin(2A) = 2\sin A \cos A$
- $\sin(A - B) = \sin A \cos B - \cos A \sin B$
- $\cos(A + B) = \cos A \cos B - \sin A \sin B \Rightarrow \cos(A + A) = \cos A \cos A - \sin A \sin A \Rightarrow \cos 2A = \cos^2 A - \sin^2 A$
 $\cos 2A = 2\cos^2 A - 1$
 $\cos 2A = 1 - 2\sin^2 A$
- $\cos(A - B) = \cos A \cos B + \sin A \sin B$
- $\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B} \Rightarrow \tan(A + A) = \frac{\tan A + \tan A}{1 - \tan A \tan A} \Rightarrow \tan(2A) = \frac{2 \tan A}{1 - \tan^2 A}$
- $\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

Traditional or Online classes

Mr. S.V. Swarnaraja

(Team Leader, Marking Examiner & Author)

Mobile: +94 777 304755

email: swa@swanash.com

www.swanash.com