

Street Drug Pharmacology 2020

J. Randall Webber, MPH, CADC, QMHP

JRW Behavioral Health Services

www.randallwebber.com

www.linkedin.com

Emerging Drugs of Abuse discussion group



Dissociative anesthetics


- Phencyclidine (PCP)
- Ketamine
- Dextromethorphan (DXM)

Dissociative anesthetics: Basics

- Addiction potential low to moderate
- Tolerance develops
- Withdrawal symptoms mild
- Moderate to high potential for immediate physical toxicity
- Potential for long-term physical toxicity varies
- Potential for acute and chronic psychiatric impairment moderate to high
- NMDA/glutamate antagonist

PCP

- Angel dust
- Sherms
- Dips
- Fry
- Embalming fluid



+500%

* The number of visits involving PCP in 2005, 2006, 2007, 2008, 2009, and 2010 is significantly different from 2011 at the .05 level.

** The number of visits involving MDMA (Ecstasy) in 2005 and 2007 is significantly different from 2011 at the .05 level.

*** The number of visits involving LSD in 2005 is significantly different from 2011 at the .05 level.

Source: 2005 to 2011 SAMHSA Drug Abuse Warning Network (DAWN).

ED visits involving PCP, LSD and MDMA: 2005-2011

PCP

- Oral (rare)
- Intranasal
- Smoked (on marijuana, mint, parsley)
- i.v. (rare)



PCP

- (Not used medically)
- Floating feeling
- Sedation
- Euphoria
- Confusion
- Disorientation
- Depersonalization
- Derealization
- Loss of ego boundaries
- Panic

PCP

- Bizarre behavior
- Aggressive behavior
- Hallucinations (rare)

PCP

- Numbness
- Ataxia
- Increased blood pressure
- Nystagmus
- Seizures

Ketamine (Ketalar)

- “Special K”
- Intravenous anesthetic
- Post-operative pain management with an opioid
- Also used in veterinary medicine
- Many of the same effects as PCP
- Fewer negative effects
- Short-acting (20-60 minutes)

Ketamine

- Addiction rare in U.S., More Common in Europe
- Heavy use →→→→→ Severe bladder infection/destruction

Ketamine: Treatment of depression

- Single sub-anesthetic dose
- Provide relief for 1-2 weeks

DXM

- Dextromethorphan
- In OTC cough medicines
- Often kept behind counter
- Robotussin DM
- Coricidin cough and cold = “Skittles”, “3-C”
- Similar to ketamine and PCP, but less intense effects except in large doses
- “Robotripping”

IMPORTANT QUESTIONS

WHAT ARE THESE DRUGS?

WHAT EFFECTS DO THEY HAVE?

WHAT ARE THE DANGERS?

WHAT IS THEIR ADDICTION POTENTIAL?

ARE THEY IN MY COMMUNITY?