

ATRAUMATIC ORTHODONTIC BANDING PROCEDURE- INDIRECT BANDING TECHNIQUE

Lakshman Chowdary Basam¹, Prasad Mandava², Vijay Kumar Chava³, Revathi Peddu⁴, Ram Chowdary Basam⁵, Sravanthi Tammineedi⁶

¹Senior lecturer, Orthodontics, Sibar Institute of Dental Sciences, Guntur, AP, India

²Head and Professor, Orthodontics, Narayana Dental College and Hospital, Nellore, AP, India

³Head and Professor, Periodontics, Narayana Dental College and Hospital, Nellore, AP, India

⁴Head and Professor, Orthodontics, Sibar Institute of Dental Sciences, Guntur, AP, India

⁵Senior lecturer, Conservative Dentistry and Endodontics, Sibar Institute of Dental Sciences, Guntur, AP, India

⁶Senior lecturer, Conservative Dentistry and Endodontics, Sibar Institute of Dental Sciences, Guntur, AP, India

ABSTRACT:

In orthodontics, banding of molars is a common procedure during treatment. Banding usually requires increased chairside time and causes fatigue to the orthodontist. Direct banding causes laceration of gingiva, extensive damage to the interdental col, which leads to bleeding, pain and discomfort to the patient. In the present article we are introducing an innovative technique showing a simple and novel laboratory procedure for indirect banding.

Keywords: Biological width, Indirect Banding.



INTRODUCTION:

Banding is the most frequently done procedure during fixed orthodontic treatment. Band construction by any direct method, including the use of preformed bands and contoured band strips has always been a time consuming and fatiguing procedure [3]. The level of anticipated pain before orthodontic treatment is higher than the real pain experienced after orthodontic procedures [1].

This simple laboratory procedure for banding, could improve chair side time, precision in banding, decrease fatigue, preventing injury to interdental col, thereby improving treatment efficiency. This technique is particularly useful in these COVID – 19 pandemic days, where there is increased risk of exposure to contagion while performing direct banding.

Here we are presenting a novel technique to band molars where orthodontist himself can do this simple technique.

Description of technique: (Figure:1-5)

Band construction by this procedure includes FOUR major steps,

i] Impression taking: This procedure was carried out by taking the single impression with irreversible hydrocolloid [alginate]. From this impression accurate working models are made with plaster of Paris.

ii] Model slicing: After drying the model, teeth adjacent to the intended tooth to be banded is sliced for about 0.5 to 1.0mm using diamond cutting disk [Figure-1]. The slicing should be done on adjacent teeth with disk positioned parallel to the long axis of the teeth intended to be banded. Care should be taken while Slicing not to extend more than 2mm beyond the gingival margin in

order to maintain the normal biological width. As average biologic width consists of the epithelial attachment (0.97 mm) plus connective tissue attachment (1.07 mm) amounting to 2.04 mm. Any violation in biologic width may alter defence mechanisms leading to inflammatory response and accelerating bone loss.^[2]

iii] Band fabrication: Band contouring and adaptation should be done accurately along with creating space for accommodation of interdental col at gingival part of band in a concave fashion [Figure-2]. Buccal tubes are welded at appropriate position on bands. Separators are placed on first appointment itself after taking first impression so that space will be available for trial in of bands in next appointment, thereby decreasing the number of appointments.

iv] Try in: Try in should be done prior to the fabrication of Transpalatal arch (TPA) [Figure-3, 4]. Cementation of TPA with accurate band adaptation is depicted in [Figure-5].

Indications for this technique:

1. All fixed orthodontic treatment procedures,

REFERENCES:

1. Ngan P, Kess B, Wilson S. Perception of discomfort by patients undergoing orthodontic treatment. Am J Orthod Dentofacial Orthop. 1989 Jul; 96(1):47-53.
2. Garguilo AW, Wentz FM, Orban B. Dimensions and relations of the

2. Uncooperative patients undergoing fixed orthodontic treatment,
3. Patients with mental disabilities,
4. Patients with limited mouth opening.

Advantages:

- 1] Easy, cost effective and simple,
- 2] Better band adaptation when compared to direct banding,
- 3] Banding of second molars for better accessibility,
- 4] Minimal trauma to gingiva compared to direct banding,
- 5] Decreases patient discomfort and fatigue,
- 6] Reduced chair side time and appointments.
- 7] This procedure is uncomplicated and involves fewer steps when compared to indirect band technique proposed by John T. Lindquist.

Disadvantages:

- 1] Technique sensitive,
- 2] Banding of adjacent teeth during the same appointment is not possible.

CONCLUSION:

This novel technique provides easy, efficient way of banding by improving access and precision and is beneficial to patient and orthodontist.

dentogingival junction in humans. J Periodontol. 1961; 32:261-7.

3. John T. Lindquist, Indirect banding technique, Angle orthodontist Vol.29, no.2. April, 1959.

FIGURES:




Figure 1: Model slicing



Figure 2: Band fabrication



Figure 3: Try-in



Figure 4: Try-in



Figure 5: Transpalatal arch (TPA) cementation