

## Laser Iridotomy Patient Information

Your eyes have been diagnosed with (or having the potential to develop) problems with the circulation of the fluid within the eye. This can lead to a very painful type of glaucoma called *angle closure glaucoma*.

Laser treatment to be performed on your eye is to help the fluid pass more freely within your eye. This is called YAG-Iridotomy.

The YAG laser is a powerful beam of light used to make small holes into the iris- the coloured part of the eye. The fluid inside your eye can then circulate more freely through this small hole. This helps to control the eye pressure. This treatment can easily be performed in the outpatients department and is carried out on a machine similar to the slit lamp used by the ophthalmologist to examine your eyes in clinic. Because this type of glaucoma can happen in both eyes, the treatment is usually performed on both eyes sequentially. It is better to prevent this type of glaucoma from happening rather than wait until you develop a painful eye outside the hospital where immediate treatment may not be available.

### On the day of your treatment:

- Do not drive yourself to the appointment.
- Your vision will be blurred for several hours after the treatment.
- Your ophthalmologist may ask you to stop existing eye drops and start new ones.
- You may need to stay at the clinic for an hour after the treatment · sometimes the eye pressure needs to be rechecked one hour after treatment.
- It is important that you do not rub your eyes for two hours after the treatment as the surface of your eye is numbed and you might inadvertently injure yourself (blotting excess tears away with the eyelid closed is possible).

### What will happen just before the laser treatment?

- The procedure will be explained to you and once you are happy to go ahead, you will be asked to sign a consent form.
- Drops are put in the eye to be treated to make the pupil smaller (Pilocarpine 2%) and to prevent a pressure rise after the procedure (lopidine stat dose). These eye drops may cause a slight headache.
- They will take about 20-30 minutes to work.
- Feel free to ask about anything you are unsure about.

Mr Madhav Vempali FRCSEd FRCOphth  
Consultant Ophthalmic Surgeon

---

- You will be seated at the laser machine.
- Anaesthetic drops are put in the eye to be treated to avoid any discomfort during the laser treatment.

### During the laser treatment

- A contact lens is placed on the front of the eye to keep the eye open and focus the laser beam.
- This may feel like 'swimming under water with open eyes'. It feels a bit uncomfortable but should not be painful.
- The laser treatment takes a few minutes. During this time, you may hear some clicking sounds and may see some bright lights. You may be aware of a pressure feeling.
- The contact lens is removed.
- An anti-inflammatory drops are prescribed. You should administer this drop four times a day for six days.

### Complications of laser treatment

These are generally uncommon:

- The pressure in the eye can rise. This may be treated with drops or tablets.
- Bleeding inside the eye from the created opening - this will clear in a few days.
- Very rare: localised cataract or adhesions between the iris and the lens - this would need a second procedure to be resolved (Cataract surgery).

**I have watched the video covering the above benefits and risks.**

**For more information please visit [www.vempali.com](http://www.vempali.com), patient information section.**

**By signing below, I acknowledge that I have read and that I understand this information**

**Signature:**

**Date:**